

ENTOMOLOGY

Heteroptera collected in Valsesia, Northern Italy

P. Dioli,¹ C.M.T. Boggio,² L. Limonta³

¹Museo Civico di Storia Naturale, Milan; ²Amateur Entomologist, Galliate (NO); ³Department of Food, Environmental and Nutritional Sciences, University of Milan, Italy

Abstract

The survey of Heteroptera carried out in 2016 along the nature path “Bosco dei Tigli” (Lime Trees Wood) in Piode (Piedmont, 900 m a.s.l.), highlighted 74 species, belonging to 68 genera in total. The number of species of each family well represents the Italian Heteroptera composition, with the prevalence of Miridae and Pentatomidae. The Miridae *Criocoris nigripes* var. *apicalis* (Fieber, 1861), a new record in the Alps, and *Dicyphus flavoviridis* (Tamanini, 1949), an Italian endemic taxon, were collected. *Atractotomus parvulus* (Reuter, 1878) and *Orthotylus viridinervis* (Kirschbaum, 1856) were recorded for the first time in Piedmont. Piode is the most Northern area, with Sondrio, where the Mediterranean Lygaeidae *Oxycaenus lavatae* (Fabricius, 1784) was found.

Introduction

The historical and nature-oriented importance of natural trails and of the surrounding habitat has received more attention recent-

ly, in order to enhance and protect the environment. Researches on biodiversity highlighted the great variety of animal and plant species that inhabit these areas (IPLA, 1989; Regione Piemonte, 2004, 2013).

Hemiptera Heteroptera presents a high number of species in Italy, more than 1500, inhabiting all trophic niches. This taxon was the object of this research as it is a good indicator of the ecosystem's health. In fact, the Heteroptera species recorded in a biotope give information on the characteristics of the environment. Samples were collected in 2016 in Valsesia along the path “Il Bosco dei Tigli”, recovered with funding from Piedmont Region and the European Community. The trail is in Piode Municipality (Vercelli Province), located 900 m a.s.l.

Materials and Methods

The species of Heteroptera were collected along the trail “Il Bosco dei Tigli” from May to November 2016.

The first sampling area (1), close to the village, was characterized by sessile oak (*Quercus petraea* (Matt) Liebl.), and linden (*Tilia cordata* Mill.). The contiguous area (2) was a meadow/pasture at 900 m a.s.l, with the prevalence of *Phyteuma betonicifolium* Vill., *Ranunculus acris* L., *Gentiana kochiana* E.P.Perrier & Songeon, *Fragaria vesca* L., *Salvia glutinosa* L., and *Primula vulgaris* Huds. The third sampling area (3) was a deciduous forest composed by *Populus tremula* L., *T. cordata*, *Q. petraea*, *Castanea sativa* Mill., *Betula pendula* Roth, *Corylus avellana* L., *Sorbus aucuparia* L., and *Laburnum alpinum* (Mill.) Bercht & J. Presl. The last sampling area (4) was close to a waterfall with *Festuca scabriculum* (Hack.) K.Richt., typical of Valsesia, *Phyteuma scheuchzeri* All., *Saxifraga cuneifolia* L., *Saponaria ocymoides* L., *Primula hirsuta* All., *Calluna vulgaris* (L.) Hull, *Cytisus scoparius* L., and *Crataegus monogyna* Jacq.

Species were identified using the main dichotomous keys of Heteroptera (Wagner & Weber, 1964; Péricart 1972, 1987, 1998; Moulet 1995) and verified by comparison with the specimen in the collection of the Museum of Natural Science in Milan (P. Dioli).

Results

The species collected in the four sampling areas along the trail are reported in Table 1.

Correspondence: Lidia Limonta, Dipartimento di Scienze per gli Alimenti la Nutrizione, l'Ambiente, Università degli Studi di Milano, Via G. Celoria 2, 20133 Milano, Italia.
E-mail: lidia.limonta@unimi.it

Key words: Hemiptera, *Orius* (*Heterorius*) *laticollis*, *Atractotomus parvulus*, *Criocoris nigripes* var. *apicalis*, *Dicyphus* (*Dicyphus*) *flavoviridis*.

Contributions: the authors contributed equally.

Conflict of interest: the authors declare no potential conflict of interest.

Funding: none.

Received for publication: 11 December 2019.

Accepted for publication: 31 December 2019.

©Copyright: the Author(s), 2019
Licensee PAGEPress, Italy
Journal of Entomological and Acarological Research 2019; 51:8752
doi:10.4081/jea.2019.8752

This article is distributed under the terms of the Creative Commons Attribution Noncommercial License (by-nc 4.0) which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

Table 1. Hemiptera Heteroptera collected in the four areas along the trail “Il Bosco dei Tigli”. The roman numerals provide the month. Corotypes (COR) according to Vigna Taglianti *et al.* (1999).

Taxa	1	2	3	4	COR
Anthocoridae					
<i>Anthocoris nemoralis</i> (Fabricius, 1794)			VII		ASE
<i>Dufouriellus ater</i> (Dufour, 1833)		VII			EUM
<i>Orius (Heterorius) laticollis</i> (Reuter, 1884)		VII	VIII		EUR
<i>Orius (Orius) niger</i> (Wolff, 1811)		VIII			TEM
Nabidae					
<i>Himacerus (Aptus) mirmicoides</i> (O. Costa, 1834)	VIII	V,VI,VII,VIII	VII,VIII, IX	IX	EUM
<i>Himacerus (Himacerus) apterus</i> (Fabricius, 1798)			VII, VIII		ASE
<i>Nabis (Nabis) rugosus</i> (Linnaeus, 1758)		VI, VII, VIII, IX	VIII	VI, VIII, IX	SIE
Miridae					
<i>Adelphocoris lineolatus</i> (Goeze, 1778)			VIII		OLA
<i>Adelphocoris seticornis</i> (Fabricius, 1775)		VIII			SIE
<i>Atractotomus parvulus</i> Reuter, 1878		VII			CEU
<i>Blepharidopterus angulatus</i> (Fallen, 1807)			VII		SIE
<i>Bryocoris pteridis</i> (Fallen, 1807)			VIII		SIE
<i>Capsus ater</i> (Linnaeus, 1758)		VI			EUM
<i>Charagochilus (Charagochilus) weberi</i> Wagner, 1953			VII		EUR
<i>Closterotomus norwegicus</i> (Gmelin, 1790)			VIII		SEM
<i>Criocoris nigripes</i> Fieber, 1861 (var. <i>apicalis</i> , Fieber, 1864)		VI			CEU
<i>Deraeocoris (Deraeocoris) ruber</i> (Linnaeus, 1758)		VII, VIII			OLA
<i>Deraeocoris (Camptobrochis) serenus</i> (Douglas & Scott, 1868)		VIII	VII, IX		TEM
<i>Dicyphus (Dicyphus) flavoviridis</i> Tamanini, 1949		V, VIII	V, IX		END
<i>Halticus apterus</i> (Linnaeus, 1758)		VII			TEM
<i>Harpocera thoracica</i> (Fallen, 1807)			VI		TEM
<i>Heterocordylus (Heterocordylus) tibialis</i> (Hahn, 1833)			V		EUR
<i>Lepidargyrus ancorifer</i> (Fieber, 1858)		VI			MED
<i>Liocoris tripustulatus</i> (Fabricius, 1781)		VII			PAL
<i>Lygus gemellatus</i> (Herrich-Schäffer, 1835)				VI	MED
<i>Monalocoris (Monalocoris) filicis</i> (Linnaeus, 1758)				VI	SIE
<i>Neolygus contaminatus</i> (Fallen, 1807)		VII			SIE
<i>Orthotylus (Orthotylus) viridinervis</i> (Kirschbaum, 1856)		VII			EUR
<i>Phylus (Phylus) coryli</i> (Linnaeus, 1758)			VIII		EUR
<i>Phytocoris (Ktenocoris) austriacus</i> Wagner, 1954			VIII	VIII	CEU
<i>Pinalitus cervinus</i> (Herrich-Schäffer, 1841)			VII		CEM
<i>Plagiognathus (Plagiognathus) arbustorum</i> (Fabricius, 1794)		VI			PAL
<i>Plagiognathus (Plagiognathus) chrysanthemi</i> (Wolff, 1804)			VI		PAL
<i>Stenodema (Stenodema) holsata</i> (Fabricius, 1787)		V, VI, VII, IX	VI		PAL
<i>Stenodema (Stenodema) laevigata</i> (Linnaeus, 1758)		V, VI, VII, VIII	V, VII, VIII	V	PAL
<i>Stenodema (Stenodema) sericans</i> (Fieber, 1861)		V	VIII, IX	VIII	EUR
<i>Stenodema (Stenodema) virens</i> (Linnaeus, 1767)		IV			OLA
<i>Stenotus binotatus</i> (Fabricius, 1794)			VIII		OLA
Tingidae					
<i>Corythucha arcuata</i> (Say, 1832)	VIII	V, VIII	V, VIII	V, VII	OLA
<i>Dictyla echii</i> (Schränk, 1782)			VIII		PAL
<i>Physatocheila dumetorum</i> (Herrich-Schäffer, 1838)			VII		CEM
Alydidae					
<i>Alydus calcaratus</i> (Linnaeus, 1758)			VIII		OLA
Coreidae					
<i>Ceraleptus gracilicornis</i> (Herrich-Schäffer, 1835)		VI			TEM
<i>Coreus marginatus</i> (Linnaeus, 1758)		IX			ASE

To be continued on next page

The survey showed differences among the four areas sampled, underlining the ecological characteristics of the *Heteroptera* species collected.

The first area, the closest to the village, was characterized by species widespread in Italy, namely *Himacerus (Aptus) mirmicoides* (O. Costa, 1834), *Aphanus rolandri* (Linnaeus, 1758), and *Pyrrhocoris apterus* (Linnaeus, 1758). Also, the Nearctic *Corythucha arcuata* (Say, 1832), since its introduction in Europe, is common on oak in North Italy, while *Tritomegas rotundipennis* (Dohrn, 1862) is recorded locally in Italy, Spain, France, Germany, Austria, and Slovenia.

Meadows and pastures of area 2 presented the highest number

of species (Figure 1). The abundance of plant species favored the presence of *Pentatomidae* and *Scutelleridae*, mainly collected in this part of the trail. Particularly noteworthy were *Myrmus miriformis* (Fallen, 1807), *Lygaeosoma sardeum* Spinola, 1837, and *Macroplax preyssleri* (Fieber, 1837), typical of dry meadows. A predator endemic species, *Dicyphus (Dicyphus) flavoviridis* Tamanini 1949 was collected on *Salvia glutinosa* L. in May and August.

In the undergrowth of the third sampling area, *A. mirmicoides*, *Oxycarenus (Oxycarenus) lavaterae* (Fabricius, 1787), and *Palomena prasina* (Linnaeus, 1761) were collected on *Cytisus scoparius* L. and *Avenella flexuosa* (L.) Drejer. *Drymus (Sylvadrymus)*

Table 1. Continued from previous page.

<i>Coriomeris denticulatus</i> (Scopoli, 1763)		IX			SEM
<i>Syromastus rhombeus</i> (Linnaeus, 1767)		VI			TEM
Rhopalidae					
<i>Corizus hyoscyami</i> (Linnaeus, 1758)			VI		ASE
<i>Myrmus miriformis</i> (Fallen, 1807)		VIII			ASE
<i>Rhopalus (Rhopalus) subrufus</i> (Gmelin, 1790)			VII		COS
<i>Stictopleurus punctatonevus</i> (Goeze, 1778)		VIII			EUR
Lygaeidae					
<i>Aphanus rolandri</i> (Linnaeus, 1758)	VII				TUM
<i>Drymus (Sylvadrymus) ryeii</i> Douglas & Scott, 1865			VI		EUR
<i>Kleidocerys resedae</i> (Panzer, 1797)		VIII	V, VII, VIII		SIE
<i>Lygaeosoma sardeum</i> Spinola, 1837		VIII			TEM
<i>Macroplax preyssleri</i> (Fieber, 1837)		VII			EUR
<i>Oxycarenus (Oxycarenus) lavaterae</i> (Fabricius, 1787)	VI, VII, VIII		VII	VII	MED
<i>Rhyparochromus pini</i> (Linnaeus, 1758)		VIII			ASE
<i>Scolopostethus thomsoni</i> Reuter, 1875		VII			OLA
<i>Spilostethus saxatilis</i> (Scopoli, 1763)			VI		TEM
Acanthosomatidae					
<i>Elasmucha grisea</i> (Linnaeus, 1758)			VI		SIE
Cydnidae					
<i>Tritomegas rotundipennis</i> (Dohrn, 1862)	VIII				EUM
Pentatomidae					
<i>Aelia acuminata</i> (Linnaeus, 1758)				V	SEM
<i>Carpocoris (Carpocoris) purpureipennis</i> (De Geer, 1773)		VI			EUR
<i>Dolycoris baccarum</i> (Linnaeus, 1758)		VI			EUR
<i>Eurydema (Eurydema) oleracea</i> (Linnaeus, 1758)		VI			PAL
<i>Eysarcoris ventralis</i> (Westwood, 1837)		VIII			PAL
<i>Graphosoma italicum italicum</i> (O.F. Müller, 1766)		VI			EUM
<i>Palomena prasina</i> (Linnaeus, 1761)	VI, VII, VIII		IV, V, VII, VIII, IX	VII, VIII	SEM
<i>Pentatoma (Pentatoma) rufipes</i> (Linnaeus, 1758)			VI		ASE
<i>Picromerus bidens</i> (Linnaeus, 1758)				IX	OLA
<i>Piezodorus lituratus</i> (Fabricius, 1794)			VIII		TEM
<i>Staria lunata</i> (Hahn, 1835)		VIII			TUM
Scutelleridae					
<i>Eurygaster testudinaria</i> (Geoffroy, 1785)		VI			PAL
Pyrrhocoridae					
<i>Pyrrhocoris apterus</i> (Linnaeus, 1758)	IV, V, VII, VIII	V, VI, VIII	IV, V, VII	IV, VII	OLA

ryeii Douglas & Scott, 1865 and *Stenodema (Stenodema) laevigata* (Linnaeus, 1758) were characteristic of the wet undergrowth. *Elasmucha grisea* (Linnaeus, 1758) in June and *Kleidocerys resedae* (Panzer, 1797) from May to August were recorded on birch and heather. *Corythucha arcuata* (Say, 1832) was the predominant species on the crown of the tree, together with *Himacerus (Himacerus) apterus* (Fabricius, 1798), an important predator of microarthropods.

Around the waterfall in the fourth sampling area two predators were collected, *Nabis rugosus* (Linnaeus, 1758), on Ericaceae, and *Picromerus bidens* (Linnaeus, 1758), which preys on moths' larvae.

Pyrrhocoris apterus (Linnaeus, 1758) was recorded in foliar debris in all the sampling area.

Family Miridae was the most represented, including 42% of the species collected, which was 31 species out of 74 Heteroptera species. Pentatomidae and Lygeidae species constituted 15% and 12% respectively. Considering the diet of the species, 77% were phytophagous, 13% predatory, 7% omnivorous, and 3% granivorous.

The following species are worth mentioning for their significance in the considered area.

Orius laticollis (Reuter, 1884). Species recorded in Italy in the Northern and Southern Regions and in Sicily, in the wet areas and the edge of streams, where it predaes mainly on *Salix* and, to a lesser extent on *Alnus* Mill., *Populus* L., *Ulmus* L., *Quercus* (L.) and *Sorbus* L. (Pericart 1972).

Atractotomus parvulus (Reuter, 1878). The first record in Piedmont, it was previously generically cited for the Northern regions of Italy (Kerzhner & Josifov 1999). Monovoltine, it occurs on conifers (*Pinus* L., *Abies* Mill.), and overwinters as an egg (Wagner & Weber 1964). This species is very close to *Atractotomus marcoi* (Carapezza, 1982) recorded in Sicily and Russia (Carapezza, 1982), recognizable by the smaller size, the shorter second antennal segment, and the structure of the male genitalia, especially the vesica. The female of *A. parvulus* from Valsesia has the second antennal segment longer than the width of the head, more strongly inflated and fusiform. Moreover, the basal part of exocorium is clear. The distribution of these two species must be better clarified on the basis of a series of specimens, as they may have been confused one with the other in the past (Stonedahl, 1990).




Figure 1. Number of species of Heteroptera collected in the four sampling areas along the natural path in Valsesia (NO).

Criocoris nigripes var. *apicalis* Fieber, 1864. The variety and the nominal form were recorded in Friuli, Trentino (Servadei 1967), and Piedmont (Dioli 1980). The specimen collected in Valsesia was the first in the Alpine area of Piedmont. In fact, previously, one adult was recorded in spring in Rovasenda (Novara) moorland, therefore two generations per year are assumed (Dioli, 1980). The species overwinters as an egg, and it colonizes *Galium* L. Adults are active in June-July (Wagner & Weber, 1964). *C. nigripes* var. *apicalis* is a new combination, previously associated with *C. crassicornis* (Hahn, 1834).

Orthotylus viridinervis (Kirschbaum, 1856). Sporadically recorded in Northern and Southern Italy, this is the first report in Piedmont. The species overwinters as an egg, thrives on *Fraxinus ornus* L., seldom on other plants. Adults are active from July to August (Wagner & Weber 1964).

Corythucha arcuata (Say, 1832). Nearctic species, recorded in the Alps and in the plain in Northern Italy, since 2000 (Bernardinelli and Zandigiacomo, 2000), later in Switzerland, Bulgaria, Turkey, and Iran. Spontaneous and planted oaks host this species (Dioli *et al.*, 2007).

Drymus ryeii (Douglas & Scott, 1865). Infrequent in Italy, recorded North of Po River in broadleaves residues and under stones (Dioli, 1974).

Oxycarenus lavaterae (Fabricius, 1787). Mediterranean species, the presence in the alpine and sub-alpine areas is considered occasional. Overwintering adults caused alarm several times as they aggregate on the bark of broadleaved plants (Capra 1961; Perini & Tamanini, 1961; Tamanini 1961; Ciampolini & Trematerra, 1986-1987).

Conclusions

In the survey, carried out along the nature path “Bosco dei Tigli” (Lime Tree Wood) at 900 m a.s.l. in Valsesia (Piedmont), 74 species of Heteroptera, belonging to 68 genera and 13 families, were collected in total. The number of species for each family represented the composition of the Italian Heteroptera, with the predominance of Miridae and Pentatomidae.

The chorology showed that 50% of the species collected are Euro-Asiatic *sensu lato*. Palearctic and European species constituted 25% and 17%, respectively. Cosmopolitan species were insignificant, 1.1% and only one endemic species was recorded, *Dicyphus (Dicyphus) flavoviridis*.

The ubiquitous or pest species *Dolycoris baccarum* (L.), *Stenodema (Stenodema) sericans* (Fieber), *S. (S.) virens* (L.), *S. (S.) laevigata* (L.), and *Coriomeris denticulatus* (Scopoli) were mainly detected in the wide meadow areas, once cultivated, characterized by Poaceae, Fabaceae, and Asteraceae. The presence of phytophagous species was balanced with predator species.

The only allochthonous species *Corythucha arcuata* (Say), incidentally introduced in Italy in 2000 (Bernardinelli and Zandigiacomo, 2000), was collected in all the sampling areas.

In the area considered, of particular note were the first record in Piedmont of *Atractotomus parvulus* (Reuter), the second record in Piedmont and the first in the Alpine area of *Criocoris nigripes* (Fieber), the new in Piedmont and second record in northern Italy of *Orthotylus viridinervis* (Kirschbaum), and the record of *Oxycarenus lavaterae* (F.) as Piode is the most norther-

ly locality in Italy, together with Sondrio, where the species was recently collected also at considerable height (M. Salvetti communication).

The species collected were mainly typical of the Alps, but Mediterranean species were also collected in the sunny area, in the meadows or on *Cytisus scoparius* L., while European and Siberian species were common in linden, hazelnut, oak, and chestnut sciophilous forest.

References

- BERNARDINELLI I., ZANDIGIACOMO P., 2000 - Prima segnalazione di *Corythucha arcuata* (Say) (Heteroptera, Tingidae) in Europa. - Inf. Fitopat. 50: 47-49.
- CAPRA F., 1961 - Note su ammassamenti di *Oxycarenus lavaterae* (F.) nel Polesine (Hemiptera, Heteroptera, Lygaeidae). - Studi Trentini Sci. Nat., XXXVIII: 182-187.
- CARAPEZZA, A., 1982 - Il genere *Atractotomus* in Sicilia e in Calabria (Heteroptera, Miridae). - Naturalista Siciliano 6: 41-48.
- CIAMPOLINI M., TREMATERRA P., 1986/1987 - Biological studies on *Oxycarenus lavaterae* (F.) (Rhynchota Heteroptera Lygaeidae). - Boll. Zool. Agr. Bachic. II 19: 187-197.
- DIOLI P., 1974 - Emitteri Eterotteri nuovi o poco noti della Valtellina (Hemiptera, Heteroptera). - Mem. Soc. Entom. Ital., Genova 53: 30-38.
- DIOLI P., 1980 - Eterotteri della Brughiera di Rovasenda (Piemonte). Quaderni sulla "Struttura delle zoocenosi terrestri". 1. La Brughiera pedemontana. - Roma. Collana del Programma finalizzato "Promozione della qualità dell'ambiente" del C.N.R.: 35-56.
- DIOLI P., GIACALONE FORINI I., MORETTI M., SALVETTI M., 2007 - Note sulla distribuzione di *Corythucha arcuata* (Insecta, Heteroptera, Tingidae) in Cantone Ticino (Svizzera), Valtellina e alto Lario (Lombardia, Italia). - Il Naturalista Valtellinese - Atti museo civ. Storia naturale Morbegno 18:59-68.
- IPLA, 1985 - Piano Naturalistico del Parco Naturale Alta Valsesia - Regione Piemonte.
- KERZNER I. M., JOSIFOV MOULET M., 1999 - Cimicomorpha II. Miridae. In: AUKEMA B. & RIEGER, C. (Eds.) Catalogue of the Heteroptera of the Palaearctic Region Vol. 3. Netherlands Entomological Society, Amsterdam.
- MOULET P., 1995 - Hemiptères Coreoidea (Coreidae, Rhopalidae, Alydidae), Pyrrhocoridae, Stenocephalidae euro-mediterranéens. - Faune de France 81: 336 pp.
- PERICART J., 1972 - Hémiptères Anthocoridae, Cimicidae, Microphysidae de l'Ouest-Paléarctique. Faune de l'Europe et du Bassin Méditerrané Vol.7: IV + 402 pp. Masson et C.ie, Paris.
- PERINI T., TAMANINI L., 1961 - Osservazioni sulla comparsa in massa di *Oxycarenus lavaterae* (F.) (Hemiptera, Heteroptera, Lygaeidae). - Studi Trentini Sc. Nat. XXXVIII: 57-66.
- REGIONE PIEMONTE 2004 - Indagini e studi finalizzati alla predisposizione del piano di tutela delle acque. D. Lgs. 152/99.
- REGIONE PIEMONTE (provincia di Vercelli, Parco Naturale alta Valsesia), 2013 - Carta degli habitat forestali di interesse comunitario nei SIC IT1120028 "Alta Valsesia" e IT1120006 "Val Mastallone".
- SERVADEI, A., 1967 - Fauna d'Italia: Rhyncota, Heteroptera e Homoptera Auchenorrhyncha. Catalogo topografico. Bologna, Calderini, 1967.
- STONEDAHL, G.M., 1990 - Revision and cladistic analysis of the Holarctic genus *Atractotomus* Fieber (Heteroptera: Miridae: Phylinae). - Bull. Am. Mus. Nat. Hist. 198: 1-88.
- TAMANINI L., 1961 - Alcune osservazioni sulla biologia dell'*Oxycarenus lavaterae* (Fabr.) (Heteroptera, Lygaeidae). - Mem. Soc. Ent. Ital. Genova, 40: 141-143.
- VIGNA TAGLIANTI A., AUDISIO A. P., BIONDI M., BOLOGNA M.A., CARPANETO G.M., DE BIASE A., FATTORINI S., PIATTELLA E., SINDACO R., VENCHI A., ZAPPAROLI M., 1999 - A proposal for a chorotype classification of the Near East fauna, in the framework of the Western Palearctic region. - Biogeographia J. Integr. Biogeogr. 20: 31-59.
- WAGNER E., WEBER H.H., 1964 - Hétéroptères Miridae. - Faune de France 67: 1-592.