



Aging hampers neutrophil extracellular traps (NETs) efficacy

Maurizio Sabbatini¹ · Elisa Bona² · Giorgia Novello² · Mario Migliario³ · Filippo Renò³ 

Received: 7 October 2021 / Accepted: 17 July 2022
© The Author(s) 2022

Abstract

Background NETosis is a neutrophil-mediated defense mechanism during which DNA and enzymes are extruded forming a network (NETs) trapping and killing different pathogens. NETosis is reduced in both mice and humans during aging.

Aims We explored the difference in the efficacy of NETs released in elderly (> 65 years) versus adults (20–50 years) subjects in inhibiting *Staphylococcus aureus* growth and activating the growth of keratinocytes.

Methods Neutrophil granulocytes, obtained from venous blood both in healthy elderly and adult subjects, were stimulated by LPS (0–250 µg/ml) to induce the formation of NET. NETs were quantified by SYBR Green staining and growth inhibition of *S. aureus* was evaluated by disk diffusion test. Furthermore, NETs (0–500 ng/ml) were added to immortalized human keratinocytes (HaCaT cells), and their proliferation was evaluated by MTT assay after 24 h. Finally, the DNA size of NETs was evaluated by flow cytometry after SYBR Green staining.

Results Greater production of NETs was observed in elderly subjects than in adults, but these NETs showed reduced bactericidal capacity and HaCaT cells' proliferation stimulation. The activities of the NETs are related to the size of the extruded DNA threads, and when NETs size was analyzed, DNA from elderly showed a higher size compared to that obtained by adults.

Discussion Unexpected results showed aging-related NETs structural modification resulting in both a lower antimicrobial activity and keratinocyte proliferation stimulation compared to NETs obtained from adults.

Conclusions The NETs DNA size observed in elderly subjects has not been previously reported and could be part of other pathogenic mechanisms observed in aging.

Keywords Aging · NETosis · LPS · HaCaT · DNA fragmentation

Introduction

Neutrophil granulocytes play a crucial role in the innate immune defense against bacteria, fungi and virus and they are potentially harmful to the host as well.

Their activation and microbicidal activities are strictly controlled by a plethora of stimuli, and recent evidence

suggests that they are quite versatile and can perform previously unsuspected functions, such as reverse transmigration, crosstalk and regulation of other leukocyte populations [1].

The antimicrobial and cytotoxic action mechanisms of neutrophils consist in phagocytosis, in the generation of reactive oxygen species (ROS), and in the degranulation of several microbicidal factors (α -defensins, cathelicidin, elastase, cathepsin G and lactoferrin).

An additional antimicrobial action of neutrophil granulocytes was reported by Brinkmann and colleagues [2] who observed the extrusion by neutrophils of a meshwork of chromatin fibers decorated with granule-derived antimicrobial peptides, and enzymes capable to kill Gram-positive and Gram-negative bacteria. This defensive meshwork has received the denomination of neutrophil extracellular traps (NETs). NETs are composed of highly decondensed chromatin fibers having diameter from 15 to 17 nm derived from nuclear components accompanied by histone proteins

✉ Filippo Renò
filippo.reno@med.uniupo.it

¹ Department of Sciences and Innovative Technology, Università del Piemonte Orientale, Viale T. Michel 11, 15121 Alessandria, Italy

² Department of Sciences and Innovative Technology, Università del Piemonte Orientale, Piazza San Eusebio 5, 13100 Vercelli, Italy

³ Department of Health Sciences and Innovative Research Laboratory for Wound Healing, Università del Piemonte Orientale, via Solaroli, 17, 28100 Novara, Italy

complexed with microbicidal globular proteins, such as elastase, cathepsin G, myeloperoxidase, normally stored in neutrophil granules [2, 3].

NETs are released into the extracellular space where the chromatin meshwork traps microbes limiting their diffusion and collecting the neutrophil factors, thus increasing the microbicidal effects [4]. NETs release becomes fundamental as a defense mechanism when the size of the pathogens makes phagocytosis an unreliable process [5].

A further ability with which NETs carry out their microbicidal activity is related to the ability of DNA to induce the chelation of manganese and other ions. As a consequence of the chelating activity of the DNA, present in the NETs meshwork, the altered transport of the ions prevents the survival of microbes [6, 7]. Furthermore, it has been observed that NETs do not have an active function in eliminating pathogens only, but also, they regulate the local inflammatory process [8, 9]. Furthermore, the importance of NETs structure in fighting microbial invasion has been demonstrated in mice infected with pathogens which showed greater bacterial dissemination when treated with exogenous DNases [5, 10].

In summary, NETs act as a complex defence system using DNA threads, microbicidal enzymes trapped in the DNA mesh and human complement system [11] and by enhancing the activity of bystander T cells [12]. Furthermore, NETs help the tissue regeneration process, particularly in wound healing, by stimulating keratinocytes proliferation [13].

This complex modularity of NETs explains why their qualitative or quantitative alteration could cause pathologies or aggravate organ damage in several pathologies [14].

During aging, immune performance is reduced and neutrophils activity is also impaired, increasing susceptibility to microbial invasion [15]. In particular, in elderly mice NETs production induced by both PMA and *S. aureus* was strongly reduced [16], as well as in human neutrophils from elderly subjects primed with TNF- α and stimulated with LPS [17]. However, the production of NETs in healthy elderly human subjects has not been extensively studied, especially in non-preconditioned cells.

Likewise, no information about the length of DNA present in NETs was produced; on the other hand, particular attention was paid to the dimension and structure of NET meshwork [2]. LPS stimulation activates the TLR4-JNK pathway that is critical in determining the NETotic fate of neutrophils in response to bacterial invasion [18]. Furthermore, TLR4 is a cell membrane receptor widely present on the cell that has been shown to be involved in wound healing by affecting keratinocytes' migration [19].

In this paper, to clarify the effect of aging on NET activity, we measured NETs production in neutrophils obtained from adult and elderly subjects after physiological stimulation with LPS and tested their antimicrobial capacity

against *S. aureus* and their healing ability in stimulating the growth of human keratinocytes.

Materials and methods

Granulocyte isolation

Human blood from healthy volunteer donors was used with prior informed consent and after receiving the approval of the Hospital "Maggiore della Carità" Ethic Committee (C.E. 61/10).

Donors were divided into two groups named adult (10 subjects, range 20–50 years, mean \pm S.D. = 35 ± 10.9) and the elderly (8 subjects over 65 years of age, range 71–75, mean \pm S.D. = 71.8 ± 1.9).

Venous blood samples were collected in Vacutainer tubes with lithium-heparin anticoagulants and were processed to separate blood to its various cellular components (erythrocytes, leukocytes).

Initially, the blood was diluted 1:2 with PBS and subsequently stratified on Lymphocyte (PromoCell, Heidelberg, Germany) previously dispensed in falcon tubes.

The three upper phases were discarded, whereas the part with erythrocytes and granulocytes was added with ammonium chloride solution (150 mM NH_4Cl , 10 mM NaHCO_3 , 1 mM EDTA, pH = 7.4) and conserved in ice for 20 min to isolate granulocytes from erythrocytes.

Once the sample was lysed, centrifugation at 800 rpm ($120 \times g$) was performed for 5 min to obtain a pellet at the bottom of the tube containing granulocytes.

At the end, they were suspended in RPMI at 10% of FBS (Euroclone, Milan, Italy) at 10^6 cells/ml concentration.

NETs fluorescence microscopy visualization

NETosis is identified by the production of neutrophil extracellular traps (NETs) formed by a network of DNA and mainly histone proteins. To assess the presence of NETs using fluorescence microscopy visualization, 1×10^5 neutrophils obtained from the two experimental groups, were seeded on 24 multi-wells and treated with LPS 250 $\mu\text{g}/\text{ml}$ for 4 h to induce NETs production (positive control). At the end of the experiments, cells were fixed overnight at 4 °C in the dark using a 3.7% formaldehyde/3% sucrose solution in PBS and stained with SYBR Green dye (10 $\mu\text{g}/\text{ml}$) (Thermo Fisher Scientific, Waltham, Massachusetts, USA). Representative images were then acquired using a fluorescence microscope (Leica, Italy), and analyzed with Image J software at an original magnification of $\times 40$.

NETs quantification

NETs were produced according to the literature procedure [20]. Briefly, 1×10^5 neutrophils/well were seeded into a 96 multi-well culture plate in Hanks' Balanced Salt solution (HBSS) and stimulated with different LPS concentrations (0–250 $\mu\text{g/ml}$, to activate NETs formation), Triton X-100 (to evaluate total DNA content) or left untreated (negative control and DNase digested samples). After 2 h, 5 U DNase was added to all wells (excluding controls and Triton X-100 treated samples) and 45 min after DNase addition, cell-impermeable fluorescent DNA dye Sytox Green (Thermo Fisher Scientific, Waltham, MA, USA) was added to each well and the plate was incubated for further 15 min. At the end of the incubation, Sytox Green fluorescence ($\lambda_{\text{ex}}504$ nm/ $\lambda_{\text{em}}523$ nm) was quantified using a microplate reader (Victor X4, Perkin-Elmer, Waltham, MA, USA). Extracellular DNA content was expressed as percentage of total DNA obtained by subtracting the fluorescence intensity of the DNase containing wells from the comparative control and dividing the obtained result by the fluorescence intensity of Triton X-100 containing wells as indicated by the following formula:

$$\% \text{extracellular DNA} = \frac{\text{DNA stimulated with LPS} - \text{digested DNA by DNase}}{\text{DNA treated with Triton} \times 100}.$$

Disk diffusion assay

The effect of NETs against bacteria was tested according to the "EUCAST Disk diffusion method 6.0". A disk *Diffusion Assay* was performed against *Staphylococcus aureus* NCTC 6571. The bacterial samples were plated in the Muller Hinton Agar medium and paper disks (6.0 mm diameter) soaked with 10 μl of NETs were placed on the plates and incubated at 37 °C for 24 h. NETs concentrations were normalized at 2 $\mu\text{g/ml}$. Inhibition halos diameter was measured in millimeter using a caliber.

Keratinocyte proliferation

Both NETs isolated from adult and elderly volunteers were tested on human spontaneously immortalized keratinocyte (HaCaT) proliferation.

Neutrophils obtained by adult and elderly volunteers were seeded in 60 mm Petri dishes, and stimulated with 250 $\mu\text{g/ml}$ LPS (Sigma Aldrich, Saint Louis, MO, USA). The neutrophils were incubated for 4 h in a humidified atmosphere containing 5% CO_2 at 37 °C, then supernatant containing

NETs was collected. NETs concentration was measured as previously reported [13].

HaCaT was incubated in cell culture medium without phenol red (Euroclone, Milan, Italy) to avoid colorimetric interference. HaCaT cells (10×10^3 cells/well) were seeded into a 96 multi-well plate, treated with increasing NETs concentrations (0–500 ng/ml of DNA) and incubated for 48 h in a humidified atmosphere containing 5% CO_2 at 37 °C. Upon the incubation, 0.5 mg/ml MTT (Thermo Fisher Scientific, Waltham, MA, USA) was added to wells for 3 h at 37 °C to allow formazan salts' precipitation. The resulting insoluble purple precipitate was then dissolved in DMSO (Carlo Erba Reagents, Cornaredo, Italy) and the absorbance was read at 570 nm using a microplate reader (Victor X4, Perkin-Elmer, Waltham, MA, USA).

NETs analysis by flow cytometry

To evaluate whether the different activities of NETs isolated from adult and elderly volunteers were due to their structural difference, NETs fragmentation was assessed by flow cytometry using a modification of the method of Delobel and Tesnière [21].

NETs obtained both from adult and elderly subjects (5 μg) were diluted in 2 ml PBS, incubated for 30 min at 37 °C with SYBR Green dye (10 $\mu\text{g/ml}$) and then analyzed using the flow cytometer (FACSCALIBUR, BD bioscience, San Jose, CA) equipped with a 15 mW air-cooled argon ion laser operating at 488 nm. Green fluorescence was collected through a 525 nm BP filter and fluorescence data were displayed on a four-decade logarithmic scale. A minimum of 10,000 events per sample were collected at a low sample flow rate setting (12 $\mu\text{l/min}$) to improve the coefficient of variation on the DNA histograms. Data were collected using the CellQuest software (BD Bioscience, San Jose, CA, USA), and then, analyzed using Flowing Software 2.5.2 (University of Turku, Finland).

Statistical analysis

Statistical analysis was performed with the GraphPad statistical software (GraphPad Software Inc., CA, USA). The statistical significance of the data obtained was evaluated using the ANOVA-Bonferroni test. The p values < 0.05 was considered statistically significant.

Results

LPS induced a high NETs release in elderly subjects compared to adult ones

Neutrophils extracted from two different groups of healthy volunteers (adult and elderly) were stimulated with LPS.


Stimulation was performed using LPS (0–250 $\mu\text{g}/\text{ml}$) for 4 h (Fig. 1A). Both groups produced NETs in a dose-dependent manner, but elderly subjects generated a larger quantity of NETs than adults (Fig. 1A, B). This difference in the amount of NETs between the two groups became statistically significant starting from an LPS concentration of 50 $\mu\text{g}/\text{ml}$, when the cells from elderly volunteers released an amount of extracellular DNA equal to $30 \pm 9\%$ of total DNA present in the cell, against the $9 \pm 4\%$ produced by neutrophils from adult subjects. Furthermore, using 100 $\mu\text{g}/\text{ml}$ LPS, NETs produced by the elderly were $50 \pm 4.5\%$ compared to $25\% \pm 0.9\%$ of adults. Finally, stimulation with 250 $\mu\text{g}/\text{ml}$ LPS induced the release of $95 \pm 3\%$ of the total DNA of elderly volunteers, which represents almost all the cellular DNA, whereas the adults released

only the $33 \pm 0.7\%$ of the total DNA (Fig. 1A). Differences in 250 $\mu\text{g}/\text{ml}$ LPS-induced NETs production in adult and elderly subjects were also evident in samples stained with SYBR Green dye and observed in fluorescence microscopy (Fig. 1B).

NETs produced by neutrophils from elderly people displayed a lower antimicrobial activity compared to adult subjects

NETs antimicrobial activity was tested using a disk diffusion assay performed against *Staphylococcus aureus*. To test the effective microbicidal action of NETs, produced by both adult and elderly subjects, NETs concentrations were normalized at 2 $\mu\text{g}/\text{ml}$. Disks soaked using NETs

Fig. 1 NETs released from human granulocytes after LPS stimulation. **A** Extracellular DNA quantification. LPS (0–250 $\mu\text{g}/\text{ml}$) was used to stimulate NETs production in 1×10^5 neutrophils per well into a 96 multi-well culture plate. White bars = adult subjects, gray bars = elderly subjects *** $p < 0.001$ compared to adult; **B** representative fluorescence images of LPS-induced NETs production obtained after 10 $\mu\text{g}/\text{ml}$ SYBR Green staining. Original magnification 40 \times . Ct = unstimulated granulocytes; LPS = granulocytes stimulated with 250 $\mu\text{g}/\text{ml}$ LPS



from adults created an inhibition halo with a diameter equal to 10.33 ± 0.48 mm; while those soaked with NETs obtained from elderly people created a halo with a significantly smaller diameter (6.33 ± 0.14 mm $p < 0.05$) (Fig. 2A, B).

HaCaT cells proliferate when stimulated only by NETs isolated from adult volunteers

NETs have been proved to induce proliferation of human keratinocytes via Toll-Like Receptor 9 (TLR 9) and NF- κ B pathway activation [13]. To test the influence of NETs on wound healing process during aging, an MTT proliferation test on spontaneously immortalized human keratinocytes (HaCaT) was performed using different concentrations of NETs obtained from adult and elderly subjects.

In the adult group, the proliferation of HaCaT cells was stimulated by NETs in a concentration range of 0.1–5 ng/ml; then, the proliferation slowed down from a concentration of 10 ng/ml, while over 100 ng/ml, a cytotoxic effect became evident (Fig. 3). On the contrary, in elderly group, no proliferative effect was observed at any NETs concentration (Fig. 3).

Flow cytometric analysis showed a different fragmentation pattern in NETs from adult and elderly subjects

A simple flow cytometric DNA fragmentation analysis was performed to assess whether the different antimicrobial and proliferative abilities observed in the NETs from adults and the elderly were due to a structural modification of the NETs. NETs were stained with SYBR Green and the green




Fig. 3 Human keratinocytes proliferation induced by NETs. MTT assay was to evaluate cell proliferation. MTT was performed on HaCaT cells stimulated with NETs (0–500 ng/ml) obtained by granulocytes from adult (continuous line) and elderly (dashed line) for 48 h. *** $p < 0.001$ referred to elderly subject

fluorescence obtained from the DNA-linked dye was read using a logarithmic scale (Fig. 4A). The DNA of NETs from adults appeared mainly distributed along a Gaussian-like curve with a low fluorescence intensity (Fig. 4A, Region 1), while the DNA of the elderly was distributed over two different populations: a “low intensity” population, formed by small sized DNA (Region 1, R1), and a “high intensity” population consisted of large DNA less fragmented (Region 2, R1). Quantitative analysis of the DNA distribution in R1 and R2 (Fig. 4B) showed that the $43.0 \pm 4.1\%$ of DNA from NETs obtained from elderly subjects was in R2 compared to the $19.3 \pm 2.6\%$ of the NETs from adult. Therefore, a real shift in the DNA size of the NETs during aging was observed (Fig. 4A).





Fig. 2 Evaluation of antimicrobial activity of NETs against *S. aureus*. **A** Representative images of inhibition growth halo induced in *S. aureus* by the NETs from adult and elderly subjects. **B** Quantification of halo dimensions induced by NETs from adult and elderly groups. *** $p < 0.001$




Fig. 4 Flow cytometric analysis of NETs DNA size. **A** Representative fluorescence histograms relative to DNA presents in NETs obtained from adult and elderly subjects. DNA was staining with SYBR Green dye (10 $\mu\text{g/ml}$) and green fluorescence obtained was displayed in a logarithmic scale. Region R1 was created to include the majority (more than 70%) of fluorescence obtained in adult samples; in samples from elderly subjects, DNA showed a different fluorescence dis-

tribution with a population showing higher fluorescence and, therefore, higher size (collected in Region R2) compared to adult samples. To stress the shift in DNA size during the aging, an overlapped histogram was added (adult/elderly). **B** Quantification of fluorescence events distribution between R1 and R2 in adult and elderly groups expressed as % of total events. *** $p < 0.001$ referred to R1

Discussion

During aging, the human immune system undergoes a process of decline in its protective function, making our body more susceptible to infections [22, 23]. This decline affects also neutrophils activities, and their immuno-senescence is rather related to a reduced efficiency in killing microbes than to a reduced number of circulating granulocytes or to a reduced efficiency of chemotaxis [15, 24–26]. Indeed, aging-related alterations observed in neutrophils appear

to be related to altered signaling pathways, such as the JAK/Stat and PI3-K/Akt/ERK1/2, resulting in an impaired production of cytokines and ROS [27, 28].

The alteration of NETosis due to aging has been poorly investigated although with controversial results.

We observed that neutrophils from elderly subjects respond to LPS stimulation by producing a greater amount of NETs compared to neutrophils obtained from adult subjects.

This effect could be linked to an altered redox activity in granulocytes obtained from elderly subjects, as a similar

effect was observed in aged mice. In fact, when granulocytes from aged mice were stimulated by 7-ketocholesterol, an athero-relevant stimulus showed an increased production of NETs linked to an increase in oxidative stress, caused by mitochondrial ROS production [29]. Therefore, mitochondria in neutrophils could act as powerful ROS generators to facilitate the innate immune function via NADPH-independent formation of NETs [30, 31].

Indeed, it has been reported that in aging, the key enzymatic activity responsible for NETs production, the Protein Arginine Deiminases-4 (PAD4), spontaneously generates NETs, and NETosis resulted exacerbated after stimulation [32, 33].

The observed increase in NETs production in elderly subjects is in contrast with the results obtained by Tseng et al. [16], who, in a murine model of *Staphylococcus aureus* infection, observed lower NETs production in neutrophils obtained by elderly animals compared to young ones. Moreover, Hazeldine et al. [17] observed in TNF α -primed human PMN a reduction of NETs production in aging. A possible explanation for this discrepancy could be due to both the LPS concentrations used for NETs production (10 ng/ml) and cells' stimulation condition used (TNF- α primed human neutrophils) [15, 22, 27]. Neutrophils from the elderly have shown a pre-activated baseline state with increased ROS production compared to neutrophils from young subjects [34, 35]. This may account for the differences observed between activated and resting neutrophils [36] and explain why aged-primed neutrophils produced a lower quantity of NETs than young ones, while aged not-primed neutrophils produce more NETs than young ones.

Tseng et al. [16] stimulated murine PMN directly with *S. aureus* and PMA (20 nM). Although PMA has been extensively used to study the biology of NETs in many publications, it is not a physiological stimulus [37], and it does not activate the JNK-dependent pathway [18].

Moreover, it could be interesting to compare in our experimental model the different quantification methods of NETs used in the literature to clarify the difference in NETs production in healthy elderly and adult subjects. However, despite increased NETs production in elderly subjects, these NETs showed less antimicrobial activity against *S. aureus*.

NETs antimicrobial activity is due both to the presence of high concentrations of antimicrobial molecules linked to the DNA meshwork and DNA ability to chelate manganese (Mn²⁺) and other divalent cations (Ca²⁺, Mg²⁺). Mn²⁺ acts as an electron transporter on the bacterial wall, playing an important role in microbial survival and proliferation, and its chelation prevents this transport [6, 7, 38]. DNA chelating activity is linked to its thread length, as long DNA fragments showed low chelating activity and bactericidal effect [39]. Even very small fragments of DNA showed a low

bactericidal effect, as bacteria extracellular DNase activity facilitates escape from NETs [40].

We analyzed soluble NETs in flow cytometry and observed that DNA from elderly subjects exhibited a shift towards larger fragments compared to DNA size distribution observed in adult subjects.

The presence of large DNA fragments in NETs could justify the low bactericidal activity observed.

We have previously demonstrated that NETs induce proliferation in human keratinocytes (HaCaT) [13], NETs act on keratinocytes through the internalization of double-strand DNA by TLR9 receptors, which induce a NF- κ B-dependent proliferation of keratinocytes [16]. This phenomenon is NET-concentration-dependent, with low physiological NETs concentration increasing keratinocyte proliferation, while high NETs concentration reduces their proliferation [13]. In our experiments, NETs from elderly failed to stimulate cell proliferation.

In addition, in this case, large DNA fragments have been found to be less efficient in the interaction with and activation of TLR9 [39]. The presence of large (or perhaps less decondensed) DNA fragments in NETs could be due to the epigenetic structure of the DNA such as age-related DNA methylation patterns [41, 42]. Other possible epigenetic mechanism is change in chromatin structure so that genomic DNA is wrapped within a chromatin structure that both compacts and protects DNA from interaction with surrounding factors. The basic unit of chromatin (nucleosome) packages the DNA with histone proteins such as H3 and H4. Methylated DNA gets wrapped around nucleosomes in a tight conformation which makes it relatively inaccessible [41, 42]. Therefore, some methylated chromatin areas could produce large fragments of NETs.

This hypothesis may explain our results, being larger DNA threads less effective in inducing both microbial toxicity and keratinocyte proliferation.

The presence of large size of DNA threads in NETs from elderly subjects has not been previously reported and it needs a more careful analysis. In fact, NETs degradation is found to be impaired in several autoimmune diseases, such as SLE [43–45] mainly due to DNase I deficiency, and compromised NET clearance in turn leads to inflammation and NETs autoantigens production [46]. Large size DNA fragments in NETs could be more difficult to clear becoming a trigger for an increased inflammation and autoimmune response in aging. Therefore, since NETs contain various active enzymes and DAMPs [47], their physiological clearance by DNase 1 action and macrophage phagocytosis is crucial [43, 48].

In conclusion, our results show a less explored aspect of the NETs linked to aging, consisting in a modification of their structure, resulting in both less antimicrobial activity and stimulation of keratinocyte proliferation, compared to

NETs obtained from adult subjects. Both effects could have a major impact on wound healing process in healthy elderly subjects.

Acknowledgements The authors thank Prof. Elisa Gamalero from the Department of Sciences and Innovative Technology (DISIT), Università del Piemonte Orientale for comments and discussion on portions of the manuscript. The authors thank M.P. Fortunato for her precious help with the language.

Author contributions All the authors contributed to the study conception and design. Material preparation, data collection and analysis were performed by EB, GN, MM and MS. The first draft of the manuscript was written by MS and FR, and all the authors commented on previous versions of the manuscript. All the authors read and approved the final manuscript.

Funding Open access funding provided by Università degli Studi del Piemonte Orientale Amedeo Avogadro within the CRUI-CARE Agreement.

Declarations

Conflict of interest On behalf of all the authors, the corresponding author states that there is no conflict of interest.

Statement of human rights The project was conducted with the approval of the Hospital “Maggiore della Carità” Ethic Committee (C.E. 61/10).

Informed consent All the participants underwent an informed consent process prior to participating.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Scapini P, Cassatella MA (2014) Social networking of human neutrophils within the immune system. *Blood* 124:710–719
- Brinkmann V, Reichard U, Goosmann C et al (2004) Neutrophil extracellular traps kill bacteria. *Science* 303:1532–1535
- Wang Y, Li M, Stadler S et al (2009) Histone hypercitrullination mediates chromatin decondensation and neutrophil extracellular trap formation. *J Cell Biol* 184:205–213
- Ito T (2014) PAMPs and DAMPs as triggers for DIC. *J Intensive Care* 2:67
- Yipp BG, Petri B, Salina D et al (2012) Infection-induced NETosis is a dynamic process involving neutrophil multitasking in vivo. *Nat Med* 18:1386–1393
- Polyanichko AM, Andrushchenko VV, Chikhirzhina EV et al (2004) The effect of manganese(II) on DNA structure: electronic and vibrational circular dichroism studies. *Nucleic Acids Res* 32:989–996
- Juttukonda LJ, Skaar EP (2015) Manganese homeostasis and utilization in pathogenic bacteria. *Mol Microbiol* 97:216–228
- Fuchs TA, Abed U, Goosmann C et al (2007) Novel cell death program leads to neutrophil extracellular traps. *J Cell Biol* 176:231–241
- Steinberg BE, Grinstein S (2007) Unconventional roles of the NADPH oxidase: signalling, ion homeostasis, and cell death. *Sci STKE* 2007:pe11
- Meng W, Paunel-Gorgulu A, Flohe S et al (2012) Depletion of neutrophil extracellular traps in vivo results in hypersusceptibility to polymicrobial sepsis in mice. *Crit Care* 16:R137
- Oehmcke S, Morgelin M, Herwald H (2009) Activation of the human contact system on neutrophil extracellular traps. *J Innate Immun* 1:225–230
- Tillack K, Breiden P, Martin R et al (2012) T lymphocyte priming by neutrophil extracellular traps links innate and adaptive immune responses. *J Immunol* 188:3150–3159
- Tonello S, Rizzi M, Migliario M et al (2017) Low concentrations of neutrophil extracellular traps induce proliferation in human keratinocytes via NF- κ B activation. *J Dermatol Sci* 88:110–116
- Sabbatini M, Magnelli V, Renò F (2021) NETosis in wound healing: when enough is enough. *Cells* 10:494–506
- Montgomery RR (2017) Age-related alterations in immune responses to West Nile virus infection. *Clin Exp Immunol* 187:26–34
- Tseng CW, Kyme PA, Arruda A et al (2012) Innate immune dysfunctions in aged mice facilitate the systemic dissemination of methicillin-resistant *S. aureus*. *PLoS ONE* 7:e41454
- Hazeldine J, Harris P, Chapple IL et al (2014) Impaired neutrophil extracellular trap formation: a novel defect in the innate immune system of aged individuals. *Aging Cell* 13:690–698
- Khan MA, Farahvash A, Douda ND et al (2017) JNK activation turns on LPS- and gram-negative bacteria-induced NADPH oxidase-dependent suicidal NETosis. *Sci Rep* 7:3409
- Chen L, Guo S, Ranzer MJ et al (2013) Toll-like receptor 4 has an essential role in early skin wound healing. *J Invest Dermatol* 133:258–267
- Vong L, Sherman PM, Glogauer M (2013) Quantification and visualization of neutrophil extracellular traps (NETs) from murine bone marrow-derived neutrophils. *Methods Mol Biol* 1031:41–50
- Delobel P, Tesnière C (2014) A simple FCM method to avoid misinterpretation in *Saccharomyces cerevisiae* cell cycle assessment between G0 and sub-G1. *PLoS ONE* 9:e84645
- Simon AK, Hollander GA, McMichael A (2015) Evolution of the immune system in humans from infancy to old age. *Proc Biol Sci* 282:20143085
- Montecino-Rodriguez E, Berent-Maoz B, Dorshkind K (2013) Causes, consequences, and reversal of immune system aging. *J Clin Invest* 123:958–965
- Butcher SK, Chahal H, Nayak L et al (2001) Senescence in innate immune responses: reduced neutrophil phagocytic capacity and CD16 expression in elderly humans. *J Leukoc Biol* 70:881–886
- Wenisch C, Patruta S, Daxbock F et al (2000) Effect of age on human neutrophil function. *J Leukoc Biol* 67:40–45
- Shaw AC, Goldstein DR, Montgomery RR (2013) Age-dependent dysregulation of innate immunity. *Nat Rev Immunol* 13:875–887
- Fulop T, Le Page A, Fortin C et al (2014) Cellular signalling in the aging immune system. *Curr Opin Immunol* 29:105–111
- Tortorella C, Simone O, Piazzolla G et al (2006) Role of phosphoinositide 3-kinase and extracellular signal-regulated kinase pathways in granulocyte macrophage-colony-stimulating factor failure

- to delay fas-induced neutrophil apoptosis in elderly humans. *J Gerontol A Biol Sci Med Sci* 61:1111–1118
29. Wang Y, Wang W, Wang N et al (2017) Mitochondrial oxidative stress promotes atherosclerosis and neutrophil extracellular traps in aged mice. *Arterioscler Thromb Vasc Biol* 37:e99–e107
 30. Ravindran M, Khan MA, Palaniyar N (2019) Neutrophil extracellular trap formation: physiology, pathology, and pharmacology. *Biomolecules* 9:365
 31. Wang C-H, Wu S-B, Wu Y-T et al (2013) Oxidative stress response elicited by mitochondrial dysfunction: implication in the pathophysiology of aging. *Exp Biol Med* 238:450–460
 32. Martinod K, Demers M, Fuchs TA et al (2013) Neutrophil histone modification by peptidylargininedeiminase 4 is critical for deep vein thrombosis in mice. *PNAS USA* 110:8674–8679
 33. Wong SL, Wagner DD (2018) Peptidylarginine deiminase 4: a nuclear button triggering neutrophil extracellular traps in inflammatory diseases and aging. *FASEB J* 32:6358–6370
 34. Kovalenko EI, Boyko AA, Semenov VF et al (2014) ROS production, intracellular HSP70 levels and their relationship in human neutrophils: effects of age. *Oncotarget* 5:11800–11812
 35. Nogueira-Neto J, Cardoso AS, Monteiro HP et al (2016) Basal neutrophil function in human aging: implications in endothelial cell adhesion. *Cell Biol Int* 40:796–802
 36. Jackaman C, Tomay F, Duong L et al (2017) Aging and cancer: the role of macrophages and neutrophils. *Ageing Res Rev* 36:105–116
 37. van der Linden M, Westerlaken GHA, van der Vlist M et al (2017) Differential signalling and kinetics of neutrophil extracellular trap. Release revealed by quantitative live imaging. *Sci Rep* 7:6529
 38. Ehrlich HL (2008) Are gram-positive bacteria capable of electron transfer across their cell wall without an externally available electron shuttle? *Geobiology* 6:220–224
 39. Bhongir RKV, Kasetty G, Papareddy P et al (2017) DNA-fragmentation is a source of bactericidal activity against *Pseudomonas aeruginosa*. *Biochem J* 474:411–425
 40. Berends ETM, Horswill AR, Haste NM et al (2010) Nuclease expression by *Staphylococcus aureus* facilitates escape from neutrophil extracellular traps. *J Innate Immun* 2:576–586
 41. Xiao F-H, Wang H-T, Kong Q-P (2019) Dynamic DNA methylation during aging: a “prophet” of age-related outcomes. *Front Genet* 10: Article 107
 42. Dor Yand Cedar H (2018) Principles of DNA methylation and their implications for biology and medicine. *Lancet* 392:777–786
 43. Hakkim A, Fürnrohr BG, Amann K et al (2010) Impairment of neutrophil extracellular trap degradation is associated with lupus nephritis. *Proc Natl Acad Sci USA* 107:9813–9818
 44. Leffler J, Martin M, Gullstrand B et al (2012) Neutrophil extracellular traps that are not degraded in systemic lupus erythematosus activate complement exacerbating the disease. *J Immunol* 188:3522–3531
 45. Mahajan A, Herrmann M, Muñoz LE (2016) Clearance deficiency and cell death pathways: a model for the pathogenesis of SLE. *Front Immunol* 7:35
 46. Lee KH, Kronbichler A, Park DD-Y et al (2017) Neutrophil extracellular traps (NETs) in autoimmune diseases: a comprehensive review. *Autoimmun Rev* 16:1160–1173
 47. Chapman EA, Lyon M, Simpson D et al (2019) Caught in a trap? Proteomic analysis of neutrophil extracellular traps in rheumatoid arthritis and systemic lupus erythematosus. *Front Immunol* 10:423
 48. Lazzaretto B, Fadeel B (2019) Intra- and extra-cellular degradation of neutrophil extracellular traps by macrophages and dendritic cells. *J Immunol* 203:2276–2290

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.