

Modular and duality properties of surface operators in $\mathcal{N} = 2^*$ gauge theories

S.K. Ashok,^{a,b} M. Billò,^c E. Dell'Aquila,^{a,b} M. Frau,^c R.R. John^{a,b} and A. Lerda^d

^a*Institute of Mathematical Sciences,
C.I.T. Campus, Taramani, Chennai, 600113 India*

^b*Homi Bhabha National Institute,
Training School Complex, Anushakti Nagar, Mumbai, 400085 India*

^c*Università di Torino, Dipartimento di Fisica and I.N.F.N. — sezione di Torino,
Via P. Giuria 1, I-10125 Torino, Italy*

^d*Università del Piemonte Orientale, Dipartimento di Scienze e Innovazione Tecnologica and
I.N.F.N. — sezione di Torino,
Via P. Giuria 1, I-10125 Torino, Italy*

E-mail: sashok@imsc.res.in, billo@to.infn.it, edellaquila@imsc.res.in,
frau@to.infn.it, renjan@imsc.res.in, lerda@to.infn.it

ABSTRACT: We calculate the instanton partition function of the four-dimensional $\mathcal{N} = 2^*$ $SU(N)$ gauge theory in the presence of a generic surface operator, using equivariant localization. By analyzing the constraints that arise from S-duality, we show that the effective twisted superpotential, which governs the infrared dynamics of the two-dimensional theory on the surface operator, satisfies a modular anomaly equation. Exploiting the localization results, we solve this equation in terms of elliptic and quasi-modular forms which resum all non-perturbative corrections. We also show that our results, derived for monodromy defects in the four-dimensional theory, match the effective twisted superpotential describing the infrared properties of certain two-dimensional sigma models coupled either to pure $\mathcal{N} = 2$ or to $\mathcal{N} = 2^*$ gauge theories.

KEYWORDS: Duality in Gauge Field Theories, Extended Supersymmetry, Supersymmetry and Duality, D-branes

ARXIV EPRINT: [1702.02833](https://arxiv.org/abs/1702.02833)

Contents

1	Introduction	2
2	Instantons and surface operators in $\mathcal{N} = 2^* \text{SU}(N)$ gauge theories	4
3	Partition functions for ramified instantons	7
3.1	Summing over fixed points and characters	11
3.2	Map between parameters	12
3.3	Extracting the prepotential and the twisted superpotential	13
4	Modular anomaly equation for the twisted superpotential	15
4.1	S-duality constraints	17
5	Surface operators in $\mathcal{N} = 2^* \text{SU}(2)$ theory	20
5.1	Relation to CFT results	23
6	Surface operators in $\mathcal{N} = 2^* \text{SU}(N)$ theories	24
6.1	Simple surface operators	24
6.2	Surface operators of type $\{p, N - p\}$	26
6.3	Surface operators of general type	27
7	Duality between surface operators	28
7.1	The pure $\mathcal{N} = 2 \text{SU}(N)$ theory	28
7.2	The $\mathcal{N} = 2^* \text{SU}(N)$ theory	31
7.3	Some remarks on the results	33
8	Conclusions	34
A	Useful formulas for modular forms and elliptic functions	35
B	Generalized instanton number in the presence of fluxes	38
C	Ramified instanton moduli and their properties	41
C.1	$\text{SU}(2)$	43
D	Prepotential coefficients for the $\text{SU}(N)$ gauge theory	45

1 Introduction

The study of how a quantum field theory responds to the presence of defects is a very important subject, which has received much attention in recent years especially in the context of supersymmetric gauge theories. In this paper we study a class of two-dimensional defects, also known as surface operators, on the Coulomb branch of the $\mathcal{N} = 2^* \text{SU}(N)$ gauge theory in four dimensions.¹ Such surface operators can be introduced and analyzed in different ways. They can be defined by the transverse singularities they induce in the four-dimensional fields [2, 3], or can be characterized by the two-dimensional theory they support on their world-volume [4, 5].

A convenient way to describe four-dimensional gauge theories with $\mathcal{N} = 2$ supersymmetry is to consider M5 branes wrapped on a punctured Riemann surface [6, 7]. From the point of view of the six-dimensional $(2, 0)$ theory on the M5 branes, surface operators can be realized by means of either M5' or M2 branes giving rise, respectively, to codimension-2 and codimension-4 defects. While a codimension-2 operator extends over the Riemann surface wrapped by the M5 brane realizing the gauge theory, a codimension-4 operator intersects the Riemann surface at a point. Codimension-2 surface operators were systematically studied in [8] where, in the context of the of the $4d/2d$ correspondence [9], the instanton partition functions of $\mathcal{N} = 2 \text{SU}(2)$ super-conformal quiver theories with surface operators were mapped to the conformal blocks of a two-dimensional conformal field theory with an affine $\text{sl}(2)$ symmetry. These studies were later extended to $\text{SU}(N)$ quiver theories whose instanton partition functions in the presence of surface operators were related to conformal field theories with an affine $\text{sl}(N)$ symmetry [10]. The study of codimension-4 surface operators was pioneered in [11] where the instanton partition function of the conformal $\text{SU}(2)$ theory with a surface operator was mapped to the Virasoro blocks of the Liouville theory, augmented by the insertion of a degenerate primary field. Many generalizations and extensions of this have been considered in the last few years [12–19].

Here we study $\mathcal{N} = 2^*$ theories in the presence of surface operators. The low-energy effective dynamics of the bulk four-dimensional theory is completely encoded in the holomorphic prepotential which at the non-perturbative level can be very efficiently determined using localization [20] along with the constraints that arise from S-duality. The latter turn out to imply [21, 22] a modular anomaly equation [23] for the prepotential, which is intimately related to the holomorphic anomaly equation occurring in topological string theories on local Calabi-Yau manifolds [24–27].² Working perturbatively in the mass of the adjoint hypermultiplet, the modular anomaly equation allows one to resum all instanton corrections to the prepotential into (quasi)-modular forms, and to write the dependence on the Coulomb branch parameters in terms of particular sums over the roots of the gauge group, thus making it possible to treat any semi-simple algebra [41, 42].

¹For a review of surface operators see [1].

²Modular anomaly equations have been studied in various contexts, such as the Ω -background [21, 22, 28–34], the $4d/2d$ correspondence [35–37], SQCD theories with fundamental matter [21, 22, 38–40] and in $\mathcal{N} = 2^*$ theories [21, 22, 41–44].

In this paper we apply the same approach to study the effective twisted superpotential which governs the infrared dynamics on the world-volume of the two-dimensional surface operator in the $\mathcal{N} = 2^*$ theory. For simplicity, we limit ourselves to $SU(N)$ gauge groups and consider half-BPS surface defects that, from the six-dimensional point of view, are codimension-2 operators. These defects introduce singularities characterized by the pattern of gauge symmetry breaking, *i.e.* by a Levi decomposition of $SU(N)$, and also by a set of continuous (complex) parameters. In [45] it has been shown that the effect of these surface operators on the instanton moduli action is equivalent to a suitable orbifold projection which produces structures known as ramified instantons [45–47]. Actually, the moduli spaces of these ramified instantons were already studied in [48] from a mathematical point of view in terms of representations of a quiver that can be obtained by performing an orbifold projection of the usual ADHM moduli space of the standard instantons. In section 2 we explicitly implement such an orbifold procedure on the non-perturbative sectors of the theory realized by means of systems of D3 and D(−1) branes [49, 50]. In section 3 we carry out the integration on the ramified instanton moduli via equivariant localization. The logarithm of the resulting partition function exhibits both a $4d$ and a $2d$ singularity in the limit of vanishing Ω deformations.³ The corresponding residues are regular in this limit and encode, respectively, the prepotential \mathcal{F} and the twisted superpotential \mathcal{W} . The latter depends, in addition to the Coulomb vacuum expectation values and the adjoint mass, on the continuous parameters of the defect.

In section 4 we show that, as it happens for the prepotential, the constraints arising from S-duality lead to a modular anomaly equation for \mathcal{W} . In section 5, we solve this equation explicitly for the $SU(2)$ theory and prove that the resulting \mathcal{W} agrees with the twisted superpotential obtained in [35] in the framework of the $4d/2d$ correspondence with the insertion of a degenerate field in the Liouville theory. Since this procedure is appropriate for codimension-4 defects [11], the agreement we find supports the proposal of a duality between the two classes of defects recently put forward in [52]. In section 6, we turn our attention to generic surface operators in the $SU(N)$ theory and again, order by order in the adjoint mass, solve the modular anomaly equations in terms of quasi-modular elliptic functions and sums over the root lattice.

We also consider the relation between our findings and what is known for surface defects defined through the two-dimensional theory they support on their world-volume. In [5] the coupling of the sigma-models defined on such defects to a large class of four-dimensional gauge theories was investigated and the twisted superpotential governing their dynamics was obtained. Simple examples for pure $\mathcal{N} = 2$ $SU(N)$ gauge theory include the linear sigma-model on $\mathbb{C}\mathbb{P}^{N-1}$, that corresponds to the so-called simple defects with Levi decomposition of type $\{1, N - 1\}$, and sigma-models on Grassmannian manifolds corresponding to defects of type $\{p, N - p\}$. The main result of [5] is that the Seiberg-Witten geometry of the four-dimensional theory can be recovered by analyzing how the vacuum structure of these sigma-models is fibered over the Coulomb moduli space. Independent

³We actually calculate the effective superpotential in the Nekrasov-Shatashvili limit [51] in which only one of the Ω -deformation parameters is turned on.

analyses based on the $4d/2d$ correspondence also show that the twisted superpotential for the simple surface operator is related to the line integral of the Seiberg-Witten differential over the punctured Riemann surface [11]. In section 7, we test this claim in detail by considering first the pure $\mathcal{N} = 2$ gauge theory. Since this theory can be recovered upon decoupling the massive adjoint hypermultiplet, we take the decoupling limit on our $\mathcal{N} = 2^*$ results for \mathcal{W} and precisely reproduce those findings. Furthermore, we show that for simple surface defects the relation between the twisted superpotential and the line integral of the Seiberg-Witten differential holds prior to the decoupling limit, *i.e.* in the $\mathcal{N} = 2^*$ theory itself. The agreement we find provides evidence for the proposed duality between the two types of descriptions of the surface operators.

Finally, in section 8 we present our conclusions and discuss possible future perspectives. Some useful technical details are provided in four appendices.

2 Instantons and surface operators in $\mathcal{N} = 2^*$ $SU(N)$ gauge theories

The $\mathcal{N} = 2^*$ theory is a four-dimensional gauge theory with $\mathcal{N} = 2$ supersymmetry that describes the dynamics of a vector multiplet and a massive hypermultiplet in the adjoint representation. It interpolates between the $\mathcal{N} = 4$ super Yang-Mills theory, to which it reduces in the massless limit, and the pure $\mathcal{N} = 2$ theory, which is recovered by decoupling the matter hypermultiplet. In this paper, we will consider for simplicity only special unitary gauge groups $SU(N)$. As is customary, we combine the Yang-Mills coupling constant g and the vacuum angle θ into the complex coupling

$$\tau = \frac{\theta}{2\pi} + i \frac{4\pi}{g^2}, \tag{2.1}$$

on which the modular group $SL(2, \mathbb{Z})$ acts in the standard fashion:

$$\tau \rightarrow \frac{a\tau + b}{c\tau + d} \tag{2.2}$$

with $a, b, c, d \in \mathbb{Z}$ and $ad - bc = 1$. In particular under S-duality we have

$$S(\tau) = -\frac{1}{\tau}. \tag{2.3}$$

The Coulomb branch of the theory is parametrized by the vacuum expectation value of the adjoint scalar field ϕ in the vector multiplet, which we take to be of the form

$$\langle \phi \rangle = \text{diag}(a_1, a_2, \dots, a_N) \quad \text{with} \quad \sum_{u=1}^N a_u = 0. \tag{2.4}$$

The low-energy effective dynamics on the Coulomb branch is entirely described by a single holomorphic function \mathcal{F} , called the prepotential, which contains a classical term, a perturbative 1-loop contribution and a tail of instanton corrections. The latter can be obtained from the instanton partition function

$$Z_{\text{inst}} = \sum_{k=0}^{\infty} q^k Z_k \tag{2.5}$$

where

$$q = e^{2\pi i \tau} \tag{2.6}$$

and Z_k is the partition function in the k -instanton sector that can be explicitly computed using localization methods.⁴ For later purposes, it is useful to recall that the weight q^k in (2.5) originates from the classical instanton action

$$S_{\text{inst}} = -2\pi i \tau \left(\frac{1}{8\pi^2} \int_{\mathbb{R}^4} \text{Tr } F \wedge F \right) = -2\pi i \tau k \tag{2.7}$$

where in the last step we used the fact that the second Chern class of the gauge field strength F equals the instanton charge k . Hence, the weight q^k is simply $e^{-S_{\text{inst}}}$.

Let us now introduce a surface operator which we view as a non-local defect D supported on a two-dimensional plane inside the four-dimensional (Euclidean) space-time (see appendix B for more details). In particular, we parametrize $\mathbb{R}^4 \simeq \mathbb{C}^2$ by two complex variables (z_1, z_2) , and place D at $z_2 = 0$, filling the z_1 -plane. The presence of the surface operator induces a singular behavior in the gauge connection A , which has the following generic form [8, 45]:

$$A = A_\mu dx^\mu \simeq - \text{diag} \left(\underbrace{\gamma_1, \dots, \gamma_1}_{n_1}, \underbrace{\gamma_2, \dots, \gamma_2}_{n_2}, \dots, \underbrace{\gamma_M, \dots, \gamma_M}_{n_M} \right) d\theta \tag{2.8}$$

as $r \rightarrow 0$. Here (r, θ) denotes the set of polar coordinates in the z_2 -plane, and the γ_I 's are constant parameters, where $I = 1, \dots, M$. The M integers n_I satisfy

$$\sum_{I=1}^M n_I = N \tag{2.9}$$

and define a vector \vec{n} that identifies the type of the surface operator. This vector is related to the breaking pattern of the gauge group (or Levi decomposition) felt on the two-dimensional defect D , namely

$$\text{SU}(N) \rightarrow \text{S}[\text{U}(n_1) \times \text{U}(n_2) \times \dots \times \text{U}(n_M)]. \tag{2.10}$$

The type $\vec{n} = \{1, 1, \dots, 1\}$ corresponds to what are called full surface operators, originally considered in [8]. The type $\vec{n} = \{1, N - 1\}$ corresponds to simple surface operators, while the type $\vec{n} = \{N\}$ corresponds to no surface operators and hence will not be considered.

In the presence of a surface operator, one can turn on magnetic fluxes for each factor of the gauge group (2.10) and thus the instanton action can receive contributions also from the corresponding first Chern classes. This means that (2.7) is replaced by [2, 8, 11, 45]

$$S_{\text{inst}}[\vec{n}] = -2\pi i \tau \left(\frac{1}{8\pi^2} \int_{\mathbb{R}^4} \text{Tr } F \wedge F \right) - 2\pi i \sum_{I=1}^M \eta_I \left(\frac{1}{2\pi} \int_D \text{Tr } F_{\text{U}(n_I)} \right) \tag{2.11}$$

⁴Our conventions are such that $Z_0 = 1$.

where η_I are constant parameters. As shown in detail in appendix B, given the behavior (2.8) of the gauge connection near the surface operator, one has

$$\begin{aligned} \frac{1}{8\pi^2} \int_{\mathbb{R}^4} \text{Tr} F \wedge F &= k + \sum_{I=1}^M \gamma_I m_I, \\ \frac{1}{2\pi} \int_D \text{Tr} F_{U(n_I)} &= m_I \end{aligned} \tag{2.12}$$

with $m_I \in \mathbb{Z}$. As is clear from the second line in the above equation, each m_I represents the flux of the U(1) factor in each subgroup U(n_I) in the Levi decomposition (2.10); furthermore, these fluxes satisfy the constraint

$$\sum_{I=1}^M m_I = 0. \tag{2.13}$$

Using (2.12), we easily find

$$S_{\text{inst}}[\vec{n}] = -2\pi i \tau k - 2\pi i \sum_{I=1}^M (\eta_I + \tau \gamma_I) m_I = -2\pi i \tau k - 2\pi i \vec{t} \cdot \vec{m} \tag{2.14}$$

where in the last step we have combined the electric and magnetic parameters (η_I, γ_I) to form the M -dimensional vector

$$\vec{t} = \{t_I\} = \{\eta_I + \tau \gamma_I\}. \tag{2.15}$$

This combination has simple duality transformation properties under $\text{SL}(2, \mathbb{Z})$. Indeed, as shown in [2], given an element \mathcal{M} of the modular group the electro-magnetic parameters transform as

$$(\gamma_I, \eta_I) \rightarrow (\gamma_I, \eta_I) \mathcal{M}^{-1} = (d\gamma_I - c\eta_I, a\eta_I - b\gamma_I). \tag{2.16}$$

Combining this with the modular transformation (2.2) of the coupling constant, it is easy to show that

$$t_I \rightarrow \frac{t_I}{c\tau + d}. \tag{2.17}$$

In particular under S-duality we have

$$S(t_I) = -\frac{t_I}{\tau}. \tag{2.18}$$

Using (2.14), we deduce that the weight of an instanton configuration in the presence of a surface operator of type \vec{n} is

$$e^{-S_{\text{inst}}[\vec{n}]} = q^k e^{2\pi i \vec{t} \cdot \vec{m}}, \tag{2.19}$$

so that the instanton partition function can be written as

$$Z_{\text{inst}}[\vec{n}] = \sum_{k, \vec{m}} q^k e^{2\pi i \vec{t} \cdot \vec{m}} Z_{k, \vec{m}}[\vec{n}]. \tag{2.20}$$

In the next section, we will describe the computation of $Z_{k, \vec{m}}[\vec{n}]$ using equivariant localization.

3 Partition functions for ramified instantons

As discussed in [45], the $\mathcal{N} = 2^*$ theory with a surface defect of type $\vec{n} = \{n_1, \dots, n_M\}$, which has a six-dimensional representation as a codimension-2 surface operator, can be realized with a system of D3-branes in the orbifold background

$$\mathbb{C} \times \mathbb{C}^2 / \mathbb{Z}_M \times \mathbb{C} \times \mathbb{C} \tag{3.1}$$

with coordinates (z_1, z_2, z_3, z_4, v) on which the \mathbb{Z}_M -orbifold acts as

$$(z_2, z_3) \rightarrow (\omega z_2, \omega^{-1} z_3), \quad \text{where } \omega = e^{\frac{2\pi i}{M}}. \tag{3.2}$$

Like in the previous section, the complex coordinates z_1 and z_2 span the four-dimensional space-time where the gauge theory is defined (namely the world-volume of the D3-branes), while the z_1 -plane is the world-sheet of the surface operator D that sits at the orbifold fixed point $z_2 = 0$. The (massive) deformation which leads from the $\mathcal{N} = 4$ to the $\mathcal{N} = 2^*$ theory takes place in the (z_3, z_4) -directions. Finally, the v -plane corresponds to the Coulomb moduli space of the gauge theory.

Without the \mathbb{Z}_M -orbifold projection, the isometry group of the ten-dimensional background is $SO(4) \times SO(4) \times U(1)$, since the D3-branes are extended in the first four directions and are moved in the last two when the vacuum expectation values (2.4) are turned on. In the presence of the surface operator and hence of the \mathbb{Z}_M -orbifold in the (z_2, z_3) -directions, this group is broken to

$$U(1) \times U(1) \times U(1) \times U(1) \times U(1). \tag{3.3}$$

In the following we will focus only on the first four $U(1)$ factors, since it is in the first four complex directions that we will introduce equivariant deformations to apply localization methods. We parameterize a transformation of this $U(1)^4$ group by the vector

$$\vec{\epsilon} = \left\{ \epsilon_1, \frac{\epsilon_2}{M}, \frac{\epsilon_3}{M}, \epsilon_4 \right\} = \{ \epsilon_1, \hat{\epsilon}_2, \hat{\epsilon}_3, \epsilon_4 \} \tag{3.4}$$

where the $1/M$ rescalings in the second and third entry, suggested by the orbifold projection, are made for later convenience. If we denote by

$$\vec{l} = \{ l_1, l_2, l_3, l_4 \} \tag{3.5}$$

the weight vector of a given state of the theory, then under $U(1)^4$ such a state transforms with a phase given by $e^{2\pi i \vec{l} \cdot \vec{\epsilon}}$, while the \mathbb{Z}_M -action produces a phase $\omega^{l_2 - l_3}$.

On top of this, we also have to consider the action of the orbifold group on the Chan-Paton factors carried by the open string states stretching between the D-branes. There are different types of D-branes depending on the irreducible representation of \mathbb{Z}_M in which this action takes place. Since there are M such representations, we have M types of D-branes, which we label with the index I already used before. On a D-brane of type I , the generator of \mathbb{Z}_M acts as ω^I , and thus the Chan-Paton factor of a string stretching between a D-brane

of type I and a D-brane of type J transforms with a phase ω^{I-J} under the action of the orbifold generator.

In order to realize the split of the gauge group in (2.10), we consider M stacks of n_I D3-branes of type I , and in order to introduce non-perturbative effects we add on top of the D3's M stacks of d_I D-instantons of type I . The latter support an auxiliary ADHM group which is

$$U(d_1) \times U(d_2) \times \cdots \times U(d_M). \quad (3.6)$$

In the resulting D3/D(-1)-brane systems there are many different sectors of open strings depending on the different types of branes to which they are attached. Here we focus only on the states of open strings with at least one end-point on the D-instantons, because they represent the instanton moduli [49, 50] on which one eventually has to integrate in order to obtain the instanton partition function.

Let us first consider the neutral states, corresponding to strings stretched between two D-instantons. In the bosonic Neveu-Schwarz sector one finds states with $U(1)^4$ weight vectors

$$\{\pm 1, 0, 0, 0\}_0, \quad \{0, \pm 1, 0, 0\}_0, \quad \{0, 0 \pm 1, 0\}_0, \quad \{0, 0, 0 \pm 1\}_0, \quad \{0, 0, 0, 0\}_{\pm 1}, \quad (3.7)$$

where the subscripts denote the charge under the last $U(1)$ factor of (3.3). They correspond to space-time vectors along the directions z_1, z_2, z_3, z_4 and v , respectively. In the fermionic Ramond sector one finds states with weight vectors

$$\left\{ \pm \frac{1}{2}, \pm \frac{1}{2}, \pm \frac{1}{2}, \pm \frac{1}{2} \right\}_{\pm \frac{1}{2}} \quad (3.8)$$

with a total odd number of minus signs due to the GSO projection. They correspond to anti-chiral space-time spinors.⁵

It is clear from (3.7) and (3.8) that the orbifold phase $\omega^{l_2-l_3}$ takes the values ω^0, ω^{+1} or ω^{-1} and can be compensated only if one considers strings of type $I-I, I-(I+1)$ or $(I+1)-I$, respectively. Therefore, the \mathbb{Z}_M -invariant neutral moduli carry Chan-Paton factors that transform in the $(\mathbf{d}_I, \bar{\mathbf{d}}_I), (\mathbf{d}_I, \bar{\mathbf{d}}_{I+1})$ or $(\mathbf{d}_{I+1}, \bar{\mathbf{d}}_I)$ representations of the ADHM group (3.6).

Let us now consider the colored states, corresponding to strings stretched between a D-instanton and a D3-brane or vice versa. Due to the twisted boundary conditions in the first two complex space-time directions, the weight vectors of the bosonic states in the Neveu-Schwarz sector are

$$\left\{ \pm \frac{1}{2}, \pm \frac{1}{2}, 0, 0 \right\}_0 \quad (3.9)$$

while those of the fermionic states in the Ramond sector are

$$\left\{ 0, 0, \pm \frac{1}{2}, \pm \frac{1}{2} \right\}_{\pm \frac{1}{2}}. \quad (3.10)$$

Assigning a negative intrinsic parity to the twisted vacuum, both in (3.9) and in (3.10) the GSO-projection selects only those vectors with an even number of minus signs. Moreover,

⁵Of course one could have chosen a GSO projection leading to chiral spinors, and the final results would have been the same.

since the orbifold acts on two of the twisted directions, the vacuum carries also an intrinsic \mathbb{Z}_M -weight. We take this to be $\omega^{-\frac{1}{2}}$ when the strings are stretched between a D3-brane and a D-instanton, and $\omega^{+\frac{1}{2}}$ for strings with opposite orientation. Then, with this choice we find \mathbb{Z}_M -invariant bosonic and fermionic states either from the $3/(-1)$ strings of type $I-I$, whose Chan-Paton factors transform in the $(\mathbf{n}_I, \bar{\mathbf{d}}_I)$ representation of the gauge and ADHM groups, or from the $(-1)/3$ strings of type $I-(I+1)$, whose Chan-Paton factors transform in the $(\mathbf{d}_I, \bar{\mathbf{n}}_{I+1})$ representation, plus of course the corresponding states arising from the strings with opposite orientation.

In appendix C we provide a detailed account of all moduli, both neutral and colored, and of their properties in the various sectors. It turns out that the moduli action, which can be derived from the interactions of the moduli on disks with at least a part of their boundary attached to the D-instantons [50], is exact with respect to the supersymmetry charge Q of weight

$$\left\{ +\frac{1}{2}, +\frac{1}{2}, +\frac{1}{2}, +\frac{1}{2} \right\}_{-\frac{1}{2}}. \tag{3.11}$$

Therefore Q can be used as the equivariant BRST-charge to localize the integral over the moduli space provided one considers $U(1)^4$ transformations under which it is invariant. This corresponds to requiring that

$$\epsilon_1 + \hat{\epsilon}_2 + \hat{\epsilon}_3 + \epsilon_4 = 0. \tag{3.12}$$

Thus we are left with three equivariant parameters, say ϵ_1 , $\hat{\epsilon}_2$ and ϵ_4 ; as we will see, the latter is related to the (equivariant) mass m of the adjoint hypermultiplet of $\mathcal{N} = 2^*$ theory.

As shown in appendix C, all instanton moduli can be paired in Q -doublets of the type $(\varphi_\alpha, \psi_\alpha)$ such that

$$Q \varphi_\alpha = \psi_\alpha, \quad Q \psi_\alpha = Q^2 \varphi_\alpha = \lambda_\alpha \varphi_\alpha \tag{3.13}$$

where λ_α are the eigenvalues of Q^2 , determined by the action of the Cartan subgroup of the full symmetry group of the theory, namely the gauge group (2.10), the ADHM group (3.6), and the residual isometry group $U(1)^4$ with parameters satisfying (3.12) in such a way that the invariant points in the moduli space are finite and isolated. The only exception to this structure of Q -doublets is represented by the neutral bosonic moduli with weight

$$\{0, 0, 0, 0\}_{-1} \tag{3.14}$$

transforming in the adjoint representation $(\mathbf{d}_I, \bar{\mathbf{d}}_I)$ of the ADHM group $U(d_I)$, which remain unpaired. We denote them as χ_I , and in order to obtain the instanton partition function we must integrate over them. In doing so, we can exploit the $U(d_I)$ symmetry to rotate χ_I into the maximal torus and write it in terms of the eigenvalues $\chi_{I,\sigma}$, with $\sigma = 1, \dots, d_I$, which represent the positions of the D -instantons of type I in the v -plane. In this way we are left with the integration over all the $\chi_{I,\sigma}$'s and a Cauchy-Vandermonde determinant

$$\mathcal{V} = \prod_{I=1}^M \prod_{\sigma,\tau=1}^{d_I} (\chi_{I,\sigma} - \chi_{I,\tau} + \delta_{\sigma\tau}). \tag{3.15}$$

More precisely, the instanton partition function in the presence of a surface operator of type \vec{n} is defined by

$$Z_{\text{inst}}[\vec{n}] = \sum_{\{d_I\}} \prod_{I=1}^M q_I^{d_I} Z_{\{d_I\}}[\vec{n}] \quad \text{with} \quad Z_{\{d_I\}}[\vec{n}] = \frac{1}{d_I!} \int \prod_{\sigma=1}^{d_I} \frac{d\chi_{I,\sigma}}{2\pi i} z_{\{d_I\}} \quad (3.16)$$

where $z_{\{d_I\}}$ is the result of the integration over all Q -doublets which localizes on the fixed points of Q^2 , and q_I is the counting parameter associated to the D-instantons of type I . With the convention that $z_{\{d_I=0\}} = 1$, we find

$$z_{\{d_I\}} = \mathcal{V} \prod_{\alpha} [\lambda_{\alpha}]^{(-)^{F_{\alpha}+1}}, \quad (3.17)$$

where the index α labels the Q -doublets and λ_{α} denotes the corresponding eigenvalue of Q^2 . This contribution goes to the denominator or to the numerator depending upon the bosonic or fermionic statistics ($F_{\alpha} = 0$ or 1 , respectively) of the first component of the doublet. Explicitly, using the data in table 1 of appendix C and the determinant (3.15), we find

$$\begin{aligned} z_{\{d_I\}} &= \prod_{I=1}^M \prod_{\sigma,\tau=1}^{d_I} \frac{(\chi_{I,\sigma} - \chi_{I,\tau} + \delta_{\sigma,\tau})(\chi_{I,\sigma} - \chi_{I,\tau} + \epsilon_1 + \epsilon_4)}{(\chi_{I,\sigma} - \chi_{I,\tau} + \epsilon_4)(\chi_{I,\sigma} - \chi_{I,\tau} + \epsilon_1)} \\ &\times \prod_{I=1}^M \prod_{\sigma=1}^{d_I} \prod_{\rho=1}^{d_{I+1}} \frac{(\chi_{I,\sigma} - \chi_{I+1,\rho} + \epsilon_1 + \hat{\epsilon}_2)(\chi_{I,\sigma} - \chi_{I+1,\rho} + \hat{\epsilon}_2 + \epsilon_4)}{(\chi_{I,\sigma} - \chi_{I+1,\rho} - \hat{\epsilon}_3)(\chi_{I,\sigma} - \chi_{I+1,\rho} + \hat{\epsilon}_2)} \\ &\times \prod_{I=1}^M \prod_{\sigma=1}^{d_I} \prod_{s=1}^{n_I} \frac{(a_{I,s} - \chi_{I,\sigma} + \frac{1}{2}(\epsilon_1 + \hat{\epsilon}_2) + \epsilon_4)}{(a_{I,s} - \chi_{I,\sigma} + \frac{1}{2}(\epsilon_1 + \hat{\epsilon}_2))} \\ &\times \prod_{I=1}^M \prod_{\sigma=1}^{d_I} \prod_{t=1}^{n_{I+1}} \frac{(\chi_{I,\sigma} - a_{I+1,t} + \frac{1}{2}(\epsilon_1 + \hat{\epsilon}_2) + \epsilon_4)}{(\chi_{I,\sigma} - a_{I+1,t} + \frac{1}{2}(\epsilon_1 + \hat{\epsilon}_2))} \end{aligned} \quad (3.18)$$

where $d_{M+1} = d_1$, $n_{M+1} = n_1$ and $a_{M+1,t} = a_{1,t}$. The integrations in (3.16) must be suitably defined and regularized. The standard prescription [41, 42, 53] is to consider $a_{I,s}$ to be real and close the contours in the upper-half $\chi_{I,\sigma}$ -planes with the choice

$$\text{Im } \epsilon_4 \gg \text{Im } \hat{\epsilon}_3 \gg \text{Im } \hat{\epsilon}_2 \gg \text{Im } \epsilon_1 > 0, \quad (3.19)$$

and enforce (3.12) at the very end of the calculations.

In this way one finds that these integrals receive contributions from the poles of $z_{\{d_I\}}$, which are in fact the critical points of Q^2 . Such poles can be put in one-to-one correspondence with a set of N Young tableaux $Y = \{Y_{I,s}\}$, with $I = 1, \dots, M$ and $s = 1, \dots, n_I$, in the sense that the box in the i -th row and j -th column of the tableau $Y_{I,s}$ represents one component of the critical value:

$$\chi_{I+(j-1)\text{mod}M,\sigma} = a_{I,s} + \left((i-1) + \frac{1}{2} \right) \epsilon_1 + \left((j-1) + \frac{1}{2} \right) \hat{\epsilon}_2. \quad (3.20)$$

Note that in this correspondence, a single tableau accounts for $d_I!$ equivalent ways of relabeling $\chi_{I,\sigma}$.

3.1 Summing over fixed points and characters

Summing over the Young tableaux collections Y we get all the non-trivial critical points corresponding to all possible values of $\{d_I\}$. Eq. (3.20) tells us that we get a distinct $\chi_{I,\sigma}$ for each box in the j -th column of the tableau $Y_{I+1-j \bmod M,s}$. Relabeling the index j as

$$j \rightarrow J + jM, \quad (3.21)$$

with $J = 1, \dots, M$, we have

$$d_I(Y) = \sum_{J=1}^M \sum_{s=1}^{n_{I+1-J}} \sum_j Y_{I+1-J,s}^{(J+jM)}, \quad (3.22)$$

where $Y_{I,s}^{(j)}$ denotes the height of the j -th column of the tableau $Y_{I,s}$, and the subscript index $I+1-J$ is understood modulo M .

The instanton partition function (3.16) can thus be rewritten as a sum over Young tableaux as follows

$$Z_{\text{inst}}[\vec{n}] = \sum_Y \prod_{I=1}^M q_I^{d_I(Y)} Z(Y) \quad (3.23)$$

where $Z(Y)$ is the residue of $z_{\{d_I\}}$ at the critical point Y . This is obtained by deleting in (3.18) the denominator factors that yield the identifications (3.20), and performing these identifications in the other factors. In other terms,

$$Z(Y) = \mathcal{V}(Y) \prod_{\alpha: \lambda_\alpha(Y) \neq 0} [\lambda_\alpha(Y)]^{(-)^{F_\alpha+1}}, \quad (3.24)$$

where $\mathcal{V}(Y)$ and $\lambda_\alpha(Y)$ are the Vandermonde determinant and the eigenvalues of Q^2 evaluated on (3.20).

A more efficient way to encode the eigenvalues $\lambda_\alpha(Y)$ is to employ the character of the action of Q^2 , which is defined as follows

$$X_{\{d_I\}} = \sum_{\alpha} (-)^{F_\alpha} e^{i\lambda_\alpha}. \quad (3.25)$$

If we introduce

$$V_I = \sum_{\sigma=1}^{d_I} e^{i\chi_{I,\sigma} - \frac{i}{2}(\epsilon_1 + \hat{\epsilon}_2)}, \quad W_I = \sum_{s=1}^{n_I} e^{ia_{I,s}} \quad (3.26)$$

and

$$T_1 = e^{i\epsilon_1}, \quad T_2 = e^{i\hat{\epsilon}_2}, \quad T_4 = e^{i\epsilon_4}, \quad (3.27)$$

we can write the contributions to the character from the various Q -doublets as in the last column of table 1 in appendix C. Then, by summing over all doublets and adding also the contribution of the Vandermonde determinant, we obtain

$$X_{\{d_I\}} = (1 - T_4) \sum_{I=1}^M \left[-(1 - T_1) V_I^* V_I + (1 - T_1) V_{I+1}^* V_I T_2 + V_I^* W_I + W_{I+1}^* V_I T_1 T_2 \right]. \quad (3.28)$$

As we have seen before, through (3.22) and (3.20) each set Y determines both the dimensions $d_I(Y)$ and the eigenvalues $\lambda_\alpha(Y)$. Thus, the character $X(Y)$ associated to a set of Young tableaux is obtained from $X_{\{d_I\}}$ by substituting (3.20) into the definitions of V_I , namely

$$V_I = \sum_{J=1}^M \sum_{s=1}^{n_{I+1-J}} e^{ia_{I+1-J,s}} T_2^J \sum_{(i,J+jM) \in Y_{I+1-J,s}} T_1^{i-1} T_2^{jM-1}. \quad (3.29)$$

By analyzing $X(Y)$ obtained in this way we can extract the explicit expression for the eigenvalues $\lambda_s(Y)$ and finally write the instanton partition function. This procedure is easily implemented in a computer program, and yields the results we will use in the next sections. In appendix (C.1), as an example, we illustrate these computations for the SU(2) gauge theory.

In our analysis we worked with the moduli action that describes D-branes probing the orbifold geometry. An alternative approach works with the resolution of the orbifold geometry [54, 55]. This involves analyzing a gauged linear sigma-model that describes a system of D1 and D5-branes in the background $\mathbb{C} \times \mathbb{C}/\mathbb{Z}_M \times T^*S^2 \times \mathbb{R}^2$. One then uses the localization formulas for supersymmetric field theories defined on the 2-sphere [56, 57] to obtain exact results. This is a very powerful approach since it also includes inherently stringy corrections to the partition function arising from world-sheet instantons [54]. The results for the instanton partition function of the $\mathcal{N} = 2^*$ theory in the presence of surface operators obtained in [55] are equivalent to our results in (3.18).

3.2 Map between parameters

One of the key points that needs to be clarified is the map between the microscopic counting parameters q_I which appear in (3.23), and the parameters (q, t_I) which were introduced in section 2 in discussing SU(N) gauge theories with surface operators. To describe this map, we start by rewriting the partition function (3.16) in terms of the total instanton number k and the magnetic fluxes m_I of the gauge groups on the surface operator which are related to the parameters d_I as follows [8, 45]:

$$d_1 = k, \quad d_{I+1} = d_I + m_{I+1}. \quad (3.30)$$

Therefore, instead of summing over $\{d_I\}$ we can sum over k and \vec{m} and find

$$Z_{\text{inst}}[\vec{n}] = \sum_{k, \vec{m}} (q_1 \cdots q_M)^k (q_2 \cdots q_M)^{m_2} (q_3 \cdots q_M)^{m_3} \cdots (q_M)^{m_M} Z_{k, \vec{m}}[\vec{n}] \quad (3.31)$$

Furthermore, if we set

$$\begin{aligned} q_I &= e^{2\pi i(t_I - t_{I+1})} \quad \text{for } I \in \{2, \dots, M-1\}, \\ q_M &= e^{2\pi i(t_M - t_1)} \quad \text{and } q = \prod_{I=1}^M q_I, \end{aligned} \quad (3.32)$$

we easily get

$$Z_{\text{inst}}[\vec{n}] = \sum_{k, \vec{m}} q^k e^{2\pi i \sum_{I=2}^M m_I(t_I - t_1)} Z_{k, \vec{m}} = \sum_{k, \vec{m}} q^k e^{2\pi i \vec{t} \cdot \vec{m}} Z_{k, \vec{m}}[\vec{n}] \quad (3.33)$$

where in the last step we introduced m_1 such that $\sum_I m_I = 0$ (see (2.13)) in order to write the result in a symmetric form. This is precisely the expected expression of the partition function in the presence of a surface operator as shown in (2.20) and justifies the map (3.32) between the parameters of the two descriptions. From (3.33) we see that only differences of the parameters t_I appear in the partition function so that it may be convenient to use as independent parameters q and the $(M - 1)$ variables

$$z_J = t_J - t_1 \quad \text{for } J \in \{2, \dots, M\}. \tag{3.34}$$

This is indeed what we are going to see in the next sections where we will show how to extract relevant information from the instanton partition functions described above.

3.3 Extracting the prepotential and the twisted superpotential

The effective dynamics on the Coulomb branch of the four-dimensional $\mathcal{N} = 2^*$ gauge theory is described by the prepotential \mathcal{F} , while the infrared physics of the two-dimensional theory defined on the world-sheet of the surface operator is governed by the twisted superpotential \mathcal{W} . The non-perturbative terms of both \mathcal{F} and \mathcal{W} can be derived from the instanton partition function previously discussed, by considering its behavior for small deformation parameters ϵ_1 and ϵ_2 and, in particular, in the so-called Nekrasov-Shatashvili (NS) limit [51].

To make precise contact with the gauge theory quantities, we set

$$\epsilon_4 = -m - \frac{\epsilon_1}{2} \tag{3.35}$$

where m is the mass of the adjoint hypermultiplet, and then take the limit for small ϵ_1 and ϵ_2 . In this way we find [8]:

$$\log Z_{\text{inst}}[\vec{n}] \simeq -\frac{\mathcal{F}_{\text{inst}}(\epsilon_1)}{\epsilon_1 \epsilon_2} + \frac{\mathcal{W}_{\text{inst}}(\epsilon_1)}{\epsilon_1} + \mathcal{O}(\epsilon_2). \tag{3.36}$$

The two leading singular contributions arise, respectively, from the (regularized) equivariant volume parts coming from the four-dimensional gauge theory and from the two-dimensional degrees of freedom supported on the surface defect D . This can be understood from the fact that, in the Ω -deformed theory, the respective super-volumes are finite and given by [1, 58]:

$$\int_{\mathbb{R}^4_{\epsilon_1, \epsilon_2}} d^4x d^4\theta \longrightarrow \frac{1}{\epsilon_1 \epsilon_2} \quad \text{and} \quad \int_{\mathbb{R}^2_{\epsilon_1}} d^2x d^2\theta \longrightarrow \frac{1}{\epsilon_1}. \tag{3.37}$$

The non-trivial result is that the functions $\mathcal{F}_{\text{inst}}$ and $\mathcal{W}_{\text{inst}}$ defined in this way are analytic in the neighborhood of $\epsilon_1 = 0$. As an illustrative example, we now describe in some detail the $SU(2)$ theory.

SU(2). When the gauge group is $SU(2)$, the only surface operators are of type $\vec{n} = \{1, 1\}$, the Coulomb branch is parameterized by

$$\langle \phi \rangle = \text{diag}(a, -a), \tag{3.38}$$

and the map (3.32) can be written as

$$q_1 = \frac{q}{x}, \quad q_2 = x = e^{2\pi i z} \tag{3.39}$$

where, for later convenience, we have defined $z = (t_2 - t_1)$. Using the results presented in appendix C.1 and their extension to higher orders, it is possible to check that the instanton prepotential arising from (3.36), namely

$$\mathcal{F}_{\text{inst}} = - \lim_{\epsilon_2 \rightarrow 0} \left(\epsilon_1 \epsilon_2 \log Z_{\text{inst}}[1, 1] \right) \tag{3.40}$$

is, as expected, a function only of the instanton counting parameter q and not of x . Expanding in inverse powers of a , we have

$$\mathcal{F}_{\text{inst}} = \sum_{\ell=1}^{\infty} f_{\ell}^{\text{inst}} \tag{3.41}$$

where $f_{\ell} \sim a^{2-\ell}$. The first few coefficients of this expansion are

$$\begin{aligned} f_{2\ell+1}^{\text{inst}} &= 0 \quad \text{for } \ell = 0, 1, \dots, \\ f_2^{\text{inst}} &= - \left(m^2 - \frac{\epsilon_1^2}{4} \right) \left(2q + 3q^2 + \frac{8}{3}q^3 + \dots \right), \\ f_4^{\text{inst}} &= \frac{1}{2a^2} \left(m^2 - \frac{\epsilon_1^2}{4} \right)^2 \left(q + 3q^2 + 4q^3 + \dots \right), \\ f_6^{\text{inst}} &= \frac{1}{16a^4} \left(m^2 - \frac{\epsilon_1^2}{4} \right)^2 \left(2\epsilon_1^2 q - 3(4m^2 - 7\epsilon_1^2)q^2 - 8(8m^2 - 9\epsilon_1^2)q^3 + \dots \right). \end{aligned} \tag{3.42}$$

One can check that this precisely agrees with the NS limit of the prepotential derived for example in [21, 22]. This complete match is a strong and non-trivial check on the correctness and consistency of the whole construction.

Let us now consider the non-perturbative superpotential, which according to (3.36) is

$$\mathcal{W}_{\text{inst}} = \lim_{\epsilon_2 \rightarrow 0} \left(\epsilon_1 \log Z_{\text{inst}}[1, 1] + \frac{\mathcal{F}_{\text{inst}}}{\epsilon_2} \right). \tag{3.43}$$

Differently from the prepotential, $\mathcal{W}_{\text{inst}}$ is, as expected, a function both of q and x . If we expand it as

$$\mathcal{W}_{\text{inst}} = \sum_{\ell=1}^{\infty} w_{\ell}^{\text{inst}} \tag{3.44}$$

with $w_{\ell}^{\text{inst}} \sim a^{1-\ell}$, using the results of appendix C.1 we find

$$\begin{aligned} w_1^{\text{inst}} &= - \left(m - \frac{\epsilon_1}{2} \right) \left[\left(x + \frac{x^2}{2} + \frac{x^3}{3} + \frac{x^4}{4} + \dots \right) + \left(\frac{1}{x} + 2 + x + \dots \right) q \right. \\ &\quad \left. + \left(\frac{1}{2x^2} + \frac{1}{x} + 3 + \dots \right) q^2 + \dots \right], \end{aligned} \tag{3.45a}$$

$$w_2^{\text{inst}} = -\frac{1}{a} \left(m^2 - \frac{\epsilon_1^2}{4} \right) \left[\left(\frac{x}{2} + \frac{x^2}{2} + \frac{x^3}{2} + \frac{x^4}{2} + \dots \right) + \left(\frac{x}{2} - \frac{1}{2x} + \dots \right) q - \left(\frac{1}{2x^2} + \frac{1}{2x} + \dots \right) q^2 + \dots \right], \quad (3.45b)$$

$$w_3^{\text{inst}} = -\frac{\epsilon_1}{a^2} \left(m^2 - \frac{\epsilon_1^2}{4} \right) \left[\left(\frac{x}{4} + \frac{x^2}{2} + \frac{3x^3}{4} + x^4 + \dots \right) + \left(\frac{1}{4x} + \frac{x}{4} + \dots \right) q + \left(\frac{1}{2x^2} + \frac{1}{4x} + \dots \right) q^2 + \dots \right], \quad (3.45c)$$

and so on. For later convenience we explicitly write down the logarithmic derivatives with respect to x , namely

$$w_1' = -\left(m - \frac{\epsilon_1}{2} \right) \left[\left(x + x^2 + x^3 + x^4 + \dots \right) - \left(\frac{1}{x} - x + \dots \right) q - \left(\frac{1}{x^2} + \frac{1}{x} + \dots \right) q^2 + \dots \right], \quad (3.46a)$$

$$w_2' = -\frac{1}{a} \left(m^2 - \frac{\epsilon_1^2}{4} \right) \left[\left(\frac{x}{2} + x^2 + \frac{3x^3}{2} + 2x^4 + \dots \right) + \left(\frac{x}{2} + \frac{1}{2x} + \dots \right) q + \left(\frac{1}{x^2} + \frac{1}{2x} + \dots \right) q^2 + \dots \right], \quad (3.46b)$$

$$w_3' = -\frac{\epsilon_1}{a^2} \left(m^2 - \frac{\epsilon_1^2}{4} \right) \left[\left(\frac{x}{4} + x^2 + \frac{9x^3}{4} + 4x^4 + \dots \right) - \left(\frac{1}{4x} - \frac{x}{4} + \dots \right) q - \left(\frac{1}{x^2} + \frac{1}{4x} + \dots \right) q^2 + \dots \right] \quad (3.46c)$$

where $w_\ell' := x \frac{\partial}{\partial x} (w_\ell^{\text{inst}})$. In the coming sections we will show that these expressions are the weak-coupling expansions of combinations of elliptic and quasi-modular forms of the modular group $\text{SL}(2, \mathbb{Z})$.

4 Modular anomaly equation for the twisted superpotential

In [21, 22] it has been shown for the $\mathcal{N} = 2^* \text{SU}(2)$ theory that the instanton expansions of the prepotential coefficients (3.42) can be resummed in terms of (quasi-) modular forms of the duality group $\text{SL}(2, \mathbb{Z})$ and that the behavior under S-duality severely constrains the prepotential \mathcal{F} which must satisfy a modular anomaly equation. This analysis has been later extended to $\mathcal{N} = 2^*$ theories with arbitrary classical or exceptional gauge groups [34, 41, 42], and also to $\mathcal{N} = 2$ SQCD theories with fundamental matter [38, 39]. In this section we use a similar approach to study how S-duality constrains the form of the twisted superpotential \mathcal{W} .

For simplicity and without loss of generality, in the following we consider a full surface operator of type $\vec{n} = \{1, 1, \dots, 1\}$ with electro-magnetic parameters $\vec{t} = \{t_1, t_2, \dots, t_N\}$. Indeed, surface operators of other type correspond to the case in which these parameters are not all different from each other and form M distinct sets, namely

$$\vec{t} = \left\{ \underbrace{t_1, \dots, t_1}_{n_1}, \underbrace{t_2, \dots, t_2}_{n_2}, \dots, \underbrace{t_M, \dots, t_M}_{n_M} \right\}. \quad (4.1)$$

Thus they can be simply recovered from the full ones with suitable identifications.

Before analyzing the S-duality constraints it is necessary to take into account the classical and the perturbative 1-loop contributions to the prepotential and the superpotential.

The classical contribution. Introducing the notation $\vec{a} = \{a_1, a_2, \dots, a_N\}$ for the vacuum expectation values, the classical contributions to the prepotential and the superpotential are given respectively by

$$\mathcal{F}_{\text{class}} = \pi i \tau \vec{a} \cdot \vec{a} \tag{4.2}$$

and

$$\mathcal{W}_{\text{class}} = 2\pi i \vec{t} \cdot \vec{a}. \tag{4.3}$$

Note that if we use the tracelessness condition (2.4), $\mathcal{W}_{\text{class}}$ can be rewritten as

$$\mathcal{W}_{\text{class}} = 2\pi i \sum_{I=2}^N z_I a_I \tag{4.4}$$

where z_I is as defined in (3.34).

These classical contributions have very simple behavior under S-duality. Indeed

$$S(\mathcal{F}_{\text{class}}) = -\mathcal{F}_{\text{class}}, \tag{4.5a}$$

$$S(\mathcal{W}_{\text{class}}) = -\mathcal{W}_{\text{class}}. \tag{4.5b}$$

To show these relations one has to use the S-duality rules (2.3) and (2.18), and recall that

$$S(\vec{a}) = \vec{a}_D := \frac{1}{2\pi i} \frac{\partial \mathcal{F}}{\partial \vec{a}} \quad \text{and} \quad S(\vec{a}_D) = -\vec{a}, \tag{4.6}$$

which for the classical prepotential simply yield $S(\vec{a}) = \tau \vec{a}$.

The 1-loop contribution. The 1-loop contribution to the partition function of the Ω -deformed gauge theory in the presence of a full surface operator of type $\{1, 1, \dots, 1\}$ can be written in terms of the function

$$\gamma(x) := \log \Gamma_2(x|\epsilon_1, \epsilon_2) = \frac{d}{ds} \left(\frac{\Lambda^s}{\Gamma(s)} \int_0^\infty dt \frac{t^{s-1} e^{-tx}}{(e^{-\epsilon_1 t} - 1)(e^{-\epsilon_2 t} - 1)} \right) \Big|_{s=0}, \tag{4.7}$$

where Γ_2 is the Barnes double Γ -function and Λ an arbitrary scale. Indeed, as shown for example in [55], the perturbative contribution is

$$\log Z_{\text{pert}}[1, 1, \dots, 1] = \sum_{\substack{u,v=1 \\ u \neq v}}^N \left[\gamma \left(a_{uv} + \left\lceil \frac{v-u}{N} \right\rceil \epsilon_2 \right) - \gamma \left(a_{uv} + m + \frac{\epsilon_1}{2} + \left\lceil \frac{v-u}{N} \right\rceil \epsilon_2 \right) \right] \tag{4.8}$$

where $a_{uv} = a_u - a_v$, and the ceiling function $\lceil y \rceil$ denotes the smallest integer greater than or equal to y . The first term in (4.8) represents the contribution of the vector multiplet, while the second term is the contribution of the massive hypermultiplet. Expanding (4.8)

for small $\epsilon_{1,2}$ and using the same definitions (3.36) used for the instanton part, we obtain the perturbative contributions to the prepotential and the superpotential in the NS limit:

$$\begin{aligned}\mathcal{F}_{\text{pert}} &= - \lim_{\epsilon_2 \rightarrow 0} \left(\epsilon_1 \epsilon_2 \log Z_{\text{pert}}[1, 1, \dots, 1] \right), \\ \mathcal{W}_{\text{pert}} &= \lim_{\epsilon_2 \rightarrow 0} \left(\epsilon_1 \log Z_{\text{pert}}[1, 1, \dots, 1] + \frac{\mathcal{F}_{\text{pert}}}{\epsilon_2} \right).\end{aligned}\tag{4.9}$$

Exploiting the series expansion of the γ -function, one can explicitly compute these expressions and show that $\mathcal{F}_{\text{pert}}$ precisely matches the perturbative prepotential in the NS limit obtained in [34, 41], while the contribution to the superpotential is novel. For example, in the case of the SU(2) theory we obtain

$$\mathcal{F}_{\text{pert}} = \frac{1}{2} \left(m^2 - \frac{\epsilon_1^2}{4} \right) \log \frac{4a^2}{\Lambda^2} - \frac{1}{48a^2} \left(m^2 - \frac{\epsilon_1^2}{4} \right)^2 - \frac{1}{960a^4} \left(m^2 - \frac{\epsilon_1^2}{4} \right)^2 \left(m^2 - \frac{3\epsilon_1^2}{4} \right) + \dots,\tag{4.10a}$$

$$\mathcal{W}_{\text{pert}} = -\frac{1}{4a} \left(m^2 - \frac{\epsilon_1^2}{4} \right) - \frac{1}{96a^3} \left(m^2 - \frac{\epsilon_1^2}{4} \right)^2 - \frac{1}{960a^5} \left(m^2 - \frac{\epsilon_1^2}{4} \right)^2 \left(m^2 - \frac{3\epsilon_1^2}{4} \right) + \dots.\tag{4.10b}$$

Note that, unlike the prepotential, the twisted superpotential has no logarithmic term.⁶ Furthermore, it is interesting to observe that

$$\mathcal{W}_{\text{pert}} = -\frac{1}{4} \frac{\partial \mathcal{F}_{\text{pert}}}{\partial a}.\tag{4.11}$$

4.1 S-duality constraints

We are now in a position to discuss the constraints on the twisted superpotential arising from S-duality. Adding the classical, the perturbative and the instanton terms described in the previous sections, we write the complete prepotential and superpotential in the NS limit as

$$\begin{aligned}\mathcal{F} &= \mathcal{F}_{\text{class}} + \mathcal{F}_{\text{pert}} + \mathcal{F}_{\text{inst}} = \pi i \tau \vec{a} \cdot \vec{a} + \sum_{\ell=1}^{\infty} f_{\ell}(\tau, \vec{a}), \\ \mathcal{W} &= \mathcal{W}_{\text{class}} + \mathcal{W}_{\text{pert}} + \mathcal{W}_{\text{inst}} = 2\pi i \sum_{I=2}^N z_I a_I + \sum_{\ell=1}^{\infty} w_{\ell}(\tau, z_I, \vec{a})\end{aligned}\tag{4.12}$$

where for later convenience, we have kept the classical terms separate. The quantum coefficients f_{ℓ} and w_{ℓ} scale as $a^{2-\ell}$ and $a^{1-\ell}$, respectively, and account for the perturbative and instanton contributions. While f_{ℓ} depend on the coupling constant τ , the superpotential coefficients w_{ℓ} are also functions of the surface operator variables z_I , as we have explicitly seen in the SU(2) theory considered in the previous section.

⁶This fact is due to the superconformal invariance, and is no longer true in the pure $\mathcal{N} = 2$ SU(2) gauge theory, for which we find

$$\mathcal{W}_{\text{pert}} = -\left(2 - 2 \log \frac{2a}{\Lambda} \right) a + \frac{\epsilon_1^2}{24a} - \frac{\epsilon_1^4}{2880a^3} + \frac{\epsilon_1^6}{40320a^5} + \dots$$

The coefficients f_ℓ have been explicitly calculated in terms of quasi-modular forms in [34, 41] and we list the first few of them in appendix D. Their relevant properties can be summarized as follows:

- All f_ℓ with ℓ odd vanish, while those with ℓ even are homogeneous functions of \vec{a} and satisfy the scaling relation⁷

$$f_{2\ell}(\tau, \lambda \vec{a}) = \lambda^{2-2\ell} f_{2\ell}(\tau, \vec{a}). \tag{4.13}$$

Since the prepotential has mass-dimension two, the $f_{2\ell}$ are homogeneous polynomials of degree 2ℓ , in m and ϵ_1 .

- The coefficients $f_{2\ell}$ depend on the coupling constant τ only through the Eisenstein series $E_2(\tau)$, $E_4(\tau)$ and $E_6(\tau)$, and are quasi-modular forms of $\text{SL}(2, \mathbb{Z})$ of weight $2\ell - 2$, such that

$$f_{2\ell}\left(-\frac{1}{\tau}, \vec{a}\right) = \tau^{2\ell-2} f_{2\ell}(\tau, \vec{a}) \Big|_{E_2 \rightarrow E_2 + \delta} \tag{4.14}$$

where $\delta = \frac{6}{\pi i \tau}$. The shift δ in E_2 is due to the fact that the second Eisenstein series is a quasi-modular form with an anomalous modular transformation (see (A.4)).

- The coefficients $f_{2\ell}$ satisfy a modular anomaly equation

$$\frac{\partial f_{2\ell}}{\partial E_2} + \frac{1}{24} \sum_{n=1}^{\ell-1} \frac{\partial f_{2n}}{\partial \vec{a}} \cdot \frac{\partial f_{2\ell-2n}}{\partial \vec{a}} = 0 \tag{4.15}$$

which can be solved iteratively.

Using the above properties, it is possible to show that S-duality acts on the prepotential \mathcal{F} in the NS limit as a Legendre transform [41, 42].

Let us now turn to the twisted superpotential \mathcal{W} . As we have seen in (4.5), S-duality acts very simply at the classical level but some subtleties arise in the quantum theory. We now make a few important points, anticipating some results of the next sections. It turns out that \mathcal{W} receives contributions so that the coefficients w_ℓ do not have a well-defined modular weight. However, these anomalous terms depend only on the coupling constant τ and the vacuum expectation values \vec{a} . In particular, they are independent of the continuous parameters z_I that characterize the surface operator. For this reason it is convenient to consider the z_I derivatives of the superpotential:

$$\mathcal{W}^{(I)} := \frac{1}{2\pi i} \frac{\partial \mathcal{W}}{\partial z_I} = a_I + \sum_{\ell=1}^{\infty} w_\ell^{(I)}(\tau, z_I, \vec{a}) \tag{4.16}$$

where, of course, $w_\ell^{(I)} := \frac{1}{2\pi i} \frac{\partial w_\ell}{\partial z_I}$.

Combining intuition from the classical S-duality transformation (4.5b) with the fact that the z_I -derivative increases the modular weight by one, and introduces an extra factor of $(-\tau)$ under S-duality, we are naturally led to propose that

$$S(\mathcal{W}^{(I)}) = \tau \mathcal{W}^{(I)}. \tag{4.17}$$

⁷To be precise, one should also scale $\Lambda \rightarrow \lambda \Lambda$ in the logarithmic term of f_2 .

This constraint can be solved if we assume that the coefficients $w_\ell^{(I)}$ satisfy the following properties (which are simple generalizations of those satisfied by f_ℓ):

- They are homogeneous functions of \vec{a} and satisfy the scaling relation

$$w_\ell^{(I)}(\tau, z_I, \lambda \vec{a}) = \lambda^{1-\ell} w_\ell^{(I)}(\tau, z_I, \vec{a}). \quad (4.18)$$

Given that the twisted superpotential has mass-dimension one, it follows that $w_\ell^{(I)}$ must be homogeneous polynomials of degree ℓ in m and ϵ_1 .

- The dependence of $w_\ell^{(I)}$ on τ and z_I is only through linear combinations of quasi-modular forms made up with the Eisenstein series and elliptic functions with total weight ℓ , such that

$$w_\ell^{(I)}\left(-\frac{1}{\tau}, -\frac{z_I}{\tau}, \vec{a}\right) = \tau^\ell w_\ell^{(I)}(\tau, z_I, \vec{a}) \Big|_{E_2 \rightarrow E_2 + \delta}. \quad (4.19)$$

We are now ready to discuss how S-duality acts on the superpotential coefficients $w_\ell^{(I)}$. Recalling that

$$S(\vec{a}) = \vec{a}_D := \frac{1}{2\pi i} \frac{\partial \mathcal{F}}{\partial \vec{a}} = \tau \vec{a} + \frac{1}{2\pi i} \frac{\partial f}{\partial \vec{a}} = \tau \left(\vec{a} + \frac{\delta}{12} \frac{\partial f}{\partial \vec{a}} \right) \quad (4.20)$$

where $f = \mathcal{F}_{\text{pert}} + \mathcal{F}_{\text{inst}}$, we have

$$\begin{aligned} S(w_\ell^{(I)}) &= w_\ell^{(I)}\left(-\frac{1}{\tau}, -\frac{z_I}{\tau}, \vec{a}_D\right) = \tau^\ell w_\ell^{(I)}(\tau, z_I, \vec{a}_D) \Big|_{E_2 \rightarrow E_2 + \delta} \\ &= \tau w_\ell^{(I)}\left(\tau, z_I, \vec{a} + \frac{\delta}{12} \frac{\partial f}{\partial \vec{a}}\right) \Big|_{E_2 \rightarrow E_2 + \delta} \end{aligned} \quad (4.21)$$

where in the last step we exploited the scaling behavior (4.18) together with (4.20). Using this result in (4.16) and formally expanding in δ , we obtain

$$\begin{aligned} \frac{1}{\tau} S(\mathcal{W}^{(I)}) &= \mathcal{W}^{(I)}\left(\tau, z_I, \vec{a} + \frac{\delta}{12} \frac{\partial f}{\partial \vec{a}}\right) \Big|_{E_2 \rightarrow E_2 + \delta} \\ &= \mathcal{W}^{(I)} + \delta \left(\frac{\partial \mathcal{W}^{(I)}}{\partial E_2} + \frac{1}{12} \frac{\partial \mathcal{W}^{(I)}}{\partial \vec{a}} \cdot \frac{\partial f}{\partial \vec{a}} \right) + \mathcal{O}(\delta^2). \end{aligned} \quad (4.22)$$

The constraint (4.17) is satisfied if

$$\frac{\partial \mathcal{W}^{(I)}}{\partial E_2} + \frac{1}{12} \frac{\partial \mathcal{W}^{(I)}}{\partial \vec{a}} \cdot \frac{\partial f}{\partial \vec{a}} = 0, \quad (4.23)$$

which also implies the vanishing of all terms of higher order in δ . This modular anomaly equation can be equivalently written as

$$\frac{\partial w_\ell^{(I)}}{\partial E_2} + \frac{1}{12} \sum_{n=0}^{\ell-1} \frac{\partial f_{\ell-n}}{\partial \vec{a}} \cdot \frac{\partial w_n^{(I)}}{\partial \vec{a}} = 0 \quad (4.24)$$

where we have defined $w_0^{(I)} = a_I$.

In the next sections we will solve this modular anomaly equation and determine the superpotential coefficients $w_\ell^{(I)}$ in terms of Eisenstein series and elliptic functions; we will also show that by considering the expansion of these quasi-modular functions we recover precisely all instanton contributions computed using localization, thus providing a very strong and highly non-trivial consistency check on our proposal (4.17) and on our entire construction. Since the explicit results are quite involved in the general case, we will start by discussing the SU(2) theory.

5 Surface operators in $\mathcal{N} = 2^* \text{SU}(2)$ theory

We now consider the simplest $\mathcal{N} = 2^*$ theory with gauge group SU(2) and solve in this case the modular anomaly equation (4.24). A slight modification from the earlier discussion is needed since for SU(2) the Coulomb vacuum expectation value of the adjoint scalar field takes the form $\langle \phi \rangle = \text{diag}(a, -a)$ and the index I used in the previous section only takes one value, namely $I = 2$. Thus we have a single z -parameter, corresponding to the unique surface operator we can have in the theory, and (4.16) becomes

$$\mathcal{W}' := \frac{1}{2\pi i} \frac{\partial \mathcal{W}}{\partial z} = -a + \sum_{\ell=1}^{\infty} w'_\ell \tag{5.1}$$

with $w'_\ell := \frac{1}{2\pi i} \frac{\partial w_\ell}{\partial z}$, while the recurrence relation (4.24) becomes

$$\frac{\partial w'_\ell}{\partial E_2} + \frac{1}{24} \sum_{n=0}^{\ell-1} \frac{\partial f_{\ell-n}}{\partial a} \frac{\partial w'_n}{\partial a} = 0 \tag{5.2}$$

with the initial condition $w'_0 = -a$. The coefficient w_1 and its z -derivative w'_1 do not depend on a and are therefore irrelevant for the IR dynamics on the surface operator. Moreover, w'_1 drops out of the anomaly equation and plays no role in determining w'_ℓ for higher values of ℓ . Nevertheless, for completeness, we observe that if we use the elliptic function

$$h_1(z|\tau) = \frac{1}{2\pi i} \frac{\partial}{\partial z} \log \theta_1(z|\tau) \tag{5.3}$$

where $\theta_1(z|\tau)$ is the first Jacobi θ -function, and exploit the expansion reported in (A.16), comparing with the instanton expansion (3.46a) obtained from localization, we are immediately led to,

$$w'_1 = \left(m - \frac{\epsilon_1}{2}\right) \left(h_1 + \frac{1}{2}\right). \tag{5.4}$$

By expanding h_1 to higher orders one can “predict” all higher instanton contributions to w'_1 . We have checked that these predictions perfectly match the explicit results obtained from localization methods involving Young tableaux with up to six boxes.

The first case in which the modular anomaly equation (5.2) shows its power is the case $\ell = 2$. Recalling that the prepotential coefficients f_n with n odd vanish, we have

$$\frac{\partial w'_2}{\partial E_2} + \frac{1}{24} \frac{\partial f_2}{\partial a} \frac{\partial w'_0}{\partial a} = 0. \tag{5.5}$$

Using the initial condition $w'_0 = -a$, substituting the exact expression for f_2 given in (D.1) and then integrating, we get

$$w'_2 = \frac{1}{24a} \left(m^2 - \frac{\epsilon_1^2}{4} \right) (E_2 + \text{modular term}). \tag{5.6}$$

At this juncture, it is important to observe that the elliptic and modular forms of $SL(2, \mathbb{Z})$, which are allowed to appear in the superpotential coefficients, are polynomials in the ring generated by the Weierstraß function $\wp(z|\tau)$ and its z -derivative $\wp'(z|\tau)$, and by the Eisenstein series E_4 and E_6 . These basis elements have weights 2, 3, 4 and 6 respectively. We refer to appendix A for a collection of useful formulas for these elliptic and modular forms and for their perturbative expansions. Since w'_2 must have weight 2, the modular term in (5.6) is restricted to be proportional to the Weierstraß function, namely

$$w'_2 = \frac{1}{24a} \left(m^2 - \frac{\epsilon_1^2}{4} \right) \left(E_2 + \alpha \frac{\wp}{4\pi^2} \right) \tag{5.7}$$

where α is a constant. Therefore our proposal works only if by fixing a *single* parameter α we can match *all* the microscopic contributions to w'_2 computed in the previous sections. Given the many constraints that this requirement puts, it is not at all obvious that it works. But actually it does! Indeed, using the expansions of E_2 and $\tilde{\wp} = \frac{\wp}{4\pi^2}$ given in (A.2) and (A.17) respectively, and comparing with (3.46b), one finds a perfect match if $\alpha = 12$. Thus, the *exact* expression of w'_2 is

$$w'_2 = \frac{1}{24a} \left(m^2 - \frac{\epsilon_1^2}{4} \right) (E_2 + 12 \tilde{\wp}). \tag{5.8}$$

We have checked up to order six that the all instanton corrections predicted by this formula completely agree with the microscopic results obtained from localization.

Let us now consider the modular anomaly equation (5.2) for $\ell = 3$. In this case since w'_1 is a -independent and the coefficients f_n with n odd vanish, we simply have

$$\frac{\partial w'_3}{\partial E_2} = 0 \tag{5.9}$$

According to our proposal, w'_3 must be an elliptic function with modular weight 3, and in view of (5.9), the only candidate is the derivative of the Weierstraß function \wp' . By comparing the expansion (A.18) with the semi-classical results (3.46c) we find a perfect match and obtain

$$w'_3 = \frac{\epsilon_1}{4a^2} \left(m^2 - \frac{\epsilon_1^2}{4} \right) \tilde{\wp}'. \tag{5.10}$$

Again we have checked that the higher order instanton corrections predicted by this formula agree with the localization results up to order six.

A similar analysis can be done for higher values of ℓ without difficulty. Obtaining the anomalous behavior by integrating the modular anomaly equation, and fixing the coefficients of the modular terms by comparing with the localization results, after a bit of

elementary algebra, we get

$$\begin{aligned}
 w'_4 &= \frac{1}{1152a^3} \left(m^2 - \frac{\epsilon_1^2}{4} \right) \left[\left(m^2 - \frac{\epsilon_1^2}{4} \right) (2E_2^2 - E_4 + 24E_2 \tilde{\varphi} + 144\tilde{\varphi}^2) + 6\epsilon_1^2 (E_4 - 144\tilde{\varphi}^2) \right], \\
 w'_5 &= \frac{\epsilon_1}{48a^4} \left(m^2 - \frac{\epsilon_1^2}{4} \right) \left[\left(m^2 - \frac{\epsilon_1^2}{4} \right) (E_2 + 12\tilde{\varphi}) \tilde{\varphi}' - 36\epsilon_1^2 \tilde{\varphi} \tilde{\varphi}' \right], \\
 w'_6 &= \frac{1}{138240a^5} \left(m^2 - \frac{\epsilon_1^2}{4} \right) \left[\left(m^2 - \frac{\epsilon_1^2}{4} \right)^2 (20E_2^3 - 11E_2E_4 - 4E_6 + 240E_2^2 \tilde{\varphi} - 60E_4 \tilde{\varphi} \right. \\
 &\quad \left. + 2160E_2 \tilde{\varphi}^2 + 8640\tilde{\varphi}^3) + 2 \left(m^2 - \frac{\epsilon_1^2}{4} \right) \epsilon_1^2 (39E_2E_4 + 56E_6 + 1440E_4 \tilde{\varphi} \right. \\
 &\quad \left. - 6480E_2 \tilde{\varphi}^2 - 120960\tilde{\varphi}^3) - 240\epsilon_1^4 (E_6 + 27E_4 \tilde{\varphi} - 2160\tilde{\varphi}^3) \right],
 \end{aligned} \tag{5.11}$$

and so on. The complete agreement with the microscopic localization results of the above expressions provides very strong and highly non-trivial evidence for the validity of the modular anomaly equation and the S-duality properties of the superpotential, and hence of our entire construction.

Exploiting the properties of the function h_1 defined in (5.3) and its relation with the Weierstraß function (see appendix A), it is possible to rewrite the above expressions as total z -derivatives. Indeed, we find

$$\begin{aligned}
 w'_2 &= \frac{1}{2a} \left(m^2 - \frac{\epsilon_1^2}{4} \right) h'_1, & w'_3 &= \frac{\epsilon_1}{4a^2} \left(m^2 - \frac{\epsilon_1^2}{4} \right) h''_1, \\
 w'_4 &= \frac{1}{48a^3} \left(m^2 - \frac{\epsilon_1^2}{4} \right) \left[\left(m^2 - \frac{\epsilon_1^2}{4} \right) (E_2 h_1 - h''_1) + 6\epsilon_1^2 h''_1 \right]', \\
 w'_5 &= \frac{\epsilon_1}{8a^4} \left(m^2 - \frac{\epsilon_1^2}{4} \right) \left[\left(m^2 - \frac{\epsilon_1^2}{4} \right) (h'_1)^2 + \frac{\epsilon_1^2}{2} (E_2 - 6h'_1) h'_1 \right]'.
 \end{aligned} \tag{5.12}$$

We have checked that the same is also true for w'_6 (and for a few higher coefficients as well), which however we do not write explicitly for brevity. Of course this is to be expected since they are the coefficients of the expansion of the derivative of the superpotential. The latter can then be simply obtained by integrating with respect to z and fixing the integration constants by comparing with the explicit localization results. In this way we obtain⁸

$$\mathcal{W} = -2\pi i z a + \sum_n w_n \tag{5.13}$$

with

$$\begin{aligned}
 w_2 &= \frac{1}{2a} \left(m^2 - \frac{\epsilon_1^2}{4} \right) h_1, & w_3 &= \frac{\epsilon_1}{4a^2} \left(m^2 - \frac{\epsilon_1^2}{4} \right) h'_1, \\
 w_4 &= \frac{1}{48a^3} \left(m^2 - \frac{\epsilon_1^2}{4} \right) \left[\left(m^2 - \frac{\epsilon_1^2}{4} \right) (E_2 h_1 - h''_1) + 6\epsilon_1^2 h''_1 + \frac{1}{2} \left(m^2 - \frac{\epsilon_1^2}{4} \right) (E_2 - 1) \right], \\
 w_5 &= \frac{\epsilon_1}{8a^4} \left(m^2 - \frac{\epsilon_1^2}{4} \right) \left[\left(m^2 - \frac{\epsilon_1^2}{4} \right) (h'_1)^2 + \frac{\epsilon_1^2}{2} (E_2 - 6h'_1) h'_1 + \frac{1}{96} \left(m^2 - \frac{9\epsilon_1^2}{4} \right) (E_2^2 - E_4) \right],
 \end{aligned} \tag{5.14}$$

⁸We neglect the a -independent terms originating from (5.4) since they are irrelevant for the infrared dynamics on the defect.

and so on. Note that, as anticipated in the previous section, the coefficients w_n do not have a homogeneous modular weight.

5.1 Relation to CFT results

So far we have studied the twisted superpotential and its z -derivative as semi-classical expansions for large a . However, we can also arrange these expansions in terms of the deformation parameter ϵ_1 . For example, using the results in (5.8), (5.10) and (5.11), we obtain

$$\mathcal{W}' = -a + \sum_{n=0}^{\infty} \epsilon_1^n \mathcal{W}'_n \tag{5.15}$$

where

$$\begin{aligned} \mathcal{W}'_0 &= \frac{m^2}{24a} (E_2 + 12\tilde{\varphi}) + \frac{m^4}{1152a^3} (2E_2^2 - E_4 + 24E_2\tilde{\varphi} + 144\tilde{\varphi}^2) + \frac{m^6}{138240a^5} (20E_2^3 \\ &\quad - 11E_2E_4 - 4E_6 + 240E_2^2\tilde{\varphi} - 60E_4\tilde{\varphi} + 2160E_2\tilde{\varphi}^2 + 8640\tilde{\varphi}^3) + \mathcal{O}(a^{-7}), \\ \mathcal{W}'_1 &= \frac{m^2}{4a^2} \tilde{\varphi}' + \frac{m^4}{48a^4} (E_2 + 12\tilde{\varphi})\tilde{\varphi}' + \mathcal{O}(a^{-6}), \\ \mathcal{W}'_2 &= -\frac{1}{96a} (E_2 + 12\tilde{\varphi}) - \frac{m^2}{2304a^3} (2E_2^2 - 13E_4 + 24E_2\tilde{\varphi} + 1872\tilde{\varphi}^2) \\ &\quad - \frac{m^4}{110592a^5} (12E_2^3 - 69E_2E_4 - 92E_6 + 144E_2^2\tilde{\varphi} - 2340E_4\tilde{\varphi} \\ &\quad\quad\quad + 11664E_2\tilde{\varphi}^2 + 198720\tilde{\varphi}^3) + \mathcal{O}(a^{-7}), \\ \mathcal{W}'_3 &= -\frac{1}{16a^2} \tilde{\varphi}' - \frac{m^2}{96a^4} (E_2 + 84\tilde{\varphi})\tilde{\varphi}' + \mathcal{O}(a^{-6}), \end{aligned} \tag{5.16}$$

and so on. Quite remarkably, up to a sign flip $a \rightarrow -a$, these expressions precisely coincide with the results obtained in [35] from the null-vector decoupling equation for the toroidal 1-point conformal block in the Liouville theory.

We would like to elaborate a bit on this match. Let us first recall that in the so-called AGT correspondence [9] the toroidal 1-point conformal block of a Virasoro primary field V in the Liouville theory is related to the Nekrasov partition function of the $\mathcal{N} = 2^*$ SU(2) gauge theory. In [11] it was shown that the insertion of the degenerate null-vector $V_{2,1}$ in the Liouville conformal block corresponds to the partition function of the SU(2) theory in the presence of a surface operator. In the semi-classical limit of the Liouville theory (which corresponds to the NS limit $\epsilon_2 \rightarrow 0$), one has [11, 35]

$$\langle V(0)V_{2,1}(z) \rangle_{\text{torus}} \simeq \mathcal{N} \exp \left(-\frac{\mathcal{F}}{\epsilon_1\epsilon_2} + \frac{\mathcal{W}(z)}{\epsilon_1} + \dots \right), \tag{5.17}$$

where \mathcal{N} is a suitable normalization factor. In [35] the null-vector decoupling equation satisfied by the degenerate conformal block was used to explicitly calculate the prepotential \mathcal{F} and the z -derivative of the twisted effective superpotential \mathcal{W}' for the $\mathcal{N} = 2^*$ SU(2)

theory, which fully agrees with the one we have obtained using the modular anomaly equation and localization methods. It is important to keep in mind that the insertion of the degenerate field $V_{2,1}$ in the Liouville theory corresponds to the insertion of a surface operator of codimension-4 in the six-dimensional $(2,0)$ theory. In the brane picture, this defect corresponds to an M2 brane ending on the M5 branes that wrap a Riemann surface and support the gauge theory in four dimensions. On the other hand, as explained in the introduction, the results we have obtained using the orbifold construction and localization pertain to a surface operator of codimension-2 in the six dimensional theory, corresponding to an M5' intersecting the original M5 branes. The equality between our results and those of [35] supports the proposal of a duality between the two types of surface operators in [52]. This also supports the conjecture of [59], based on [10, 60, 61], that in the presence of simple surface operators the instanton partition function is insensitive to whether they are realized as codimension-2 or codimension-4 operators. In section 7.1 we will comment on such relations in the case of higher rank gauge groups and will also make contact with the results for the twisted chiral rings when the surface defect is realized by coupling two-dimensional sigma-models to pure $\mathcal{N} = 2$ SU(N) gauge theory.

6 Surface operators in $\mathcal{N} = 2^* \text{SU}(N)$ theories

We now generalize the previous analysis to SU(N) gauge groups. As discussed in section 2, in the higher rank cases there are many types of surface operators corresponding to the different partitions of N . We start our discussion from simple surface operators of type $\{1, (N - 1)\}$.

6.1 Simple surface operators

In the case of the simple partition $\{1, (N - 1)\}$, the vector \vec{t} of the electro-magnetic parameters characterizing the surface operator takes the form

$$\vec{t} = \left\{ t_1, \underbrace{t_2, \dots, t_2}_{N-1} \right\}. \tag{6.1}$$

Correspondingly, the classical contribution to the twisted effective superpotential becomes

$$\mathcal{W}_{\text{class}} = 2\pi i \vec{t} \cdot \vec{a} = 2\pi i \left(a_1 t_1 + t_2 \sum_{i=2}^N a_i \right) = -2\pi i z a_1 \tag{6.2}$$

where we have used the tracelessness condition on the vacuum expectation values and, according to (3.34), have defined $z = t_2 - t_1$.

When quantum corrections are included, one finds that the coefficients w'_ℓ of the z -derivative of the superpotential satisfy the modular anomaly equation (4.24). The solution of this equation proceeds along the same lines as in the SU(2) case, although new structures, involving the differences $a_{ij} = a_i - a_j$, appear. We omit details of the calculations and merely present the results. As for the SU(2) theory, the coefficients can be compactly written in terms of modular and elliptic functions, particularly the second Eisenstein series and

the function h_1 defined in (5.3). For clarity, and also for later convenience, we indicate the dependence on z but understand the dependence on τ in h_1 . The first few coefficients w'_ℓ are

$$w'_2 = \left(m^2 - \frac{\epsilon_1^2}{4}\right) \sum_{j=2}^N \frac{h'_1(z)}{a_{1j}}, \tag{6.3a}$$

$$w'_3 = \epsilon_1 \left(m^2 - \frac{\epsilon_1^2}{4}\right) \sum_{j=2}^N \frac{h''_1(z)}{a_{1j}^2} + \frac{1}{2} \left(m^2 - \frac{\epsilon_1^2}{4}\right) \left(m + \frac{\epsilon_1}{2}\right) \sum_{j \neq k=2}^N \frac{h''_1(z)}{a_{1j} a_{1k}}, \tag{6.3b}$$

$$\begin{aligned} w'_4 &= \frac{1}{6} \left(m^2 - \frac{\epsilon_1^2}{4}\right) \left[\left(m^2 - \frac{\epsilon_1^2}{4}\right) (E_2 h'_1(z) - h'''_1(z)) + 6 \epsilon_1^2 h'''_1(z) \right] \sum_{j=2}^N \frac{1}{a_{1j}^3} \\ &+ \epsilon_1 \left(m^2 - \frac{\epsilon_1^2}{4}\right) \left(m + \frac{\epsilon_1}{2}\right) \sum_{j \neq k=2}^N \frac{h'''_1(z)}{a_{1j}^2 a_{1k}} \\ &+ \frac{1}{6} \left(m^2 - \frac{\epsilon_1^2}{4}\right) \left(m + \frac{\epsilon_1}{2}\right)^2 \sum_{j \neq k \neq \ell=2}^N \frac{h'''_1(z)}{a_{1j} a_{1k} a_{1\ell}}, \end{aligned} \tag{6.3c}$$

and so on. We have explicitly checked the above formulas against localization results up to SU(7) finding complete agreement. It is easy to realize that for $N = 2$ only the highest order poles contribute and the corresponding expressions precisely coincide with the results in the previous section. In the higher rank cases, there are also contributions from structures with lesser order poles that are made possible because of the larger number of Coulomb parameters. Furthermore, we observe that there is no pole when a_j approaches a_k with $j, k = 2, \dots, N$.

It is interesting to observe that the above expressions can be rewritten in a suggestive form using the root system Φ of SU(N). The key observation is that using the vector \vec{t} defined in (6.1) we can select a subset of roots $\Psi \subset \Phi$ such that their scalar products with the vector \vec{a} of the vacuum expectation values produce exactly all the factors of a_{1j} in the denominators of (6.3). Defining

$$\Psi = \{ \vec{\alpha} \in \Phi \mid \vec{\alpha} \cdot \vec{t} + z = 0 \}, \tag{6.4}$$

one can verify that for any $\vec{\alpha} \in \Psi$, the scalar product $\vec{\alpha} \cdot \vec{a}$ is of the form a_{1j} . Therefore, w'_2 in (6.3a) can be written as

$$w'_2 = \left(m^2 - \frac{\epsilon_1^2}{4}\right) \sum_{\vec{\alpha} \in \Psi} \frac{h'_1(-\vec{\alpha} \cdot \vec{t})}{\vec{\alpha} \cdot \vec{a}} = \left(m^2 - \frac{\epsilon_1^2}{4}\right) \sum_{\vec{\alpha} \in \Psi} \frac{h'_1(\vec{\alpha} \cdot \vec{t})}{\vec{\alpha} \cdot \vec{a}} \tag{6.5}$$

where in the last step we used the fact that h'_1 is an even function. Similarly the other coefficients in (6.3) can also be rewritten using the roots of SU(N). Indeed, introducing the subsets of Ψ defined as⁹

$$\begin{aligned} \Psi(\vec{\alpha}) &= \{ \vec{\beta} \in \Psi \mid \vec{\alpha} \cdot \vec{\beta} = 1 \}, \\ \Psi(\vec{\alpha}, \vec{\beta}) &= \{ \vec{\gamma} \in \Psi \mid \vec{\alpha} \cdot \vec{\gamma} = \vec{\beta} \cdot \vec{\gamma} = 1 \}, \end{aligned} \tag{6.6}$$

⁹These definitions are analogous to the ones used in [41, 42] to define the root lattice sums appearing in the prepotential; see also (D.7).

we find that w'_3 in (6.3b) becomes

$$\begin{aligned}
 w'_3 &= -\epsilon_1 \left(m^2 - \frac{\epsilon_1^2}{4} \right) \sum_{\vec{\alpha} \in \Psi} \frac{h''_1(\vec{\alpha} \cdot \vec{t})}{(\vec{\alpha} \cdot \vec{a})^2} \\
 &\quad - \frac{1}{2} \left(m^2 - \frac{\epsilon_1^2}{4} \right) \left(m + \frac{\epsilon_1}{2} \right) \sum_{\vec{\alpha} \in \Psi} \sum_{\vec{\beta} \in \Psi(\vec{\alpha})} \frac{h''_1(\vec{\alpha} \cdot \vec{t})}{(\vec{\alpha} \cdot \vec{a})(\vec{\beta} \cdot \vec{a})},
 \end{aligned} \tag{6.7}$$

while w'_4 in (6.3c) is

$$\begin{aligned}
 w'_4 &= \frac{1}{6} \left(m^2 - \frac{\epsilon_1^2}{4} \right) \left[\left(m^2 - \frac{\epsilon_1^2}{4} \right) \sum_{\vec{\alpha} \in \Psi} \frac{E_2 h'_1(\vec{\alpha} \cdot \vec{t}) - h'''_1(\vec{\alpha} \cdot \vec{t})}{(\vec{\alpha} \cdot \vec{a})^3} + 6 \epsilon_1^2 \sum_{\vec{\alpha} \in \Psi} \frac{h'''_1(\vec{\alpha} \cdot \vec{t})}{(\vec{\alpha} \cdot \vec{a})^3} \right] \\
 &\quad + \epsilon_1 \left(m^2 - \frac{\epsilon_1^2}{4} \right) \left(m + \frac{\epsilon_1}{2} \right) \sum_{\vec{\alpha} \in \Psi} \sum_{\vec{\beta} \in \Psi(\vec{\alpha})} \frac{h'''_1(\vec{\alpha} \cdot \vec{t})}{(\vec{\alpha} \cdot \vec{a})^2 (\vec{\beta} \cdot \vec{a})} \\
 &\quad + \frac{1}{4} \left(m^2 - \frac{\epsilon_1^2}{4} \right) \left(m + \frac{\epsilon_1}{2} \right)^2 \left[\sum_{\vec{\alpha} \in \Psi} \sum_{\vec{\beta} \neq \vec{\gamma} \in \Psi(\vec{\alpha})} \frac{h'''_1(\vec{\alpha} \cdot \vec{t})}{(\vec{\alpha} \cdot \vec{a})(\vec{\beta} \cdot \vec{a})(\vec{\gamma} \cdot \vec{a})} \right. \\
 &\quad \quad \left. - \frac{1}{3} \sum_{\vec{\alpha} \in \Psi} \sum_{\vec{\beta} \in \Psi(\vec{\alpha})} \sum_{\vec{\gamma} \in \Psi(\vec{\alpha}, \vec{\beta})} \frac{h'''_1(\vec{\alpha} \cdot \vec{t})}{(\vec{\alpha} \cdot \vec{a})(\vec{\beta} \cdot \vec{a})(\vec{\gamma} \cdot \vec{a})} \right].
 \end{aligned} \tag{6.8}$$

We observe that the two sums in the last two lines of (6.8) are actually equal to each other and exactly reproduce the last line of (6.3c). However, for different sets of roots the two sums are different and lead to different structures. Thus, for reasons that will soon become clear, we have kept them separate even in this case.

6.2 Surface operators of type $\{p, N - p\}$

We now discuss a generalization of the simple surface operator in which we still have a single complex variable z as before, but the type is given by the following vector

$$\vec{t} = \left\{ \underbrace{t_1, \dots, t_1}_p, \underbrace{t_2, \dots, t_2}_{N-p} \right\}. \tag{6.9}$$

In this case, using the tracelessness condition on the vacuum expectation values, the classical contribution to the superpotential is

$$\mathcal{W}_{\text{class}} = 2\pi i \left(t_1 \sum_{i=1}^p a_i + t_2 \sum_{j=p+1}^N a_j \right) = -2\pi i z \sum_{i=1}^p a_i \tag{6.10}$$

where again we have defined $z = t_2 - t_1$.

It turns out that the quantum corrections to the z -derivatives of the superpotential are given exactly by the *same* formulas (6.5), (6.7) and (6.8) in which the only difference is in the subsets of the root system Φ that have to be considered in the lattice sums. These subsets are still defined as in (6.4) and (6.6) but with the vector \vec{t} given by (6.9). We observe that in this case the two last sums in (6.8) are different. We have verified these

formulas against the localization results up to $SU(7)$ finding perfect agreement. The fact that the superpotential coefficients can be formally written in the same way for all unitary groups and for all types with two entries, suggests that probably universal formulas should exist for surface operators with more than two distinct entries in the \vec{t} -vector. This is indeed what happens as we will show in the next subsection.

6.3 Surface operators of general type

A surface operator of general type corresponds to splitting the $SU(N)$ gauge group as in (2.10) which leads to the following partition of the Coulomb parameters

$$\vec{a} = \left\{ \underbrace{a_1, \dots, a_{n_1}}_{n_1}, \underbrace{a_{n_1+1}, \dots, a_{n_1+n_2}}_{n_2}, \dots, \underbrace{a_{N-n_M+1}, \dots, a_N}_{n_M} \right\}, \quad (6.11)$$

and to the following \vec{t} -vector

$$\vec{t} = \left\{ \underbrace{t_1, \dots, t_1}_{n_1}, \underbrace{t_2, \dots, t_2}_{n_2}, \dots, \underbrace{t_M, \dots, t_M}_{n_M} \right\} \quad (6.12)$$

with

$$\sum_{I=1}^M n_I = N. \quad (6.13)$$

In this case we therefore have several variables z_I defined as in (3.34), and several combinations of elliptic functions evaluated at different points. However, if we use the root system Φ of $SU(N)$ the structure of the superpotential coefficients is very similar to what we have seen before in the simplest cases. To see this, let us first define the following subsets¹⁰ of Φ :

$$\begin{aligned} \Psi_{IJ} &= \{ \vec{\alpha} \in \Phi \mid \vec{\alpha} \cdot \vec{t} + z_I - z_J = 0 \}, \\ \Psi_{IJ}(\vec{\alpha}) &= \{ \vec{\beta} \in \Psi_{IJ} \mid \vec{\alpha} \cdot \vec{\beta} = 1 \}, \\ \Psi_{IJ}(\vec{\alpha}, \vec{\beta}) &= \{ \vec{\gamma} \in \Psi_{IJ} \mid \vec{\alpha} \cdot \vec{\gamma} = \vec{\beta} \cdot \vec{\gamma} = 1 \} \end{aligned} \quad (6.14)$$

which are obvious generalizations of the definitions (6.4) and (6.6). Then, writing

$$\mathcal{W}^{(I)} = \frac{1}{2\pi i} \frac{\partial \mathcal{W}}{\partial z_I} = a_{I_1} + \dots + a_{I_{n_I}} + \sum_{\ell} w_{\ell}^{(I)}, \quad (6.15)$$

for $I = 2, \dots, M$, we find that the first few coefficients $w_{\ell}^{(I)}$ are given by

$$w_2^{(I)} = \left(m^2 - \frac{\epsilon_1^2}{4} \right) \sum_{J \neq I} \sum_{\vec{\alpha} \in \Psi_{IJ}} \frac{h_1'(\vec{\alpha} \cdot \vec{t})}{\vec{\alpha} \cdot \vec{a}}, \quad (6.16)$$

$$\begin{aligned} w_3^{(I)} &= -\epsilon_1 \left(m^2 - \frac{\epsilon_1^2}{4} \right) \sum_{J \neq I} \sum_{\vec{\alpha} \in \Psi_{IJ}} \frac{h_1''(\vec{\alpha} \cdot \vec{t})}{(\vec{\alpha} \cdot \vec{a})^2} \\ &\quad - \frac{1}{2} \left(m^2 - \frac{\epsilon_1^2}{4} \right) \left(m + \frac{\epsilon_1}{2} \right) \sum_{J \neq I} \sum_{\vec{\alpha} \in \Psi_{IJ}} \sum_{\vec{\beta} \in \Psi_{IJ}(\vec{\alpha})} \frac{h_1''(\vec{\alpha} \cdot \vec{t})}{(\vec{\alpha} \cdot \vec{a})(\vec{\beta} \cdot \vec{a})}, \end{aligned} \quad (6.17)$$

¹⁰When $J = 1$ one must take $z_1 = 0$.

$$\begin{aligned}
w_4^{(I)} = & \frac{1}{6} \left(m^2 - \frac{\epsilon_1^2}{4} \right) \left[\left(m^2 - \frac{\epsilon_1^2}{4} \right) \sum_{J \neq I} \sum_{\vec{\alpha} \in \Psi_{IJ}} \frac{E_2 h_1'(\vec{\alpha} \cdot \vec{t}) - h_1'''(\vec{\alpha} \cdot \vec{t})}{(\vec{\alpha} \cdot \vec{a})^3} \right. \\
& \left. + 6 \epsilon_1^2 \sum_{J \neq I} \sum_{\vec{\alpha} \in \Psi_{IJ}} \frac{h_1'''(\vec{\alpha} \cdot \vec{t})}{(\vec{\alpha} \cdot \vec{a})^3} \right] \\
& + \epsilon_1 \left(m^2 - \frac{\epsilon_1^2}{4} \right) \left(m + \frac{\epsilon_1}{2} \right) \sum_{J \neq I} \sum_{\vec{\alpha} \in \Psi_{IJ}} \sum_{\vec{\beta} \in \Psi_{IJ}(\vec{\alpha})} \frac{h_1'''(\vec{\alpha} \cdot \vec{t})}{(\vec{\alpha} \cdot \vec{a})^2 (\vec{\beta} \cdot \vec{a})} \\
& + \frac{1}{4} \left(m^2 - \frac{\epsilon_1^2}{4} \right) \left(m + \frac{\epsilon_1}{2} \right)^2 \left[\sum_{J \neq I} \sum_{\vec{\alpha} \in \Psi_{IJ}} \sum_{\vec{\beta} \neq \vec{\gamma} \in \Psi_{IJ}(\vec{\alpha})} \frac{h_1'''(\vec{\alpha} \cdot \vec{t})}{(\vec{\alpha} \cdot \vec{a}) (\vec{\beta} \cdot \vec{a}) (\vec{\gamma} \cdot \vec{a})} \right. \\
& \left. - \frac{1}{3} \sum_{J \neq I} \sum_{\vec{\alpha} \in \Psi_{IJ}} \sum_{\vec{\beta} \in \Psi_{IJ}(\vec{\alpha})} \sum_{\vec{\gamma} \in \Psi_{IJ}(\vec{\alpha}, \vec{\beta})} \frac{h_1'''(\vec{\alpha} \cdot \vec{t})}{(\vec{\alpha} \cdot \vec{a}) (\vec{\beta} \cdot \vec{a}) (\vec{\gamma} \cdot \vec{a})} \right] \\
& + \left(m^2 - \frac{\epsilon_1^2}{4} \right)^2 \sum_{J \neq K \neq I} \sum_{\vec{\alpha} \in \Psi_{IJ}} \sum_{\vec{\beta} \in \Psi_{IK}(\vec{\alpha})} \frac{h_1'(\vec{\alpha} \cdot \vec{t}) h_1'(\vec{\alpha} \cdot \vec{t} - \vec{\beta} \cdot \vec{t})}{(\vec{\alpha} \cdot \vec{a}) (\vec{\beta} \cdot \vec{a}) (\vec{\alpha} \cdot \vec{a} - \vec{\beta} \cdot \vec{a})}
\end{aligned} \tag{6.18}$$

where the summation indices J, K, \dots , take integer values from 1 to M . One can explicitly check that these formulas reduce to those of the previous subsections if $M = 2$ and that no singularity arises when two a 's belonging to the same subgroup in (6.11) approach each other. We have verified these expressions in many cases up to $SU(7)$, always finding agreement with the explicit localization results. Of course it is possible to write down similar expressions for the higher coefficients $w_\ell^{(I)}$, which however become more and more cumbersome as ℓ increases. Given the group theoretic structure of these formulas, it is tempting to speculate that they may be valid for the other simply laced groups of the ADE series as well, similarly to what happens for the analogous expressions of the prepotential coefficients [41]. It would be interesting to verify whether this happens or not.

7 Duality between surface operators

In this section we establish a relation between our localization results and those obtained when the surface defect is realized by coupling two-dimensional sigma-models to the four dimensional gauge theory. When the surface operators are realized in this way, the twisted chiral ring has been independently obtained by studying the two-dimensional (2, 2) theories [62, 63] and related to the Seiberg-Witten geometry of the four dimensional gauge theory [4, 5]. Building on these general results, we extract the semi-classical limit and compare it with the localization answer, finding agreement.

In order to be explicit, we will consider only gauge theories without Ω -deformation, and begin our analysis by first discussing the pure $\mathcal{N} = 2$ theory with gauge group $SU(N)$; in the end we will return to the $\mathcal{N} = 2^*$ theory.

7.1 The pure $\mathcal{N} = 2$ $SU(N)$ theory

The pure $\mathcal{N} = 2$ theory can be obtained by decoupling the adjoint hypermultiplet of the $\mathcal{N} = 2^*$ model. More precisely, this decoupling is carried out by taking the following limit

(see for example [34])

$$m \rightarrow \infty \quad \text{and} \quad q \rightarrow 0 \quad \text{such that} \quad q m^{2N} = (-1)^N \Lambda^{2N} \quad \text{is finite,} \quad (7.1)$$

where Λ is the strong coupling scale of the pure $\mathcal{N} = 2$ theory. In presence of a surface operator, this limit must be combined with a scaling prescription for the continuous variables that characterize the defect. For surface operators of type $\{p, N - p\}$, which possess only one parameter $x = e^{2\pi i z}$, this scaling is

$$m \rightarrow \infty \quad \text{and} \quad x \rightarrow 0 \quad \text{such that} \quad x m^N = (-1)^{p-1} x_0 \Lambda^N \quad \text{is finite.} \quad (7.2)$$

Here $x_0 = e^{2\pi i z_0}$ is the parameter that labels the surface operator in the pure theory à la Gukov-Witten [2–5].

Performing the limits (7.1) and (7.2) on the localization results described in the previous sections, we obtain

$$\mathcal{W}' = \sum_{i=1}^p \mathcal{W}'_i \quad (7.3)$$

where

$$\mathcal{W}'_i = -a_i - \Lambda^N \left(x_0 + \frac{1}{x_0} \right) \prod_{j \neq i}^N \frac{1}{a_{ij}} - \frac{\Lambda^{2N}}{2} \left(x_0^2 + \frac{1}{x_0^2} \right) \frac{\partial}{\partial a_i} \left(\prod_{j \neq i}^N \frac{1}{a_{ij}^2} \right) + \mathcal{O}(\Lambda^{3N}). \quad (7.4)$$

We have explicitly verified this expression in all cases up to $SU(7)$, and for the low rank groups we have also computed the higher instanton corrections.¹¹ With some simple algebra one can check that, up to the order we have worked, \mathcal{W}' is not singular for $a_i \rightarrow a_j$ when both i and j are $\leq p$ or $> p$. Furthermore, one can verify that

$$\sum_{i=1}^N \mathcal{W}'_i = 0 \quad (7.5)$$

as a consequence of the tracelessness condition on the vacuum expectation values.

We now show that this result is completely consistent with the exact twisted chiral ring relation obtained in [5]. For the pure $\mathcal{N} = 2$ $SU(N)$ theory with a surface operator parameterized by x_0 , the twisted chiral ring relation takes the form [5]

$$\mathcal{P}_N(y) - \Lambda^N \left(x_0 + \frac{1}{x_0} \right) = 0 \quad (7.6)$$

with

$$\mathcal{P}_N(y) = \prod_{i=1}^N (y - e_i) \quad (7.7)$$

¹¹For example, for $SU(2)$ and $p = 1$ we find

$$\mathcal{W}'_1 = -a - \frac{\Lambda^2}{2a} \left(x_0 + \frac{1}{x_0} \right) + \frac{\Lambda^4}{8a^3} \left(x_0^2 + \frac{1}{x_0^2} \right) - \frac{\Lambda^6}{16a^5} \left(x_0^3 + x_0 + \frac{1}{x_0} + \frac{1}{x_0^3} \right) + \frac{\Lambda^8}{128a^7} \left(5x_0^4 + 8x_0^2 + \frac{8}{x_0^2} + \frac{5}{x_0^4} \right) + \mathcal{O}(\Lambda^{10})$$

where $a = a_1$.

where e_i are the quantum corrected expectation values of the adjoint scalar. They reduce to a_i in the classical limit $\Lambda \rightarrow 0$ and parameterize the quantum moduli space of the theory. The e_i , which satisfy the tracelessness condition

$$\sum_{i=1}^N e_i = 0, \tag{7.8}$$

were explicitly computed long ago in the 1-instanton approximation in [64, 65] by evaluating the period integrals of the Seiberg-Witten differential and read

$$e_i = a_i - \Lambda^{2N} \frac{\partial}{\partial a_i} \left(\prod_{j \neq i} \frac{1}{a_{ij}^2} \right) + \mathcal{O}(\Lambda^{4N}). \tag{7.9}$$

The higher instanton corrections can be efficiently computed using localization methods [66–69], but their expressions will not be needed in the following.

Inserting (7.9) into (7.7) and systematically working order by order in Λ^N , it is possible to show that the N roots of the chiral ring equation (7.6) are

$$y_i = a_i + \Lambda^N \left(x_0 + \frac{1}{x_0} \right) \prod_{j \neq i} \frac{1}{a_{ij}} + \frac{\Lambda^{2N}}{2} \left(x_0^2 + \frac{1}{x_0^2} \right) \frac{\partial}{\partial a_i} \left(\prod_{j \neq i} \frac{1}{a_{ij}^2} \right) + \mathcal{O}(\Lambda^{3N}) \tag{7.10}$$

for $i = 1, \dots, N$. Comparing with (7.4), we see that, up to an overall sign, y_i coincide with the derivatives of the superpotential \mathcal{W}'_i we obtained from localization. Therefore, we can rewrite the left hand side of (7.6) in a factorized form and get

$$\prod_{i=1}^N (y + \mathcal{W}'_i) - \mathcal{P}_N(y) + \Lambda^N \left(x_0 + \frac{1}{x_0} \right) = 0 \tag{7.11}$$

This shows a perfect match between our localization results and the semi-classical expansion of the chiral ring relation of [5], and provides further non-trivial evidence for the equivalence of the two descriptions. Let us elaborate a bit more on this. According to [5], a surface operator of type $\{p, N-p\}$ has a dual description as a Grassmannian sigma-model coupled to the $SU(N)$ gauge theory, and all information about the twisted chiral ring of the sigma-model is contained in two monic polynomials, Q and \tilde{Q} of degree p and $(N-p)$ respectively, given by

$$Q(y) = \sum_{\ell=0}^p y^\ell \mathcal{X}_{p-\ell}, \quad \tilde{Q}(y) = \sum_{k=0}^{N-p} y^k \tilde{\mathcal{X}}_{N-p-k}. \tag{7.12}$$

with $\mathcal{X}_0 = \tilde{\mathcal{X}}_0 = 1$. Here, \mathcal{X}_ℓ are the twisted chiral ring elements of the Grassmannian sigma-model, and in particular

$$\mathcal{X}_1 = \frac{1}{2\pi i} \frac{\partial \mathcal{W}}{\partial z_0} \tag{7.13}$$

where \mathcal{W} is the superpotential of the surface operator of type $\{p, N-p\}$. The polynomial \tilde{Q} encodes the auxiliary information about the “dual” surface operator obtained by

sending $p \rightarrow (N - p)$. The crucial point is that, according to the proposal of [5], the two polynomials Q and \tilde{Q} satisfy the relation

$$Q(y) \tilde{Q}(y) - \mathcal{P}_N(y) + \Lambda^N \left(x_0 + \frac{1}{x_0} \right) = 0. \tag{7.14}$$

Comparing with (7.11), we are immediately led to the following identifications¹²

$$Q(y) = \prod_{i=1}^p (y + \mathcal{W}'_i), \quad \tilde{Q}(y) = \prod_{j=p+1}^N (y + \mathcal{W}'_j). \tag{7.15}$$

Thus, using (7.13) and (7.3), we find

$$\frac{1}{2\pi i} \frac{\partial \mathcal{W}}{\partial z_0} = \sum_{i=1}^p \mathcal{W}'_i = \mathcal{W}'. \tag{7.16}$$

This equality shows that our localization results for the superpotential of the surface operator of type $\{p, N - p\}$ in the pure $SU(N)$ theory perfectly consistent with the proposal of [5], thus proving the duality between the two descriptions. All this is also a remarkable consistency check of the way in which we have extracted the semi-classical results for the twisted chiral ring of the Grassmannian sigma-model and of the twisted superpotential we have computed.

7.2 The $\mathcal{N} = 2^* SU(N)$ theory

Inspired by the previous outcome, we now analyze the twisted chiral ring relation for simple operators in $\mathcal{N} = 2^*$ theories using the Seiberg-Witten curve and compare it with our localization results for the undeformed theory. To this aim, let us first recall from section 6.1 (see in particular (6.3) with $\epsilon_1 = 0$) that for a simple surface operator corresponding to the following partition of the Coulomb parameters

$$\left\{ a_i, \underbrace{\{a_j \text{ with } j \neq i\}}_{N-1} \right\}, \tag{7.17}$$

the z -derivative of the superpotential is

$$\begin{aligned} \mathcal{W}'_i &= -a_i + m^2 \sum_{j \neq i} \frac{h'_1}{a_{ij}} + \frac{m^3}{2} \sum_{j \neq k \neq i} \frac{h''_1}{a_{ij} a_{ik}} \\ &+ \frac{m^4}{6} \left(\sum_{j \neq i} \frac{E_2 h'_1 - h'''_1}{a_{ij}^3} + \sum_{j \neq k \neq \ell \neq i} \frac{h'''_1}{a_{ij} a_{ik} a_{i\ell}} \right) + \mathcal{O}(m^5). \end{aligned} \tag{7.18}$$

Let us now see how this information can be retrieved from the Seiberg-Witten curve of the $\mathcal{N} = 2^*$ theories. As is well known, in this case there are two possible descriptions (see [44] for a review). The first one, which we call the Donagi-Witten curve [70], is written

¹²We have chosen a specific ordering in which the first p factors correspond to the first p vacuum expectation values a_i ; of course one could as well choose a different ordering by permuting the factors.

naturally in terms of the modular covariant coordinates on moduli space, while the second, which we call the d'Hoker-Phong curve [71], is written naturally in terms of the quantum corrected coordinates on moduli space. As shown in [44], these two descriptions are linearly related to each other with coefficients depending on the second Eisenstein series E_2 .

Since our semi-classical results have been resummed into elliptic and quasi-modular forms, we use the Donagi-Witten curve, which for the $SU(N)$ gauge theory is an N -fold cover of an elliptic curve. It is described by the pair of equations:

$$Y^2 = X^3 - \frac{E_4}{48}X + \frac{E_6}{864}, \quad F_N(y, X, Y) = 0. \quad (7.19)$$

The first equation describes an elliptic curve and thus we can identify (X, Y) with the Weierstraß function and its derivative (see (A.11)). More precisely we have

$$\begin{aligned} X &= -\tilde{\wp} = -h'_1 + \frac{1}{12} E_2, \\ Y &= \frac{1}{2} \tilde{\wp}' = \frac{1}{2} h''_1 \end{aligned} \quad (7.20)$$

The second equation in (7.19) contains a polynomial in y of degree N which encodes the modular covariant coordinates A_k on the Coulomb moduli space of the gauge theory:

$$F_N(y, X, Y) = \sum_{k=0}^N (-1)^k A_k P_{N-k}(y, X, Y) \quad (7.21)$$

where P_k are the modified Donagi-Witten polynomials introduced in [44]. The first few of them are:¹³

$$\begin{aligned} P_0 &= 1, & P_1 &= y, \\ P_2 &= y^2 - m^2 X, & P_3 &= y^3 - 3 y m^2 X + 2 m^3 Y, \\ P_4 &= y^4 - 6 m^2 y^2 X + 8 y m^3 Y - m^4 \left(3 X^2 - \frac{1}{24} E_4 \right). \end{aligned} \quad (7.22)$$

On the other hand, the first few modular covariant coordinates A_k are (see [44]):

$$\begin{aligned} A_2 &= \sum_{i < j} a_i a_j + \frac{m^2}{12} \binom{N}{2} E_2 + \frac{m^4}{288} (E_2^2 - E_4) \sum_{i \neq j} \frac{1}{a_{ij}^2} + \mathcal{O}(m^6), \\ A_3 &= \sum_{i < j < k} a_i a_j a_k - \frac{m^4}{144} (E_2^2 - E_4) \sum_i \sum_{j \neq i} \frac{a_i}{a_{ij}^2} + \mathcal{O}(m^6), \\ A_4 &= \sum_{i < j < k < \ell} a_i a_j a_k a_\ell + \frac{m^2}{12} \binom{N-2}{2} E_2 \sum_{i < j} a_i a_j + \frac{m^4}{48} E_2^2 \\ &\quad + \frac{m^4}{288} (E_2^2 - E_4) \left[\sum_{i < j} \sum_{k \neq \ell} \frac{a_i a_j}{a_{k\ell}^2} + 3 \sum_i \sum_{j \neq i} \frac{a_i^2}{a_{ij}^2} - \binom{N}{2} \right] + \mathcal{O}(m^6), \end{aligned} \quad (7.23)$$

and so on.

¹³The E_4 term in P_4 is one of the modifications which in [44] were found to be necessary and is crucial also here.

We now have all the necessary ingredients to proceed. First of all, using the above expressions and performing the decoupling limits (7.1) and (7.2), one can check that the Donagi-Witten equation $F_N = 0$ reduces to the twisted chiral ring relation (7.6) of the pure theory. Of course this is not a mere coincidence; on the contrary it supports the idea that the Donagi-Witten equation actually encodes also the twisted chiral ring relation of the simple codimension-4 surface operators of the $\mathcal{N} = 2^*$ theories. Secondly, working order by order in the hypermultiplet mass m , one can verify that the N roots of the Donagi-Witten equation are given by

$$y_i = a_i - m^2 \sum_{j \neq i} \frac{h'_1}{a_{ij}} - \frac{m^3}{2} \sum_{j \neq k \neq i} \frac{h''_1}{a_{ij} a_{ik}} - \frac{m^4}{6} \left(\sum_{j \neq i} \frac{E_2 h'_1 - h'''_1}{a_{ij}^3} + \sum_{j \neq k \neq \ell \neq i} \frac{h'''_1}{a_{ij} a_{ik} a_{i\ell}} \right) + \mathcal{O}(m^5). \tag{7.24}$$

Remarkably, this precisely matches, up to an overall sign, the answer (7.18) for the simple codimension-2 surface operator we have obtained using localization. Once again, we have exhibited the equivalence of twisted chiral rings calculated for the two kinds of surface operators. Furthermore, we can rewrite the Donagi-Witten equation in a factorized form as follows

$$\prod_{i=1}^N (y + \mathcal{W}'_i) - F_N(y, X, Y) = 0 \tag{7.25}$$

which is the $\mathcal{N} = 2^*$ equivalent of the pure theory relation (7.11).

At this point one is tempted to proceed as in the pure theory and try to deduce also the superpotential for surface operators of type $\{p, N - p\}$. However, from our explicit localization results we know that in this case \mathcal{W}' is not simply the sum of the superpotentials of type $\{1, N - 1\}$, differently from what happens in the pure theory (see (7.3)). Thus, a naive extension to the $\mathcal{N} = 2^*$ of the proposal of [5] to describe the coupling of a two dimensional Grassmannian sigma-model to the four dimensional gauge theory can not work in this case. This problem as well as the coupling of a flag variety to the $\mathcal{N} = 2^*$ theory, which is relevant for surface operators of general type, remains an open question which we leave to future investigations.

7.3 Some remarks on the results

The result we obtained from the twisted superpotential in the case of simple operators is totally consistent with the proposal given in the literature for simple codimension-4 surface operators labeled by a single continuous parameter z , whose superpotential has been identified with the line integral of the Seiberg-Witten differential of the four-dimensional gauge theory along an open path [11]:

$$\mathcal{W}(z) = \int_{z^*}^z \lambda_{SW} \tag{7.26}$$

where z^* is an arbitrary reference point. Indeed, in the Donagi-Witten variables, the differential is simply $\lambda_{SW}(z) = y(z) dz$. Given that the Donagi-Witten curve is an N -fold

cover of the torus, the twisted superpotential with the classical contribution proportional to a_i can be obtained by solving for $y(z)$ and writing out the solution on the i th branch.

As we have seen in the previous subsection, the general identification in (7.26) works also in the pure $\mathcal{N} = 2$ theory, once the parameters in the Seiberg-Witten differential are rescaled by a factor of Λ^N [5]. This rescaling can be interpreted as a renormalization of the continuous parameter that labels the surface operator [72].

The agreement we find gives further evidence of the duality between defects realized as codimension-2 and codimension-4 operators that we have already discussed in section 5.1, where we showed the equality of the twisted effective superpotential computed in the two approaches for simple defects in the $SU(2)$ theory. We have extended these checks to defects of type $\{p, N - p\}$ in pure $\mathcal{N} = 2$ theories, and to simple defects in $\mathcal{N} = 2^*$ theories with higher rank gauge groups. All these checks support the proposal of [52] based on a “separation of variables” relation.

8 Conclusions

In this paper we have studied the properties of surface operators on the Coulomb branch of the four dimensional $\mathcal{N} = 2^*$ theory with gauge group $SU(N)$ focusing on the superpotential \mathcal{W} . This superpotential, describing the effective two-dimensional dynamics on the defect world-sheet, receives non-perturbative contributions, which we calculated using equivariant localization. Furthermore, exploiting the constraints arising from the non-perturbative $SL(2, \mathbb{Z})$ symmetry, we showed that in a semi-classical regime in which the mass of the adjoint hypermultiplet is much smaller than the classical Coulomb branch parameters, the twisted superpotential satisfies a modular anomaly equation that we solved order by order in the mass expansion.

We would like to remark some interesting properties of our results. If we focus on the derivatives of the superpotential, the coefficients of the various terms in the mass expansion are linear combination of elliptic and quasi-modular forms with a given weight. The explicit expression for the twisted superpotential can be written in a very general and compact form in terms of suitable restricted sums over the root lattice of the gauge algebra.

The match of our localization results with the ones obtained in [5] by studying the coupling with two-dimensional sigma models is a non-trivial check of our methods and provides evidence for the duality between the codimension-2 and codimension-4 surface operators proposed in [52]. Further evidence is given by the match of the twisted superpotentials in the $\mathcal{N} = 2^*$ theory, which we proved for the simple surface operators using the Donagi-Witten curve of the model. A key input for this match is the exact quantum expression of the chiral ring elements calculated using localization [44, 69]. It would be really important to extend the discussion of this duality to more general surface operators described by a generic Levi decomposition.

There are several possible extensions of our work. A very direct one would be to check that the general expression given for the twisted superpotential is actually valid for all simply laced groups, in analogy to what happens for the four-dimensional prepotential. A technically more challenging extension would be to study surface operators for theories

with non-simply laced gauge groups. The prepotential in these cases has been calculated in [42] using localization methods and expressed in terms of modular forms of suitable congruence subgroups of $SL(2, \mathbb{Z})$, and it would be very interesting to similarly calculate the twisted superpotential in a semi-classical expansion.

Another interesting direction would be to study surface operators in SQCD theories. For $SU(N)$ gauge groups, the prepotential as well as the action of S-duality on the infrared variables have been calculated in a special locus of the Coulomb moduli space that has a \mathbb{Z}_N symmetry [38, 39]. Of special importance was the generalized Hecke groups acting on the period integrals and the period matrix of the Seiberg-Witten curve. It would be worthwhile to explore if such groups continue to play a role in determining the twisted superpotential as well.

A related development would be to analyze the higher order terms in the ϵ_2 expansion of the partition function (see (3.36)) and check whether or not they also obey a modular anomaly equation like the prepotential and the superpotential do. This would help us in clarifying the properties of the partition function in the presence of a surface operator in a general Ω background.

There has been a lot of progress in understanding M2 brane surface operators via the $4d/2d$ correspondence. For higher rank theories, explicit results for such surface defects have been obtained in various works including [73–77]. In particular in [75], the partition functions of theories with N_f^2 free hypermultiplets on the deformed 4-sphere in the presence of surface defects have been related to specific conformal blocks in Toda conformal field theories. This has been extended in [76, 77] to study gauge theory partition functions in the presence of intersecting surface defects. It would be interesting to study such configurations directly using localization methods.

Acknowledgments

We would like to thank Dileep Jatkar, Madhusudhan Raman and especially Jan Troost for useful discussions and Matteo Beccaria for comments on the manuscript. The work of M.B. and M.F. is partially supported by the Compagnia di San Paolo contract “MAST: Modern Applications of String Theory” TO-Call3-2012-0088. The work of M.B., M.F. and A.L. is partially supported by the MIUR PRIN Contract 2015MP2CX4 “Non-perturbative Aspects Of Gauge Theories And Strings”.

A Useful formulas for modular forms and elliptic functions

In this appendix we collect some formulas about quasi-modular forms and elliptic functions that are useful to check the statements of the main text.

Eisenstein series. We begin with the Eisenstein series E_{2n} , which admit a Fourier expansion in terms of $q = e^{2\pi i\tau}$ of the form

$$E_{2n} = 1 + \frac{2}{\zeta(1-2n)} \sum_{k=1}^{\infty} \sigma_{2n-1}(k) q^k, \tag{A.1}$$

where $\sigma_p(k)$ is the sum of the p -th powers of the divisors of k . More explicitly we have

$$\begin{aligned}
 E_2 &= 1 - 24 \sum_{k=1}^{\infty} \sigma_1(k) q^k = 1 - 24q - 72q^2 - 96q^3 - 168q^4 + \dots, \\
 E_4 &= 1 + 240 \sum_{k=1}^{\infty} \sigma_3(k) q^k = 1 + 240q + 2160q^2 + 6720q^3 + 17520q^4 + \dots, \\
 E_6 &= 1 - 504 \sum_{k=1}^{\infty} \sigma_5(k) q^k = 1 - 504q - 16632q^2 - 122976q^3 - 532728q^4 + \dots.
 \end{aligned}
 \tag{A.2}$$

Under a modular transformation $\tau \rightarrow \frac{a\tau+b}{c\tau+d}$, with $a, b, c, d \in \mathbb{Z}$ and $ad - bc = 1$, the Eisenstein series transform as

$$E_2 \rightarrow (c\tau + d)^2 E_2 + \frac{6}{\pi i} c(c\tau + d), \quad E_4 \rightarrow (c\tau + d)^4 E_4, \quad E_6 \rightarrow (c\tau + d)^6 E_6.
 \tag{A.3}$$

In particular, under S-duality we have

$$\begin{aligned}
 E_2(\tau) &\rightarrow E_2\left(-\frac{1}{\tau}\right) = \tau^2 (E_2(\tau) + \delta), \\
 E_4(\tau) &\rightarrow E_4\left(-\frac{1}{\tau}\right) = \tau^4 E_4(\tau), \\
 E_6(\tau) &\rightarrow E_6\left(-\frac{1}{\tau}\right) = \tau^6 E_6(\tau)
 \end{aligned}
 \tag{A.4}$$

where $\delta = \frac{6}{\pi i \tau}$.

Elliptic functions. The elliptic functions that are relevant for this paper can all be obtained from the Jacobi θ -function

$$\theta_1(z|\tau) = \sum_{n=-\infty}^{\infty} q^{\frac{1}{2}(n-\frac{1}{2})^2} (-x)^{(n-\frac{1}{2})}
 \tag{A.5}$$

where $x = e^{2\pi iz}$. From θ_1 , we first define the function

$$h_1(z|\tau) = \frac{1}{2\pi i} \frac{\partial}{\partial z} \log \theta_1(z|\tau) = x \frac{\partial}{\partial x} \log \theta_1(z|\tau),
 \tag{A.6}$$

and the Weierstraß \wp -function

$$\wp(z|\tau) = -\frac{\partial^2}{\partial z^2} \log \theta_1(z|\tau) - \frac{\pi^2}{3} E_2(\tau).
 \tag{A.7}$$

In most of our formulas the following rescaled \wp -function appears:

$$\tilde{\wp}(z|\tau) := \frac{\wp(z, \tau)}{4\pi^2} = x \frac{\partial}{\partial x} \left(x \frac{\partial}{\partial x} \log \theta_1(z|\tau) \right) - \frac{1}{12} E_2(\tau),
 \tag{A.8}$$

which we can write also as

$$\tilde{\wp}(z|\tau) = h'_1(z|\tau) - \frac{1}{12} E_2(\tau).
 \tag{A.9}$$

Another relevant elliptic function is the derivative of the Weierstraß function, namely

$$\tilde{\wp}'(z|\tau) := \frac{1}{2\pi i} \frac{\partial}{\partial z} \tilde{\wp}(z|\tau) = x \frac{\partial}{\partial x} \tilde{\wp}(z|\tau) = h_1''(z|\tau). \quad (\text{A.10})$$

The Weierstraß function and its derivative satisfy the equation of an elliptic curve, given by

$$\tilde{\wp}'(z|\tau)^2 + 4\tilde{\wp}(z|\tau)^3 - \frac{E_4}{12}\tilde{\wp}(z|\tau) - \frac{E_6}{216} = 0. \quad (\text{A.11})$$

By differentiating this equation, we obtain

$$\tilde{\wp}''(z|\tau) = -6\tilde{\wp}(z|\tau)^2 + \frac{E_4}{24} \quad (\text{A.12})$$

which, using (A.9) and (A.10), we can rewrite as

$$h_1'''(z|\tau) = -6(h_1'(z|\tau))^2 + E_2 h_1'(z|\tau) - \frac{E_2^2 - E_4}{24}. \quad (\text{A.13})$$

The function h_1 , $\tilde{\wp}$ and $\tilde{\wp}'$ have well-known expansions near the point $z = 0$. However, a different expansion is needed for our purposes, namely the expansion for small q and x . To find such an expansion we observe that q and x variables must be rescaled differently, as is clear from the map (3.32) between the gauge theory parameters and the microscopic counting parameters. In particular for $M = 2$ this map reads (see also (3.39))

$$q = q_1 q_2 \quad , \quad x = q_2, \quad (\text{A.14})$$

so that if the microscopic parameters are all scaled equally as $q_i \rightarrow \lambda q_i$, then the gauge theory parameters scale as

$$q \rightarrow \lambda^2 q \quad x \rightarrow \lambda x. \quad (\text{A.15})$$

With this in mind, we now expand the elliptic functions for small λ and set $\lambda = 1$ in the end, since this is the relevant expansion needed to compare with the instanton calculations. Proceeding in this way, we find¹⁴

$$\begin{aligned} h_1(x|q) &= h_1(\lambda x|\lambda^2 q) \Big|_{\lambda=1} \\ &= \left[-\frac{1}{2} + \lambda \left(\frac{q}{x} - x \right) + \lambda^2 \left(\frac{q^2}{x^2} - x^2 \right) + \lambda^3 \left(\frac{q^3}{x^3} + \frac{q^2}{x} - qx - x^3 \right) \right. \\ &\quad \left. - \lambda^4 x^4 + \lambda^5 \left(\frac{q^3}{x} - q^2 x - x^5 \right) - \lambda^6 (q^2 x^2 + x^6) + \dots \right]_{\lambda=1} \\ &= -\frac{1}{2} - \left(x + x^2 + x^3 + x^4 + x^5 + x^6 + \dots \right) + \left(\frac{1}{x} - x \right) q \\ &\quad + \left(\frac{1}{x^2} + \frac{1}{x} - x - x^2 \right) q^2 + \left(\frac{1}{x^3} + \frac{1}{x} + \dots \right) q^3 + \dots, \end{aligned} \quad (\text{A.16})$$

¹⁴Depending on the context, we denote the arguments of the elliptic functions by either $(z|\tau)$ as we did so far, or by their exponentials $(x|q)$ when the expansions are being used.

$$\begin{aligned}
 \tilde{\wp}(x|q) &= \tilde{\wp}(\lambda x|\lambda^2 q)\Big|_{\lambda=1} \\
 &= \left[-\frac{1}{12} - \lambda\left(\frac{q}{x} + x\right) + \lambda^2\left(-\frac{2q^2}{x^2} + 2q - 2x^2\right) \right. \\
 &\quad \left. - \lambda^3\left(\frac{3q^3}{x^3} + \frac{q^2}{x} + qx + 3x^3\right) + \lambda^4(6q^2 - 4x^4) + \dots \right]_{\lambda=1} \tag{A.17} \\
 &= -\frac{1}{12} - (x + 2x^2 + 3x^3 + 4x^4 + \dots) - \left(\frac{1}{x} - 2 + x\right)q \\
 &\quad - \left(\frac{2}{x^2} + \frac{1}{x} - 6 + \dots\right)q^2 - \frac{3q^3}{x^3} + \dots,
 \end{aligned}$$

$$\begin{aligned}
 \tilde{\wp}'(x|q) &= \tilde{\wp}'(\lambda x|\lambda^2 q)\Big|_{\lambda=1} \\
 &= \left[\lambda\left(\frac{q}{x} - x\right) + \lambda^2\left(\frac{4q^2}{x^2} - 4x^2\right) \right. \\
 &\quad \left. + \lambda^3\left(\frac{9q^3}{x^3} + \frac{q^2}{x} - qx - 9x^3\right) - 16\lambda^4x^4 + \dots \right]_{\lambda=1} \tag{A.18} \\
 &= -(x + 4x^2 + 9x^3 + 16x^4 + \dots) + \left(\frac{1}{x} - x\right)q + \left(\frac{4}{x^2} + \frac{1}{x} + \dots\right)q^2 + \frac{9q^3}{x^3} + \dots.
 \end{aligned}$$

As a consistency check it is possible to verify that, using these expansions and those of the Eisenstein series in (A.2), the elliptic curve equation (A.11) is identically satisfied order by order in λ .

As we have seen in section 2, the modular group acts on $(z|\tau)$ as follows:

$$(z|\tau) \rightarrow \left(\frac{z}{c\tau + d} \Big| \frac{a\tau + b}{c\tau + d} \right) \tag{A.19}$$

with $a, b, c, d \in \mathbb{Z}$ and $ad - bc = 1$. Under such transformations the Weierstraß function and its derivative have, respectively, weight 2 and 3, namely

$$\begin{aligned}
 \wp(z|\tau) &\rightarrow \wp\left(\frac{z}{c\tau + d} \Big| \frac{a\tau + b}{c\tau + d}\right) = (c\tau + d)^2 \wp(z|\tau), \\
 \wp'(z|\tau) &\rightarrow \wp'\left(\frac{z}{c\tau + d} \Big| \frac{a\tau + b}{c\tau + d}\right) = (c\tau + d)^3 \wp'(z|\tau).
 \end{aligned} \tag{A.20}$$

Of course, similar relations hold for the rescaled functions $\tilde{\wp}$ and $\tilde{\wp}'$. In particular, under S-duality we have

$$\begin{aligned}
 \tilde{\wp}(z|\tau) &\rightarrow \tilde{\wp}\left(-\frac{z}{\tau} \Big| -\frac{1}{\tau}\right) = \tau^2 \tilde{\wp}(z|\tau), \\
 \tilde{\wp}'(z|\tau) &\rightarrow \tilde{\wp}'\left(-\frac{z}{\tau} \Big| -\frac{1}{\tau}\right) = -\tau^3 \tilde{\wp}'(z|\tau).
 \end{aligned} \tag{A.21}$$

B Generalized instanton number in the presence of fluxes

In this appendix we calculate the second Chern class of the gauge field in the presence of a surface operator for a generic Lie algebra \mathfrak{g} .

Surface operator Ansatz. A surface operator creates a singularity in the gauge field A . As discussed in the main text, we parametrize the space-time $\mathbb{R}^4 \simeq \mathbb{C}^2$ by two complex variables ($z_1 = \rho e^{i\phi}$, $z_2 = r e^{i\theta}$), and consider a two-dimensional defect D located at $z_2 = 0$ and filling the z_1 -plane. In this set-up, we make the following Ansatz [8]:

$$A = \widehat{A} + g(r) d\theta, \quad (\text{B.1})$$

where \widehat{A} is regular all over \mathbb{R}^4 and $g(r)$ is a \mathfrak{g} -valued function regular when $r \rightarrow 0$. The corresponding field strength is then

$$F := dA - i A \wedge A = \widehat{F} + d(g(r) d\theta) - i d\theta \wedge [g(r), \widehat{A}]. \quad (\text{B.2})$$

From this expression we obtain

$$\begin{aligned} \text{Tr } F \wedge F &= \text{Tr } \widehat{F} \wedge \widehat{F} + 2 \text{Tr} \left(d(g(r) d\theta) \wedge \widehat{F} \right) - 2i \text{Tr} \left(d\theta \wedge [g(r), \widehat{A}] \wedge \widehat{F} \right) \\ &= \text{Tr } \widehat{F} \wedge \widehat{F} + 2 \text{Tr} d(g(r) d\theta \wedge \widehat{F}) + 2 \text{Tr} \left(g(r) d\theta \wedge (d\widehat{F} - i \widehat{A} \wedge \widehat{F} - i \widehat{F} \wedge \widehat{A}) \right). \end{aligned} \quad (\text{B.3})$$

The last term vanishes due to the Bianchi identity, and thus we are left with

$$\text{Tr } F \wedge F = \text{Tr } \widehat{F} \wedge \widehat{F} + 2 \text{Tr} d(g(r) d\theta \wedge \widehat{F}) \quad (\text{B.4})$$

We now assume that the function $g(r)$ has components only along the Cartan directions of \mathfrak{g} , labeled by an index i , such that

$$\lim_{r \rightarrow 0} g_i(r) = -\gamma_i \quad \text{and} \quad \lim_{r \rightarrow \infty} g_i(r) = 0. \quad (\text{B.5})$$

This means that near the defect the gauge connection behaves as

$$A = A_\mu dx^\mu \simeq -\text{diag}(\gamma_1, \dots, \gamma_{\text{rank}(\mathfrak{g})}) d\theta \quad (\text{B.6})$$

for $r \rightarrow 0$. Using this in (B.4), we have

$$\text{Tr } F \wedge F = \text{Tr } \widehat{F} \wedge \widehat{F} + 2 \sum_i d(g_i(r) d\theta \wedge \widehat{F}_i). \quad (\text{B.7})$$

Notice that in the last term we can replace \widehat{F}_i with F_i because the difference lies entirely in the transverse directions of the surface operator and thus does not contribute in the wedge product with $d\theta$. Since the defect D effectively acts as a boundary in \mathbb{R}^4 located at $r = 0$, integrating (B.7) over \mathbb{R}^4 we have

$$\frac{1}{8\pi^2} \int_{\mathbb{R}^4} \text{Tr } F \wedge F = \frac{1}{8\pi^2} \int_{\mathbb{R}^4} \text{Tr } \widehat{F} \wedge \widehat{F} + \sum_i \frac{\gamma_i}{2\pi} \int_D F_i = k + \sum_i \gamma_i m_i. \quad (\text{B.8})$$

Here we have denoted by k the instanton number of the smooth connection \widehat{A} and taken into account a factor of 2π originating from the integration over θ . Finally, we have defined

$$m_i = \frac{1}{2\pi} \int_D F_i. \quad (\text{B.9})$$

These quantities, which we call fluxes, must satisfy a quantization condition that can be understood as follows. All fields of the gauge theory are organized in representations¹⁵ of \mathfrak{g} and, in particular, can be chosen to be eigenstates of the Cartan generators H_i with eigenvalues λ_i . These eigenvalues define a vector $\vec{\lambda} = \{\lambda_i\}$, which is an element of the weight lattice Λ_W of \mathfrak{g} . Let us now consider a gauge transformation in the Cartan subgroup with parameters $\vec{\omega} = \{\omega_i\}$. On a field with weight $\vec{\lambda}$, this transformation simply acts by a phase factor $\exp(i\vec{\omega} \cdot \vec{\lambda})$. From the point of view of the two-dimensional theory on the defect, the Cartan gauge fields A_i must approach a pure-gauge configuration at infinity so that

$$A_i \sim d\omega_i \quad \text{for } \rho \rightarrow \infty, \tag{B.10}$$

with ω_i being a function of ϕ , the polar angle in the z_1 -plane. In this situation, for the corresponding gauge transformation to be single-valued, one finds

$$\vec{\omega}(\phi + 2\pi) \cdot \vec{\lambda} = \vec{\omega}(\phi) \cdot \vec{\lambda} + 2\pi n \tag{B.11}$$

with integer n . In other words $\vec{\omega} \cdot \vec{\lambda}$ must be a map from the circle at infinity S_1^∞ into S_1 with integer winding number n . Given this, we have

$$2\pi m_i = \int_D F_i = \oint_{S_1^\infty} d\omega_i = \omega_i(\phi + 2\pi) - \omega_i(\phi). \tag{B.12}$$

Then, using (B.11), we immediately deduce that

$$\vec{m} \cdot \vec{\lambda} \in \mathbb{Z}. \tag{B.13}$$

For the group $SU(N)$ this condition amounts to say that \vec{m} must belong to the dual of the weight lattice:

$$\vec{m} \in (\Lambda_W)^*. \tag{B.14}$$

The $SU(N)$ case. For $U(N)$ the Cartan generators H_i can be taken as the diagonal $(N \times N)$ matrices with just a single non-zero entry equal to 1 in the i -th place ($i = 1, \dots, N$). The restriction to $SU(N)$ can be obtained by choosing a basis of $(N-1)$ traceless generators, for instance $(H_i - H_{i+1})/\sqrt{2}$. In terms of the standard orthonormal basis $\{\vec{e}_i\}$ of \mathbb{R}^N , the $(N-1)$ simple roots of $SU(N)$ are then $\{(\vec{e}_1 - \vec{e}_2), (\vec{e}_2 - \vec{e}_3), \dots\}$ and the root lattice Λ_R is the \mathbb{Z} -span of these simple roots. Note that Λ_R lies in a codimension-1 subspace orthogonal to $\sum_i \vec{e}_i$, and that the integrality condition for the weights is simply $\vec{\alpha} \cdot \vec{\lambda} \in \mathbb{Z}$ for any root $\vec{\alpha}$. This shows that the weight lattice is the dual of the root lattice, or equivalently that the dual of the weight lattice is the root lattice: $(\Lambda_W)^* = \Lambda_R$. Therefore, the condition (B.14) implies that the flux vector \vec{m} must be of the form

$$\vec{m} = n_1(\vec{e}_1 - \vec{e}_2) + n_2(\vec{e}_2 - \vec{e}_3) + \dots + n_{N-1}(\vec{e}_{N-1} - \vec{e}_N) \quad \text{with } n_i \in \mathbb{Z}. \tag{B.15}$$

This simply corresponds to

$$\vec{m} = \sum_i m_i \vec{e}_i \quad \text{with } m_i \in \mathbb{Z} \quad \text{and} \quad \sum_i m_i = 0. \tag{B.16}$$

The fact that the fluxes m_i are integers (adding up to zero) has been used in the main text.

¹⁵Here for simplicity we consider the gauge group G to be the universal covering group of \mathfrak{g} ; in particular for $\mathfrak{g} = A_{N-1}$, we take $G = SU(N)$.

Generic surface operator. The case in which all the γ_i 's defined in (B.5) are distinct, corresponds to the surface operator of type $[1, 1, \dots, 1]$, also called full surface operator. If instead some of the γ_i 's coincide, the surface operator has a more generic form. Let us consider for example the case in which the $SU(N)$ gauge field at the defect takes the form (see (2.8)):

$$A = A_\mu dx^\mu \simeq -\text{diag} \left(\underbrace{\gamma_1, \dots, \gamma_1}_{n_1}, \underbrace{\gamma_2, \dots, \gamma_2}_{n_2}, \dots, \underbrace{\gamma_M, \dots, \gamma_M}_{n_M} \right) d\theta, \quad (\text{B.17})$$

for $r \rightarrow 0$, which corresponds to splitting the gauge group according to

$$SU(N) \rightarrow S[U(n_1) \times U(n_2) \times \dots \times U(n_M)]. \quad (\text{B.18})$$

The calculation of the second Chern class (B.8) proceeds as before, but the result can be written as follows

$$\frac{1}{8\pi^2} \int_M \text{Tr} F \wedge F = k + \sum_{I=1}^M \gamma_I m_I \quad (\text{B.19})$$

with

$$m_I = \sum_{i=1}^{n_I} m_i = \frac{1}{2\pi} \int_D \sum_{i=1}^{n_I} F_i = \frac{1}{2\pi} \int_D \text{Tr} F_{U(n_I)}. \quad (\text{B.20})$$

Here we see that it is the magnetic flux associated with the $U(1)$ factor in each subgroup $U(n_I)$ that appears in the expression for the generalized instanton number in the presence of magnetic fluxes.

C Ramified instanton moduli and their properties

In this appendix we describe the instanton moduli in the various sectors. Our results are summarized in table 1.

Let us first consider the neutral states of the strings stretching between two D -instantons.

- **$(-1)/(-1)$ strings of type $I-I$:** all moduli of this type transform in the adjoint representation $(\mathbf{d}_I, \bar{\mathbf{d}}_I)$ of $U(d_I)$. A special role is played by the bosonic states created in the Neveu-Schwarz (NS) sector of such strings by the complex oscillator ψ^v in the last complex space-time direction, which is neutral with respect to the orbifold. We denote them by χ_I . They are characterized by a $U(1)^4$ weight $\{0, 0, 0, 0\}$ and a charge $(+1)$ with respect to the last $U(1)$. The complex conjugate moduli $\bar{\chi}_I$, with weight $\{0, 0, 0, 0\}$ and charge (-1) , are paired in a Q -doublet with the fermionic moduli $\bar{\eta}_I$ coming from the ground state of the Ramond (R) sector with weight $\{-\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\}$ and charge $(-\frac{1}{2})$. All other moduli in this sector are arranged in Q -doublets. One doublet is $(A_I^{z_1}, M_I^{z_1})$, where $A_I^{z_1}$ is from the ψ^{z_1} oscillator in the NS sector with weight $\{+1, 0, 0, 0\}$ and charge 0, and $M_I^{z_1}$ is from the R ground state $\{+\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\}$ with charge $(+\frac{1}{2})$. Another doublet is $(A_I^{z_4}, M_I^{z_4})$, where $A_I^{z_4}$ is from the ψ^{z_4} oscillator in the NS sector with weight $\{0, 0, 0, +1\}$ and charge 0, and $M_I^{z_4}$ is from the R ground state with weight $\{-\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, +\frac{1}{2}\}$ and charge $(+\frac{1}{2})$. Also the

Doublet	$(-)^{F_\alpha}$	Chan-Paton	$U(1)^4$ charge	Q^2 -eigenvalue λ_α	Character
$(\bar{\chi}_I, \bar{\eta}_I)$	+	$(\mathbf{d}_I, \bar{\mathbf{d}}_I)$	$\{0, 0, 0, 0\}$	$\chi_{I,\sigma} - \chi_{I,\tau}$	
$(A_I^{z_1}, M_I^{z_1})$	+	$(\mathbf{d}_I, \bar{\mathbf{d}}_I)$	$\{+1, 0, 0, 0\}$	$\chi_{I,\sigma} - \chi_{I,\tau} + \epsilon_1$	$V_I^* V_I T_1$
$(A_I^{z_4}, M_I^{z_4})$	+	$(\mathbf{d}_I, \bar{\mathbf{d}}_I)$	$\{0, 0, 0, +1\}$	$\chi_{I,\sigma} - \chi_{I,\tau} + \epsilon_4$	$V_I^* V_I T_4$
(λ_I, D_I)	-	$(\mathbf{d}_I, \bar{\mathbf{d}}_I)$	$\{+\frac{1}{2}, +\frac{1}{2}, +\frac{1}{2}, +\frac{1}{2}\}$	$\chi_{I,\sigma} - \chi_{I,\tau}$	
$(\lambda_I^{z_1}, D_I^{z_1})$	-	$(\mathbf{d}_I, \bar{\mathbf{d}}_I)$	$\{+\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, +\frac{1}{2}\}$	$\chi_{I,\sigma} - \chi_{I,\tau} + \epsilon_1 + \epsilon_4$	$-V_I^* V_I T_1 T_4$
$(A_I^{z_2}, M_I^{z_2})$	+	$(\mathbf{d}_I, \bar{\mathbf{d}}_{I+1})$	$\{0, +1, 0, 0\}$	$\chi_{I,\sigma} - \chi_{I+1,\rho} + \hat{\epsilon}_2$	$V_{I+1}^* V_I T_2$
$(\lambda_I^{z_2}, D_I^{z_2})$	-	$(\mathbf{d}_I, \bar{\mathbf{d}}_{I+1})$	$\{-\frac{1}{2}, +\frac{1}{2}, -\frac{1}{2}, +\frac{1}{2}\}$	$\chi_{I,\sigma} - \chi_{I+1,\rho} + \hat{\epsilon}_2 + \epsilon_4$	$-V_{I+1}^* V_I T_2 T_4$
$(\bar{A}_I^{z_3}, \bar{M}_I^{z_3})$	+	$(\mathbf{d}_I, \bar{\mathbf{d}}_{I+1})$	$\{0, 0, -1, 0\}$	$\chi_{I,\sigma} - \chi_{I+1,\rho} - \hat{\epsilon}_3$	$V_{I+1}^* V_I T_1 T_2 T_4$
$(\lambda_I^{z_3}, D_I^{z_3})$	-	$(\mathbf{d}_I, \bar{\mathbf{d}}_{I+1})$	$\{+\frac{1}{2}, +\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\}$	$\chi_{I,\sigma} - \chi_{I+1,\rho} + \epsilon_1 + \hat{\epsilon}_2$	$-V_{I+1}^* V_I T_1 T_2$
(w_I, μ_I)	+	$(\mathbf{n}_I, \bar{\mathbf{d}}_I)$	$\{+\frac{1}{2}, +\frac{1}{2}, 0, 0\}$	$a_{I,s} - \chi_{I,\sigma} + \frac{1}{2}(\epsilon_1 + \hat{\epsilon}_2)$	$V_I^* W_I$
(μ'_I, h'_I)	-	$(\mathbf{n}_I, \bar{\mathbf{d}}_I)$	$\{0, 0, -\frac{1}{2}, +\frac{1}{2}\}$	$a_{I,s} - \chi_{I,\sigma} + \frac{1}{2}(\epsilon_1 + \hat{\epsilon}_2) + \epsilon_4$	$-V_I^* W_I T_4$
$(\hat{w}_I, \hat{\mu}_I)$	+	$(\mathbf{d}_I, \bar{\mathbf{n}}_{I+1})$	$\{+\frac{1}{2}, +\frac{1}{2}, 0, 0\}$	$\chi_{I,\sigma} - a_{I+1,t} + \frac{1}{2}(\epsilon_1 + \hat{\epsilon}_2)$	$W_{I+1}^* V_I T_1 T_2$
$(\hat{\mu}'_I, \hat{h}'_I)$	-	$(\mathbf{d}_I, \bar{\mathbf{n}}_{I+1})$	$\{0, 0, -\frac{1}{2}, +\frac{1}{2}\}$	$\chi_{I,\sigma} - a_{I+1,t} + \frac{1}{2}(\epsilon_1 + \hat{\epsilon}_2) + \epsilon_4$	$-W_{I+1}^* V_I T_1 T_2 T_4$

Table 1. The spectrum of moduli, organized in doublets of the BRST charge Q (or its conjugate \bar{Q}). For each of them, we display their statistics $(-)^{F_\alpha}$, the representation of the color and ADHM groups in which they transform, their charge vector with respect to the $U(1)^4$ symmetry, the eigenvalue λ_α of Q^2 and the corresponding contribution to the character. The neutral moduli carrying a superscript z_1, z_2, z_3 or z_4 , and the colored moduli in this table are complex. The quantities appearing in the last column, namely V_I, W_I, T_1, T_2 and T_4 are defined in (3.26) and (3.27).

complex conjugate doublets are present. Finally, there is a (real) doublet (λ_I, D_I) where λ_I is from the R ground state with weight $\{+\frac{1}{2}, +\frac{1}{2}, +\frac{1}{2}, +\frac{1}{2}\}$ and charge $(-\frac{1}{2})$, and D_I is an auxiliary field, and a complex doublet $(\lambda_I^{z_1}, D_I^{z_1})$ with $\lambda_I^{z_1}$ associated to the R ground state with weight $\{+\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}, +\frac{1}{2}\}$ and charge $(-\frac{1}{2})$, and $D_I^{z_1}$ an auxiliary field.

• **$(-1)/(-1)$ strings of type $I-(I+1)$:** in this sector the moduli transform in the bi-fundamental representation $(\mathbf{d}_I, \bar{\mathbf{d}}_{I+1})$ of $U(\mathbf{d}_I) \times U(\mathbf{d}_{I+1})$. In order to cancel the phase ω^{-1} due to the different representations on the Chan-Paton indices at the two endpoints, the weights under spacetime rotations of the operators creating the states in this sector must be such that $l_2 - l_3 = 1$. In this way they can survive the \mathbb{Z}_M -orbifold projection. Applying this requirement, we find a doublet $(A_I^{z_2}, M_I^{z_2})$, $A_I^{z_2}$ is from the ψ^{z_2} oscillator in the NS sector with weight $\{0, +1, 0, 0\}$ and charge 0, and $M_I^{z_2}$ is from the R ground state $\{-\frac{1}{2}, +\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\}$ with charge $(+\frac{1}{2})$. Another doublet is $(\bar{A}_I^{z_3}, \bar{M}_I^{z_3})$ where $\bar{A}_I^{z_3}$ is from the $\bar{\psi}^{z_3}$ oscillator in the NS sector with weight $\{0, 0, -1, 0\}$ and charge 0, and $\bar{M}_I^{z_3}$ is from the R ground state $\{+\frac{1}{2}, +\frac{1}{2}, -\frac{1}{2}, +\frac{1}{2}\}$ with charge $(+\frac{1}{2})$.¹⁶ Furthermore, we find two other complex Q -doublets, $(\lambda_I^{z_2}, D_I^{z_2})$ and $(\lambda_I^{z_3}, D_I^{z_3})$ where $\lambda_I^{z_2}$ and $\lambda_I^{z_3}$ are associated to the R ground states with weights $\{-\frac{1}{2}, +\frac{1}{2}, -\frac{1}{2}, +\frac{1}{2}\}$ and $\{+\frac{1}{2}, +\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2}\}$ and charges $(-\frac{1}{2})$,

¹⁶Notice that this last doublet is actually the complex conjugate of a Q -doublet of type $(I+1)-I$, which is made of $(A_I^{z_3}, M_I^{z_3})$ with $A_I^{z_3}$ corresponding to the weight $\{0, 0, 1, 0\}$ and $M_I^{z_3}$ corresponding to the weight $\{-\frac{1}{2}, -\frac{1}{2}, +\frac{1}{2}, -\frac{1}{2}\}$.

while $D_I^{z_2}$ and $D_I^{z_3}$ are auxiliary fields. Also the complex conjugate doublets are present in the \mathbb{Z}_M -invariant spectrum, and arise from strings with the opposite orientation.

- **3/(-1) strings of type I-I:** these open strings have mixed Neumann-Dirichlet boundary conditions along the (z_1, z_2) -directions and thus the corresponding states are characterized by the action of a twist operator Δ [50]. We assign an orbifold charge $\omega^{-\frac{1}{2}}$ to this twist operator, so that the states which survive the \mathbb{Z}_M -projection are those with weights such that $l_2 - l_3 = 1/2$. The moduli in this sector belong to the bi-fundamental representation $(\mathbf{n}_I \times \bar{\mathbf{d}}_I)$ of the gauge and ADHM groups, and form two complex doublets. One is (w_I, μ_I) where the NS component w_I has weight $\{+\frac{1}{2}, +\frac{1}{2}, 0, 0\}$ and charge 0, and the R component μ_I has weight $\{0, 0, -\frac{1}{2}, -\frac{1}{2}\}$ and charge $(+\frac{1}{2})$. The other doublet is (μ'_I, h'_I) where μ'_I is associated to the R ground state with weight $\{0, 0, -\frac{1}{2}, +\frac{1}{2}\}$ and charge $(-\frac{1}{2})$, while h'_I is an auxiliary field. Also the complex conjugate doublets, associated to the $(-1)/3$ strings of type I-I, are present in the spectrum.

- **(-1)/3 strings of type I-(I+1):** these open strings have mixed Dirichlet-Neumann boundary conditions along the (z_1, z_2) -directions and transform in the bi-fundamental representation $(\mathbf{d}_I \times \bar{\mathbf{n}}_{I+1})$ of the gauge and ADHM groups. As compared to the previous case, the states in this sector are characterized by the action of an anti-twist operator $\bar{\Delta}$ which carries an orbifold parity $\omega^{+\frac{1}{2}}$. Thus the \mathbb{Z}_M -invariant configurations must have again weights with $l_2 - l_3 = 1/2$ in order to compensate for the ω^{-1} factor carried by the Chan-Paton indices. Taking this into account, we find two complex doublets: $(\hat{w}_I, \hat{\mu}_I)$ where the NS component \hat{w}_I has weight $\{+\frac{1}{2}, +\frac{1}{2}, 0, 0\}$ and charge 0, and the R component $\hat{\mu}_I$ has weight $\{0, 0, -\frac{1}{2}, -\frac{1}{2}\}$ and charge $(+\frac{1}{2})$, and $(\hat{\mu}'_I, \hat{h}'_I)$ where $\hat{\mu}'_I$ is associated to the R ground state with weight $\{0, 0, -\frac{1}{2}, +\frac{1}{2}\}$ and charge $(-\frac{1}{2})$, while \hat{h}'_I is an auxiliary field. Also the complex conjugate doublets, associated to the $3/(-1)$ strings of type $(I+1)$ -I, are present in the spectrum.

Notice that no states from the $3/(-1)$ strings of type $I-(I+1)$ or from the $(-1)/3$ strings of type $(I+1)$ -I survive the orbifold projection. Indeed, in the first case the phases $\omega^{-\frac{1}{2}}$ and ω^{-1} from the twist operator Δ and the Chan-Paton factors cannot be compensated by the NS or R weights; while in the second case the phases $\omega^{+\frac{1}{2}}$ and ω^{+1} from the anti-twist operator $\bar{\Delta}$ and the Chan-Paton factors cannot be canceled.

All the above results are summarized in table 1, which contains also other relevant information about the moduli. As an illustrative example, we now consider in detail the SU(2) theory.

C.1 SU(2)

In this case we have $M = 2$, and thus necessarily $n_1 = n_2 = 1$. Therefore, in the SU(2) theory we have only simple surface operators. Furthermore, since the index s takes only one value, we can simplify the notation and suppress this index in the following.

Each pair $Y = (Y_1, Y_2)$ of Young tableaux contributes to the instanton partition function with a weight $q_1^{d_1} q_2^{d_2}$ where d_1 and d_2 are given by (3.22), which in this case take the

simple form [8]

$$d_1 = \sum_j \left(Y_1^{2j+1} + Y_2^{2j+1} \right), \quad d_2 = \sum_j \left(Y_1^{2j+2} + Y_2^{2j+2} \right). \quad (\text{C.1})$$

with Y_I^k representing the length of the k th column of the tableau Y_I .

Let us begin by considering the case of pairs of Young tableaux with a single box. There are two such pairs that can contribute. One is $Y = (\square, \bullet)$ corresponding to $d_1 = 1$ and $d_2 = 0$. Using these values in (3.18), we find

$$z_{\{1,0\}} = \frac{(\epsilon_1 + \epsilon_4) \left(a_1 - \chi_{1,1} + \frac{1}{2}(\epsilon_1 + \hat{\epsilon}_2) + \epsilon_4 \right) \left(\chi_{1,1} - a_2 + \frac{1}{2}(\epsilon_1 + \hat{\epsilon}_2) + \epsilon_4 \right)}{\epsilon_1 \epsilon_4 \left(a_1 - \chi_{1,1} + \frac{1}{2}(\epsilon_1 + \hat{\epsilon}_2) \right) \left(\chi_{1,1} - a_2 + \frac{1}{2}(\epsilon_1 + \hat{\epsilon}_2) \right)} \quad (\text{C.2})$$

Due to the prescription (3.19), only the pole at

$$\chi_{1,1} = a_1 + \frac{1}{2}(\epsilon_1 + \hat{\epsilon}_2) \quad (\text{C.3})$$

contributes to the contour integral over $\chi_{1,1}$, yielding

$$Z_{(\square, \bullet)} = \frac{(\epsilon_1 + \epsilon_4) (a_{12} + \epsilon_1 + \hat{\epsilon}_2 + \epsilon_4)}{\epsilon_1 (a_{12} + \epsilon_1 + \hat{\epsilon}_2)} = \frac{(\epsilon_1 + \epsilon_4) (4a + 2\epsilon_1 + \epsilon_2 + 2\epsilon_4)}{\epsilon_1 (4a + 2\epsilon_1 + \epsilon_2)} \quad (\text{C.4})$$

where in the last step we used the notation $a_{12} = a_1 - a_2 = 2a$ and reintroduced $\epsilon_2 = 2\hat{\epsilon}_2$. A similar analysis can be done for the second pair of tableaux with one box that contributes, namely $Y = (\bullet, \square)$ corresponding to $d_1 = 0$ and $d_2 = 1$. In this case we find

$$Z_{(\bullet, \square)} = \frac{(\epsilon_1 + \epsilon_4) (-4a + 2\epsilon_1 + \epsilon_2 + 2\epsilon_4)}{\epsilon_1 (-4a + 2\epsilon_1 + \epsilon_2)}. \quad (\text{C.5})$$

In the case of two boxes, we have five different pairs of tableaux that can contribute. They are: $Y = (\square, \square)$, $Y = (\square\square, \bullet)$, $Y = (\bullet, \square\square)$, $Y = (\square, \bullet)$ and $Y = (\bullet, \square)$. The contributions of these five diagrams are listed below in table 2.

Multiplying all contributions with the appropriate weight factor and summing over them, we obtain the instanton partition function for the SU(2) gauge theory in the presence of the surface operator:

$$\begin{aligned} Z_{\text{inst}}[1, 1] = & 1 + q_1 \frac{(\epsilon_1 + \epsilon_4) (4a + 2\epsilon_1 + \epsilon_2 + 2\epsilon_4)}{\epsilon_1 (4a + 2\epsilon_1 + \epsilon_2)} + q_2 \frac{(\epsilon_1 + \epsilon_4) (-4a + 2\epsilon_1 + \epsilon_2 + 2\epsilon_4)}{\epsilon_1 (-4a + 2\epsilon_1 + \epsilon_2)} \\ & + q_1^2 \frac{(\epsilon_1 + \epsilon_4) (2\epsilon_1 + \epsilon_4) (4a + 2\epsilon_1 + \epsilon_2 + 2\epsilon_4) (4a + 4\epsilon_1 + \epsilon_2 + 2\epsilon_4)}{2\epsilon_1^2 (4a + 2\epsilon_1 + \epsilon_2) (4a + 4\epsilon_1 + \epsilon_2)} \\ & + q_2^2 \frac{(\epsilon_1 + \epsilon_4) (2\epsilon_1 + \epsilon_4) (-4a + 2\epsilon_1 + \epsilon_2 + 2\epsilon_4) (-4a + 4\epsilon_1 + \epsilon_2 + 2\epsilon_4)}{2\epsilon_1^2 (-4a + 2\epsilon_1 + \epsilon_2) (-4a + 4\epsilon_1 + \epsilon_2)} \\ & + q_1 q_2 \left(\frac{(\epsilon_1 + \epsilon_4) (\epsilon_2 + \epsilon_4) (4a + \epsilon_2 - 2\epsilon_4) (4a + 2\epsilon_1 + \epsilon_2 + 2\epsilon_4)}{\epsilon_1 \epsilon_2 (4a + \epsilon_2) (4a + 2\epsilon_1 + \epsilon_2)} \right. \\ & \quad + \frac{(\epsilon_1 + \epsilon_4) (\epsilon_2 + \epsilon_4) (-4a + \epsilon_2 - 2\epsilon_4) (-4a + 2\epsilon_1 + \epsilon_2 + 2\epsilon_4)}{\epsilon_1 \epsilon_2 (-4a + \epsilon_2) (-4a + 2\epsilon_1 + \epsilon_2)} \\ & \quad \left. + \frac{(\epsilon_1 + \epsilon_4)^2 (4a + \epsilon_2 + 2\epsilon_4) (-4a + \epsilon_2 + 2\epsilon_4)}{\epsilon_1^2 (4a + \epsilon_2) (-4a + \epsilon_2)} \right) + \dots \quad (\text{C.6}) \end{aligned}$$

Y	weight	poles	Z_Y
(\square, \square)	$q_1 q_2$	$\chi_{1,1} = a_1 + \frac{1}{2}(\epsilon_1 + \hat{\epsilon}_2)$ $\chi_{2,1} = a_2 + \frac{1}{2}(\epsilon_1 + \hat{\epsilon}_2)$	$\frac{(\epsilon_1 + \epsilon_4)^2 (4a + \epsilon_2 + 2\epsilon_4)(-4a + \epsilon_2 + 2\epsilon_4)}{\epsilon_1^2 (4a + \epsilon_2)(-4a + \epsilon_2)}$
$(\square\square, \bullet)$	$q_1 q_2$	$\chi_{1,1} = a_1 + \frac{1}{2}(\epsilon_1 + \hat{\epsilon}_2)$ $\chi_{2,1} = \chi_{1,1} + \hat{\epsilon}_2$	$\frac{(\epsilon_1 + \epsilon_4)(\epsilon_2 + \epsilon_4)(4a + \epsilon_2 - 2\epsilon_4)(4a + 2\epsilon_1 + \epsilon_2 + 2\epsilon_4)}{\epsilon_1 \epsilon_2 (4a + \epsilon_2)(4a + 2\epsilon_1 + \epsilon_2)}$
$(\bullet, \square\square)$	$q_1 q_2$	$\chi_{2,1} = a_2 + \frac{1}{2}(\epsilon_1 + \hat{\epsilon}_2)$ $\chi_{1,1} = \chi_{2,1} + \hat{\epsilon}_2$	$\frac{(\epsilon_1 + \epsilon_4)(\epsilon_2 + \epsilon_4)(-4a + \epsilon_2 - 2\epsilon_4)(-4a + 2\epsilon_1 + \epsilon_2 + 2\epsilon_4)}{\epsilon_1 \epsilon_2 (-4a + \epsilon_2)(-4a + 2\epsilon_1 + \epsilon_2)}$
$(\begin{smallmatrix} \square \\ \square \end{smallmatrix}, \bullet)$	q_1^2	$\chi_{1,1} = a_1 + \frac{1}{2}(\epsilon_1 + \hat{\epsilon}_2)$ $\chi_{1,2} = \chi_{1,1} + \epsilon_1$	$\frac{(\epsilon_1 + \epsilon_4)(2\epsilon_1 + \epsilon_4)(4a + 2\epsilon_1 + \epsilon_2 + 2\epsilon_4)(4a + 4\epsilon_1 + \epsilon_2 + 2\epsilon_4)}{2\epsilon_1^2 (4a + 2\epsilon_1 + \epsilon_2)(4a + 4\epsilon_1 + \epsilon_2)}$
$(\bullet, \begin{smallmatrix} \square \\ \square \end{smallmatrix})$	q_2^2	$\chi_{2,1} = a_2 + \frac{1}{2}(\epsilon_1 + \hat{\epsilon}_2)$ $\chi_{2,2} = \chi_{2,1} + \epsilon_1$	$\frac{(\epsilon_1 + \epsilon_4)(2\epsilon_1 + \epsilon_4)(-4a + 2\epsilon_1 + \epsilon_2 + 2\epsilon_4)(-4a + 4\epsilon_1 + \epsilon_2 + 2\epsilon_4)}{2\epsilon_1^2 (-4a + 2\epsilon_1 + \epsilon_2)(-4a + 4\epsilon_1 + \epsilon_2)}$

Table 2. We list the tableaux, the weight factors, the pole structure and the contribution to the partition function in all five cases with two boxes for the SU(2) theory.

where the ellipses stand for the contributions originating from tableaux with higher number of boxes, which can be easily generated with a computer program. We have explicitly computed these terms up six boxes, but we do not write them here since the raw expressions are very long and not particularly illuminating. To the extent it is possible to make comparisons, we observe that the above result agrees with the instanton partition function reported in eq. (B.6) of [8] under the following change of notation

$$q_1 \rightarrow y, \quad q_2 \rightarrow x, \quad \epsilon_4 \rightarrow -m, \quad 2a \rightarrow 2a + \frac{\epsilon_2}{2}. \quad (\text{C.7})$$

Note then that the mass m appearing in [8] is the equivariant mass of the hypermultiplet [78], which differs by ϵ -corrections from the mass we have used in this paper (see (3.35)).

D Prepotential coefficients for the SU(N) gauge theory

The prepotential \mathcal{F} of the $\mathcal{N} = 2^*$ SU(N) gauge theory has been determined in terms of quasi-modular forms in [34, 41]. Expanding \mathcal{F} as in (4.12), the first few non-zero coefficients f_ℓ in the NS limit turn out to be

$$f_2 = \frac{1}{4} \left(m^2 - \frac{\epsilon_1^2}{4} \right) \sum_{u \neq v} \log \frac{(a_u - a_v)^2}{\Lambda^2} + N \left(m^2 - \frac{\epsilon_1^2}{4} \right) \log \hat{\eta}, \quad (\text{D.1})$$

$$f_4 = -\frac{1}{24} \left(m^2 - \frac{\epsilon_1^2}{4} \right)^2 E_2 C_2, \quad (\text{D.2})$$

$$f_6 = -\frac{1}{288} \left(m^2 - \frac{\epsilon_1^2}{4} \right)^2 \left\{ \left[\frac{2}{5} \left(m^2 - \frac{\epsilon_1^2}{4} \right) (5E_2^2 + E_4) - 6\epsilon_1^2 E_4 \right] C_4 + \frac{1}{2} \left(m^2 - \frac{\epsilon_1^2}{4} \right) (E_2^2 - E_4) C_{2;1,1} \right\}, \quad (\text{D.3})$$

$$\begin{aligned}
 f_8 = & -\frac{1}{1728} \left(m^2 - \frac{\epsilon_1^2}{4}\right)^2 \left\{ \left[\frac{2}{105} \left(m^2 - \frac{\epsilon_1^2}{4}\right)^2 (175E_2^3 + 84E_2E_4 + 11E_6) \right. \right. \\
 & \left. \left. - \frac{24\epsilon^2}{35} \left(m^2 - \frac{\epsilon_1^2}{4}\right) (7E_2E_4 + 3E_6) + \frac{24\epsilon^4}{7} E_6 \right] C_6 \right. \\
 & - \frac{1}{5} \left(m^2 - \frac{\epsilon_1^2}{4}\right) \left[\left(m^2 - \frac{\epsilon_1^2}{4}\right) (5E_2^3 - 3E_2E_4 - 2E_6) - 6\epsilon^2 (E_2E_4 - E_6) \right] C_{4;2} \\
 & - \frac{1}{5} \left(m^2 - \frac{\epsilon_1^2}{4}\right) \left[\frac{1}{12} \left(m^2 - \frac{\epsilon_1^2}{4}\right) (5E_2^3 - 3E_2E_4 - 2E_6) - 3\epsilon^2 (E_2E_4 - E_6) \right] C_{3;3} \\
 & \left. + \frac{1}{24} \left(m^2 - \frac{\epsilon_1^2}{4}\right)^2 (E_2^3 - 3E_2E_4 + 2E_6) C_{2;1,1,1,1} \right\}. \tag{D.4}
 \end{aligned}$$

Here E_2 , E_4 and E_6 are the Eisenstein series and

$$\log \hat{\eta} = - \sum_{k=1}^{\infty} \frac{\sigma_1(k)}{k} q^k = -\frac{1}{24} \log q + \log \eta \tag{D.5}$$

with η being the Dedekind η -function. Finally, the root lattice sums are defined by

$$C_{n;m_1,m_2,\dots,m_k} = \sum_{\vec{\alpha} \in \Phi} \sum_{\vec{\beta}_1 \neq \vec{\beta}_2 \neq \dots \neq \vec{\beta}_k \in \Phi(\vec{\alpha})} \frac{1}{(\vec{\alpha} \cdot \vec{a})^n (\vec{\beta}_1 \cdot \vec{a})^{m_1} (\vec{\beta}_2 \cdot \vec{a})^{m_2} \dots (\vec{\beta}_k \cdot \vec{a})^{m_k}} \tag{D.6}$$

where Φ is the root system of $SU(N)$ and

$$\Phi(\vec{\alpha}) = \{ \vec{\beta} \in \Phi \mid \vec{\alpha} \cdot \vec{\beta} = 1 \}. \tag{D.7}$$

We refer to [41] for the details and the derivation of these results. Notice, however, that we have slightly changed our notation, since $f_{2\ell}^{\text{here}} = f_{\ell}^{\text{there}}$. By expanding the modular functions in powers of q and selecting $SU(2)$ as gauge group, it is easy to show that the above formulas reproduce both the perturbative part and the instanton contributions, reported respectively in (4.10a) and (3.42) of the main text.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] S. Gukov, *Surface operators*, [arXiv:1412.7127](https://arxiv.org/abs/1412.7127) [[INSPIRE](https://inspirehep.net/literature/1412712)].
- [2] S. Gukov and E. Witten, *Gauge theory, ramification, and the geometric Langlands program*, [hep-th/0612073](https://arxiv.org/abs/hep-th/0612073) [[INSPIRE](https://inspirehep.net/literature/561625)].
- [3] S. Gukov and E. Witten, *Rigid surface operators*, *Adv. Theor. Math. Phys.* **14** (2010) 87 [[arXiv:0804.1561](https://arxiv.org/abs/0804.1561)] [[INSPIRE](https://inspirehep.net/literature/770432)].
- [4] D. Gaiotto, *Surface operators in $N = 2$ 4d gauge theories*, *JHEP* **11** (2012) 090 [[arXiv:0911.1316](https://arxiv.org/abs/0911.1316)] [[INSPIRE](https://inspirehep.net/literature/820803)].

- [5] D. Gaiotto, S. Gukov and N. Seiberg, *Surface defects and resolvents*, *JHEP* **09** (2013) 070 [[arXiv:1307.2578](#)] [[INSPIRE](#)].
- [6] E. Witten, *Solutions of four-dimensional field theories via M-theory*, *Nucl. Phys. B* **500** (1997) 3 [[hep-th/9703166](#)] [[INSPIRE](#)].
- [7] D. Gaiotto, *$N = 2$ dualities*, *JHEP* **08** (2012) 034 [[arXiv:0904.2715](#)] [[INSPIRE](#)].
- [8] L.F. Alday and Y. Tachikawa, *Affine $SL(2)$ conformal blocks from 4d gauge theories*, *Lett. Math. Phys.* **94** (2010) 87 [[arXiv:1005.4469](#)] [[INSPIRE](#)].
- [9] L.F. Alday, D. Gaiotto and Y. Tachikawa, *Liouville correlation functions from four-dimensional gauge theories*, *Lett. Math. Phys.* **91** (2010) 167 [[arXiv:0906.3219](#)] [[INSPIRE](#)].
- [10] C. Kozcaz, S. Pasquetti, F. Passerini and N. Wyllard, *Affine $sl(N)$ conformal blocks from $N = 2$ $SU(N)$ gauge theories*, *JHEP* **01** (2011) 045 [[arXiv:1008.1412](#)] [[INSPIRE](#)].
- [11] L.F. Alday, D. Gaiotto, S. Gukov, Y. Tachikawa and H. Verlinde, *Loop and surface operators in $N = 2$ gauge theory and Liouville modular geometry*, *JHEP* **01** (2010) 113 [[arXiv:0909.0945](#)] [[INSPIRE](#)].
- [12] M. Taki, *On AGT conjecture for pure super Yang-Mills and W -algebra*, *JHEP* **05** (2011) 038 [[arXiv:0912.4789](#)] [[INSPIRE](#)].
- [13] A. Marshakov, A. Mironov and A. Morozov, *On AGT relations with surface operator insertion and stationary limit of beta-ensembles*, *J. Geom. Phys.* **61** (2011) 1203 [[arXiv:1011.4491](#)] [[INSPIRE](#)].
- [14] C. Kozcaz, S. Pasquetti and N. Wyllard, *A & B model approaches to surface operators and Toda theories*, *JHEP* **08** (2010) 042 [[arXiv:1004.2025](#)] [[INSPIRE](#)].
- [15] T. Dimofte, S. Gukov and L. Hollands, *Vortex counting and Lagrangian 3-manifolds*, *Lett. Math. Phys.* **98** (2011) 225 [[arXiv:1006.0977](#)] [[INSPIRE](#)].
- [16] K. Maruyoshi and M. Taki, *Deformed prepotential, quantum integrable system and Liouville field theory*, *Nucl. Phys. B* **841** (2010) 388 [[arXiv:1006.4505](#)] [[INSPIRE](#)].
- [17] M. Taki, *Surface operator, bubbling Calabi-Yau and AGT relation*, *JHEP* **07** (2011) 047 [[arXiv:1007.2524](#)] [[INSPIRE](#)].
- [18] M. Bullimore, H.-C. Kim and P. Koroteev, *Defects and quantum Seiberg-Witten geometry*, *JHEP* **05** (2015) 095 [[arXiv:1412.6081](#)] [[INSPIRE](#)].
- [19] B. Assel and S. Schäfer-Nameki, *Six-dimensional origin of $N = 4$ SYM with duality defects*, *JHEP* **12** (2016) 058 [[arXiv:1610.03663](#)] [[INSPIRE](#)].
- [20] N.A. Nekrasov, *Seiberg-Witten prepotential from instanton counting*, *Adv. Theor. Math. Phys.* **7** (2003) 831 [[hep-th/0206161](#)] [[INSPIRE](#)].
- [21] M. Billò, M. Frau, L. Gallot, A. Lerda and I. Pesando, *Deformed $N = 2$ theories, generalized recursion relations and S -duality*, *JHEP* **04** (2013) 039 [[arXiv:1302.0686](#)] [[INSPIRE](#)].
- [22] M. Billò, M. Frau, L. Gallot, A. Lerda and I. Pesando, *Modular anomaly equation, heat kernel and S -duality in $N = 2$ theories*, *JHEP* **11** (2013) 123 [[arXiv:1307.6648](#)] [[INSPIRE](#)].
- [23] J.A. Minahan, D. Nemeschansky and N.P. Warner, *Instanton expansions for mass deformed $N = 4$ super Yang-Mills theories*, *Nucl. Phys. B* **528** (1998) 109 [[hep-th/9710146](#)] [[INSPIRE](#)].

- [24] M. Bershadsky, S. Cecotti, H. Ooguri and C. Vafa, *Holomorphic anomalies in topological field theories*, *Nucl. Phys. B* **405** (1993) 279 [[hep-th/9302103](#)] [[INSPIRE](#)].
- [25] E. Witten, *Quantum background independence in string theory*, [hep-th/9306122](#) [[INSPIRE](#)].
- [26] M. Aganagic, V. Bouchard and A. Klemm, *Topological strings and (almost) modular forms*, *Commun. Math. Phys.* **277** (2008) 771 [[hep-th/0607100](#)] [[INSPIRE](#)].
- [27] M. Günaydin, A. Neitzke and B. Pioline, *Topological wave functions and heat equations*, *JHEP* **12** (2006) 070 [[hep-th/0607200](#)] [[INSPIRE](#)].
- [28] M.-X. Huang and A. Klemm, *Holomorphic anomaly in gauge theories and matrix models*, *JHEP* **09** (2007) 054 [[hep-th/0605195](#)] [[INSPIRE](#)].
- [29] T.W. Grimm, A. Klemm, M. Mariño and M. Weiss, *Direct integration of the topological string*, *JHEP* **08** (2007) 058 [[hep-th/0702187](#)] [[INSPIRE](#)].
- [30] M.-X. Huang and A. Klemm, *Holomorphicity and modularity in Seiberg-Witten theories with matter*, *JHEP* **07** (2010) 083 [[arXiv:0902.1325](#)] [[INSPIRE](#)].
- [31] M.-X. Huang and A. Klemm, *Direct integration for general Ω backgrounds*, *Adv. Theor. Math. Phys.* **16** (2012) 805 [[arXiv:1009.1126](#)] [[INSPIRE](#)].
- [32] D. Galakhov, A. Mironov and A. Morozov, *S-duality as a beta-deformed Fourier transform*, *JHEP* **08** (2012) 067 [[arXiv:1205.4998](#)] [[INSPIRE](#)].
- [33] N. Nemkov, *S-duality as Fourier transform for arbitrary ϵ_1, ϵ_2* , *J. Phys.* **47** (2014) 105401 [[arXiv:1307.0773](#)] [[INSPIRE](#)].
- [34] M. Billò et al., *Modular anomaly equations in $N = 2^*$ theories and their large- N limit*, *JHEP* **10** (2014) 131 [[arXiv:1406.7255](#)] [[INSPIRE](#)].
- [35] A.-K. Kashani-Poor and J. Troost, *The toroidal block and the genus expansion*, *JHEP* **03** (2013) 133 [[arXiv:1212.0722](#)] [[INSPIRE](#)].
- [36] A.-K. Kashani-Poor and J. Troost, *Transformations of spherical blocks*, *JHEP* **10** (2013) 009 [[arXiv:1305.7408](#)] [[INSPIRE](#)].
- [37] A.-K. Kashani-Poor and J. Troost, *Quantum geometry from the toroidal block*, *JHEP* **08** (2014) 117 [[arXiv:1404.7378](#)] [[INSPIRE](#)].
- [38] S.K. Ashok, M. Billò, E. Dell’Aquila, M. Frau, A. Lerda and M. Raman, *Modular anomaly equations and S-duality in $N = 2$ conformal SQCD*, *JHEP* **10** (2015) 091 [[arXiv:1507.07476](#)] [[INSPIRE](#)].
- [39] S.K. Ashok, E. Dell’Aquila, A. Lerda and M. Raman, *S-duality, triangle groups and modular anomalies in $N = 2$ SQCD*, *JHEP* **04** (2016) 118 [[arXiv:1601.01827](#)] [[INSPIRE](#)].
- [40] M. Beccaria, A. Fachechi, G. Macorini and L. Martina, *Exact partition functions for deformed $N = 2$ theories with $N_f = 4$ flavours*, *JHEP* **12** (2016) 029 [[arXiv:1609.01189](#)] [[INSPIRE](#)].
- [41] M. Billò, M. Frau, F. Fucito, A. Lerda and J.F. Morales, *S-duality and the prepotential in $N = 2^*$ theories (I): the ADE algebras*, *JHEP* **11** (2015) 024 [[arXiv:1507.07709](#)] [[INSPIRE](#)].
- [42] M. Billò, M. Frau, F. Fucito, A. Lerda and J.F. Morales, *S-duality and the prepotential of $N = 2^*$ theories (II): the non-simply laced algebras*, *JHEP* **11** (2015) 026 [[arXiv:1507.08027](#)] [[INSPIRE](#)].

- [43] M. Beccaria and G. Macorini, *Exact partition functions for the Ω -deformed $N = 2^*$ $SU(2)$ gauge theory*, *JHEP* **07** (2016) 066 [[arXiv:1606.00179](#)] [[INSPIRE](#)].
- [44] S.K. Ashok et al., *Chiral observables and S -duality in $N = 2^*$ $U(N)$ gauge theories*, *JHEP* **11** (2016) 020 [[arXiv:1607.08327](#)] [[INSPIRE](#)].
- [45] H. Kanno and Y. Tachikawa, *Instanton counting with a surface operator and the chain-saw quiver*, *JHEP* **06** (2011) 119 [[arXiv:1105.0357](#)] [[INSPIRE](#)].
- [46] V. Mehta and C. Seshadri, *Moduli of vector bundles on curves with parabolic structures*, *Math. Ann.* **248** (1980) 205.
- [47] I. Biswas, *Parabolic bundles as orbifold bundles*, *Duke Math. J.* **88** (1997) 305.
- [48] B. Feigin, M. Finkelberg, A. Negut and L. Rybnikov, *Yangians and cohomology rings of Laumon spaces*, *Selecta Math.* **17** (2011) 573 [[arXiv:0812.4656](#)].
- [49] M.R. Douglas, *Branes within branes*, [hep-th/9512077](#) [[INSPIRE](#)].
- [50] M. Billò, M. Frau, I. Pesando, F. Fucito, A. Lerda and A. Liccardo, *Classical gauge instantons from open strings*, *JHEP* **02** (2003) 045 [[hep-th/0211250](#)] [[INSPIRE](#)].
- [51] N.A. Nekrasov and S.L. Shatashvili, *Quantization of integrable systems and four dimensional gauge theories*, [arXiv:0908.4052](#) [[INSPIRE](#)].
- [52] E. Frenkel, S. Gukov and J. Teschner, *Surface operators and separation of variables*, *JHEP* **01** (2016) 179 [[arXiv:1506.07508](#)] [[INSPIRE](#)].
- [53] G.W. Moore, N. Nekrasov and S. Shatashvili, *D particle bound states and generalized instantons*, *Commun. Math. Phys.* **209** (2000) 77 [[hep-th/9803265](#)] [[INSPIRE](#)].
- [54] G. Bonelli, A. Sciarappa, A. Tanzini and P. Vasko, *Vortex partition functions, wall crossing and equivariant Gromov-Witten invariants*, *Commun. Math. Phys.* **333** (2015) 717 [[arXiv:1307.5997](#)] [[INSPIRE](#)].
- [55] S. Nawata, *Givental J -functions, quantum integrable systems, AGT relation with surface operator*, *Adv. Theor. Math. Phys.* **19** (2015) 1277 [[arXiv:1408.4132](#)] [[INSPIRE](#)].
- [56] N. Doroud, J. Gomis, B. Le Floch and S. Lee, *Exact results in $D = 2$ supersymmetric gauge theories*, *JHEP* **05** (2013) 093 [[arXiv:1206.2606](#)] [[INSPIRE](#)].
- [57] F. Benini and S. Cremonesi, *Partition functions of $N = (2, 2)$ gauge theories on S^2 and vortices*, *Commun. Math. Phys.* **334** (2015) 1483 [[arXiv:1206.2356](#)] [[INSPIRE](#)].
- [58] M. Billò, M. Frau, F. Fucito and A. Lerda, *Instanton calculus in RR background and the topological string*, *JHEP* **11** (2006) 012 [[hep-th/0606013](#)] [[INSPIRE](#)].
- [59] N. Wyllard, *Instanton partition functions in $N = 2$ $SU(N)$ gauge theories with a general surface operator and their W -algebra duals*, *JHEP* **02** (2011) 114 [[arXiv:1012.1355](#)] [[INSPIRE](#)].
- [60] H. Awata, H. Fuji, H. Kanno, M. Manabe and Y. Yamada, *Localization with a surface operator, irregular conformal blocks and open topological string*, *Adv. Theor. Math. Phys.* **16** (2012) 725 [[arXiv:1008.0574](#)] [[INSPIRE](#)].
- [61] N. Wyllard, *W -algebras and surface operators in $N = 2$ gauge theories*, *J. Phys. A* **44** (2011) 155401 [[arXiv:1011.0289](#)] [[INSPIRE](#)].
- [62] E. Witten, *Phases of $N = 2$ theories in two-dimensions*, *Nucl. Phys. B* **403** (1993) 159 [[hep-th/9301042](#)] [[INSPIRE](#)].

- [63] A. Hanany and K. Hori, *Branes and $N = 2$ theories in two-dimensions*, *Nucl. Phys. B* **513** (1998) 119 [[hep-th/9707192](#)] [[INSPIRE](#)].
- [64] E. D'Hoker, I.M. Krichever and D.H. Phong, *The effective prepotential of $N = 2$ supersymmetric $SU(N_c)$ gauge theories*, *Nucl. Phys. B* **489** (1997) 179 [[hep-th/9609041](#)] [[INSPIRE](#)].
- [65] S.G. Naculich, H.J. Schnitzer and N. Wyllard, *The $N = 2$ $U(N)$ gauge theory prepotential and periods from a perturbative matrix model calculation*, *Nucl. Phys. B* **651** (2003) 106 [[hep-th/0211123](#)] [[INSPIRE](#)].
- [66] N. Nekrasov and V. Pestun, *Seiberg-Witten geometry of four dimensional $N = 2$ quiver gauge theories*, [arXiv:1211.2240](#) [[INSPIRE](#)].
- [67] F. Fucito, J.F. Morales, D.R. Pacifici and R. Poghossian, *Gauge theories on Ω -backgrounds from non commutative Seiberg-Witten curves*, *JHEP* **05** (2011) 098 [[arXiv:1103.4495](#)] [[INSPIRE](#)].
- [68] F. Fucito, J.F. Morales and D. Ricci Pacifici, *Deformed Seiberg-Witten curves for ADE quivers*, *JHEP* **01** (2013) 091 [[arXiv:1210.3580](#)] [[INSPIRE](#)].
- [69] M. Beccaria, A. Fachechi and G. Macorini, *Chiral trace relations in Ω -deformed $N = 2$ theories*, *JHEP* **05** (2017) 023 [[arXiv:1702.01254](#)] [[INSPIRE](#)].
- [70] R. Donagi and E. Witten, *Supersymmetric Yang-Mills theory and integrable systems*, *Nucl. Phys. B* **460** (1996) 299 [[hep-th/9510101](#)] [[INSPIRE](#)].
- [71] E. D'Hoker and D.H. Phong, *Calogero-Moser systems in $SU(N)$ Seiberg-Witten theory*, *Nucl. Phys. B* **513** (1998) 405 [[hep-th/9709053](#)] [[INSPIRE](#)].
- [72] D. Gaiotto, G.W. Moore and A. Neitzke, *Wall-crossing in coupled 2d-4d systems*, *JHEP* **12** (2012) 082 [[arXiv:1103.2598](#)] [[INSPIRE](#)].
- [73] G. Bonelli, A. Tanzini and J. Zhao, *Vertices, vortices and interacting surface operators*, *JHEP* **06** (2012) 178 [[arXiv:1102.0184](#)] [[INSPIRE](#)].
- [74] G. Bonelli, A. Tanzini and J. Zhao, *The Liouville side of the vortex*, *JHEP* **09** (2011) 096 [[arXiv:1107.2787](#)] [[INSPIRE](#)].
- [75] J. Gomis and B. Le Floch, *$M2$ -brane surface operators and gauge theory dualities in Toda*, *JHEP* **04** (2016) 183 [[arXiv:1407.1852](#)] [[INSPIRE](#)].
- [76] J. Gomis, B. Le Floch, Y. Pan and W. Peelaers, *Intersecting surface defects and two-dimensional CFT*, [arXiv:1610.03501](#) [[INSPIRE](#)].
- [77] Y. Pan and W. Peelaers, *Intersecting surface defects and instanton partition functions*, [arXiv:1612.04839](#) [[INSPIRE](#)].
- [78] T. Okuda and V. Pestun, *On the instantons and the hypermultiplet mass of $N = 2^*$ super Yang-Mills on S^4* , *JHEP* **03** (2012) 017 [[arXiv:1004.1222](#)] [[INSPIRE](#)].