

Dependence of the layer heat potentials upon support perturbations

Matteo Dalla Riva ^{*}, Paolo Luzzini [†]

March 31, 2023

Abstract

We prove that the integral operators associated with the layer heat potentials depend smoothly upon a parametrization of the support of integration. The analysis is carried out in the optimal Hölder setting.

Keywords: Heat equation; layer heat potentials; transmission problem; shape sensitivity analysis; perturbed domain.

2020 Mathematics Subject Classification: 31B10; 47G10; 35K05; 35K20.

1 Introduction

Potential theory is a powerful tool for analyzing boundary value problems for partial differential equations, particularly in the case of elliptic and parabolic operators. Along with existence, uniqueness, and regularity issues, potential-theoretic methods can be used to study perturbation problems. For example, potential-theoretic techniques can be employed to analyze how a solution depends on a deformation of the domain. To apply this kind of approach, it is important to understand how layer potentials and other potential-type integral operators depend on variations of the support of integration.

In the literature, most of the research in this direction focuses on the elliptic case. We mention, for example, Coifman and Meyer's results [7] and Wu's results [48] on the analytic dependence of the Cauchy integral on a variable arc-length parametrized curve. Moreover, Potthast's work [40, 41, 42] and Potthast and Stratis' results [43] establish a Fréchet differentiability result for layer potentials associated with the Helmholtz operator and apply these findings to some inverse problems in acoustic and electromagnetic scattering. Lanza de Cristoforis and Preciso [28] showed that the Cauchy integral depends analytically on a parametrization of the support. Later, Lanza de Cristoforis and Rossi [29, 30] considered the case of layer potentials associated with the Laplace and Helmholtz operators and proved real analyticity results which were used in [23, 24] to study domain perturbation problems for the Laplace and Poisson equations. In [9], we have employed similar techniques for a perturbed obstacle scattering problem. The case of layer potentials associated with a general second-order elliptic operator with constant coefficients has been analyzed in [8], while the periodic and quasi-periodic cases are considered in

^{*}Dipartimento di Ingegneria, Università degli Studi di Palermo, Viale delle Scienze, Ed. 8, 90128 Palermo, Italy.

[†]Dipartimento di Matematica 'Tullio Levi Civita', Università degli Studi di Padova, Via Trieste 63, 35121 Padova, Italy.

Lanza de Cristoforis and Musolino [27] and [3]. In a collaboration with Musolino and Pukhtaievych we have exploited the results on the periodic potentials for a shape sensitivity analysis of the longitudinal flow along a periodic array of cylinders and the effective conductivity of periodic composites (see, e.g., [37, 36, 11]). More recently, Henríquez and Schwab [19] proved that the Calderón projector of the Laplacian in \mathbb{R}^2 is an holomorphic function of the shape of the support, a result that has been extended to \mathbb{R}^n in [10].

Moreover, several authors have explored elliptic domain perturbation problems using approaches other than potential theory. Examples include Bucur and Buttazzo [4], Daners [13], Delfour and Zolésio [15], Henrot and Pierre [17], Novotny and Sokołowski [38], Pironneau [39], and Sokołowski and Zolésio [45].

So, we can conclude that the literature on shape stability and regularity is fairly rich in the case of elliptic operators. The parabolic case, instead, seems to be far less understood. Daners [12] studied the stability of solutions of parabolic problems upon domain perturbation, while Chapko, Kress, and Yoon [5, 6] proved shape differentiability results for the solutions of the Dirichlet and Neumann problems for the heat equation. Then, they applied these results to some inverse problems in heat conduction. However, to the best of our knowledge, higher regularity results are still unavailable. Moreover, an analysis of the shape dependence of the integral operators arising in parabolic potential theory is lacking in the literature. To fill this gap, this paper aims to develop a high regularity theory by analyzing the dependence of layer heat potentials on variations in the shape of the support of integration. To wit, we fix a regular open subset Ω of \mathbb{R}^n , which plays the role of a reference set, and we introduce a class of diffeomorphisms $\mathcal{A}_{\partial\Omega}$ from $\partial\Omega$ to \mathbb{R}^n (see Definition 2.1 and Figure 1). Our main results prove that the maps that take a function $\phi \in \mathcal{A}_{\partial\Omega}$ to certain layer heat potentials supported on $\phi(\partial\Omega)$ are of class C^∞ .

We now describe these results with a few more details. Let $S_n : \mathbb{R}^{1+n} \setminus \{0, 0\} \rightarrow \mathbb{R}$ denote the usual fundamental solution of the heat operator, that is

$$S_n(t, x) := \begin{cases} \frac{1}{(4\pi t)^{\frac{n}{2}}} e^{-\frac{|x|^2}{4t}} & \text{if } (t, x) \in]0, +\infty[\times \mathbb{R}^n, \\ 0 & \text{if } (t, x) \in (]-\infty, 0] \times \mathbb{R}^n) \setminus \{(0, 0)\}. \end{cases}$$

Throughout the paper we find convenient to adopt the following notation: if D is a subset of \mathbb{R}^n , $T \in]0, +\infty[$ and h is a map from D to \mathbb{R}^n , we denote by h^T the map from $[0, T] \times D$ to $[0, T] \times \mathbb{R}^n$ defined by

$$h^T(t, x) := (t, h(x)) \quad \forall (t, x) \in [0, T] \times D.$$

Let $\phi \in \mathcal{A}_{\partial\Omega}$ and let μ be a continuous function from $[0, T] \times \partial\Omega$ to \mathbb{R} . In order to work with a space of densities that is not ϕ -dependent, it makes sense to consider the single layer potential with density $\mu \circ (\phi^T)^{(-1)}$:

$$\begin{aligned} & v[\mu \circ (\phi^T)^{(-1)}](t, x) \\ & := \int_0^t \int_{\phi(\partial\Omega)} S_n(t - \tau, x - y) \mu \circ (\phi^T)^{(-1)}(\tau, y) d\sigma_y d\tau \\ & = \int_0^t \int_{\phi(\partial\Omega)} S_n(t - \tau, x - y) \mu(\tau, \phi^{(-1)}(y)) d\sigma_y d\tau \quad \forall (t, x) \in [0, T] \times \mathbb{R}^n. \end{aligned}$$

Moreover, in the applications it is often convenient to consider the boundary integral operator associated with the ϕ -pullback of the single layer potential, that is

$$V_\phi[\mu] := v[\mu \circ (\phi^T)^{(-1)}] \circ \phi^T \quad \text{on } [0, T] \times \partial\Omega. \quad (1)$$

We also consider the maps

$$V_{l,\phi}[\mu](t, x) := \int_0^t \int_{\phi(\partial\Omega)} \partial_{x_l} S_n(t - \tau, \phi(x) - y) \mu \circ (\phi^T)^{(-1)}(\tau, y) d\sigma_y d\tau, \quad (2)$$

$$W_{*,\phi}[\mu](t, x) := \int_0^t \int_{\phi(\partial\Omega)} D_x S_n(t - \tau, \phi(x) - y) \cdot \nu_\phi(x) \mu \circ (\phi^T)^{(-1)}(\tau, y) d\sigma_y d\tau, \quad (3)$$

$$W_\phi[\mu](t, x) := - \int_0^t \int_{\phi(\partial\Omega)} D_x S_n(t - \tau, \phi(x) - y) \cdot \nu_\phi(y) \mu \circ (\phi^T)^{(-1)}(\tau, y) d\sigma_y d\tau, \quad (4)$$

for all $(t, x) \in [0, T] \times \partial\Omega$ and all $l \in \{1, \dots, n\}$. Here above, $\partial_{x_l} S_n$ denotes the x_l -derivate of S_n and $D_x S_n$ is the gradient of S_n with respect to the spatial variables, whereas ν_ϕ denotes the exterior unit normal field to $\phi(\partial\Omega)$. The maps $V_{l,\phi}$ and $W_{*,\phi}$ are associated with the ϕ -pullback of the x_l and normal derivatives of the single layer potential, respectively, and the map W_ϕ is associated with the ϕ -pullback of the double layer potential. The main result of this paper states that the maps from a certain subset of $\mathcal{A}_{\partial\Omega}$ to suitable spaces of operators, which take ϕ to V_ϕ , $V_{l,\phi}$, $W_{*,\phi}$, and W_ϕ , belong to the class C^∞ (see Theorem 5.4).

Our strategy to prove the result proceeds as follows. First, we characterize layer potentials as solutions of a transmission problem on a ϕ -dependent domain. We then pullback this problem to a fixed domain and obtain a new abstract ϕ -dependent problem on a fixed set. Finally, we use the real analyticity of the inversion map to establish the regularity upon shape perturbations. It's worth noting that the non-homogeneous term in the heat equation for the transmission problem includes a distributional term of the form $\partial_t f$, as described in Theorem 3.1 and Lemma 4.2. Standard results in parabolic theory do not cover this case, hence we relied on new results on the heat volume potential that the second author developed for this purpose and presented in [35]. We acknowledge that a direct change of variable, as described in Chapko, Kress and Yoon [5, 6], can avoid the issue of $\partial_t f$ in the equation. However, their approach requires the boundary to be C^2 , and it does not work in the optimal Hölder setting, i.e. with sets of class $C^{1,\alpha}$. In contrast, our approach maintains sharp assumptions on the regularity of sets and perturbations.

The strategy we have just described stems from the approach proposed by Lanza de Cristoforis and Rossi [29, 30] to analyze the shape dependence of layer potentials for the Laplace and Helmholtz operators. The extension to the heat equation is, however, by no means straightforward. On the contrary, we have to be very careful when dealing with the space-time anisotropy and we have to identify a suitable functional setting with the correct regularity for the space and the time components.

We also observe that, instead of the real and complex analyticity results that are typical in the elliptic case (see, e.g., [8] and [10]) we have to content ourselves with a smoothness result. This seems to be an unavoidable issue for the heat equation, and its origin can be traced to the regularity of the fundamental solution S_n . Specifically, S_n is real analytic in x for a fixed $t \neq 0$, but only C^∞ in t for a fixed $x \neq 0$.

In some forthcoming papers, we plan to use the results presented here to study the shape sensitivity of the solutions to linear and nonlinear boundary value problems for the heat equation.

The paper is organized as follows. In Section 2 we introduce some notation and present some known preliminary results: in Section 2.1 we fix some standard notation and definitions, in Section 2.2 we introduce certain parabolic Schauder spaces, in Section 2.3 we define the class of diffeomorphisms $\mathcal{A}_{\partial\Omega}$ and present some related results, and, finally, in Section 2.4 we recall the definition of layer heat potentials and their basic properties. In Section 3 we prove the unique solvability of a certain auxiliary transmission problem. This is the problem that later we use to characterize the layer potentials. Then, in Section 4, we ϕ -pullback the problem to a fixed domain and we analyze the dependence of the

pulled-back problem upon the shape parameter ϕ . Finally, Section 5 contains our main results on the dependence of the layer potentials upon ϕ . In Appendix A we collect some technical results on composition and integral operators.

2 Preliminaries

2.1 Some standard notation

For standard definitions of calculus in normed spaces, and in particular for the definition and properties of real analytic operators, we refer to Deimling [14]. The inverse of an invertible function f is denoted by $f^{(-1)}$, while the reciprocal of a complex-valued function g is denoted by g^{-1} . If A is a matrix, then A^\top denotes the transpose matrix of A and A_{ij} denotes the (i, j) -entry of A . If A is invertible, A^{-1} is the inverse of A and we set $A^{-\top} := (A^{-1})^\top$.

The symbol \mathbb{N} denotes the set of natural numbers including 0. Throughout the paper,

$$n \in \mathbb{N} \setminus \{0, 1\}$$

denotes the dimension of the Euclidean ambient space \mathbb{R}^n . If $\mathbb{D} \subseteq \mathbb{R}^n$, then $\overline{\mathbb{D}}$ denotes the closure of \mathbb{D} and $\partial\mathbb{D}$ denotes the boundary of \mathbb{D} . If \mathcal{X} is a normed space, then $B(\mathbb{D}, \mathcal{X})$ and $C^0(\mathbb{D}, \mathcal{X})$ denote the space of bounded and continuous functions from \mathbb{D} to \mathcal{X} , respectively. As usual, we equip $B(\mathbb{D}, \mathcal{X})$ with the sup-norm and we set $C_b^0(\mathbb{D}, \mathcal{X}) := C^0(\mathbb{D}, \mathcal{X}) \cap B(\mathbb{D}, \mathcal{X})$. If \mathcal{Y} is also a normed space, then $\mathcal{L}(\mathcal{X}, \mathcal{Y})$ denotes the space of bounded linear operators from \mathcal{X} to \mathcal{Y} equipped with the usual operator norm.

Let Ω be an open subset of \mathbb{R}^n . Then $\Omega^- := \mathbb{R}^n \setminus \overline{\Omega}$ denotes the exterior of Ω . Let $m \in \mathbb{N}$. The space of m -times continuously differentiable real-valued functions on Ω is denoted by $C^m(\Omega)$. Let $f \in C^m(\Omega)$. Then Df denotes the Jacobian matrix of f . The subspace of $C^m(\Omega)$ of those functions f whose derivatives $D^\eta f$ of order $|\eta| \leq m$ can be extended with continuity to $\overline{\Omega}$ is denoted $C^m(\overline{\Omega})$. The subspace of $C^m(\overline{\Omega})$ whose derivatives up to the m order are bounded is denoted $C_b^m(\overline{\Omega})$. As is well known, $C_b^m(\overline{\Omega})$ equipped with the norm $\|f\|_{C_b^m(\overline{\Omega})} := \sum_{|\eta| \leq m} \sup_{\overline{\Omega}} |D^\eta f|$ is a Banach space. The symbol ν_Ω denotes the outward unit normal field to $\partial\Omega$, where it exists.

Let $m \in \mathbb{N}$ and $\alpha \in]0, 1[$. For the definition of open subsets of \mathbb{R}^n of class C^m and $C^{m, \alpha}$, and of the Schauder spaces $C^{m, \alpha}(\overline{\Omega})$ and $C^{m, \alpha}(\partial\Omega)$, we refer to Gilbarg and Trudinger [16, pp. 52, 95].

2.2 Parabolic Schauder spaces

Let $\alpha, \beta \in]0, 1[$, $T \in]0, +\infty[$ and $\mathbb{D} \subseteq \mathbb{R}^n$. Then $C^{\alpha; \beta}([0, T] \times \mathbb{D})$ denotes the space of bounded continuous functions u from $[0, T] \times \mathbb{D}$ to \mathbb{R} such that

$$\begin{aligned} \|u\|_{C^{\alpha; \beta}([0, T] \times \mathbb{D})} &:= \sup_{[0, T] \times \mathbb{D}} |u| + \sup_{\substack{t_1, t_2 \in [0, T] \\ t_1 \neq t_2}} \sup_{x \in \mathbb{D}} \frac{|u(t_1, x) - u(t_2, x)|}{|t_1 - t_2|^\alpha} \\ &+ \sup_{t \in [0, T]} \sup_{\substack{x_1, x_2 \in \mathbb{D} \\ x_1 \neq x_2}} \frac{|u(t, x_1) - u(t, x_2)|}{|x_1 - x_2|^\beta} < +\infty. \end{aligned}$$

Let now Ω be an open subset of \mathbb{R}^n . Then $C^{\frac{1+\alpha}{2}; 1+\alpha}([0, T] \times \overline{\Omega})$ denotes the space of bounded continuous functions u from $[0, T] \times \overline{\Omega}$ to \mathbb{R} which are continuously differentiable with respect to the space variables and such that

$$\|u\|_{C^{\frac{1+\alpha}{2}; 1+\alpha}([0, T] \times \overline{\Omega})} := \sup_{[0, T] \times \overline{\Omega}} |u| + \sum_{i=1}^n \|\partial_{x_i} u\|_{C^{\frac{\alpha}{2}; \alpha}([0, T] \times \overline{\Omega})}$$




Figure 1: The diffeomorphism ϕ and the ϕ -dependent sets $\mathbb{I}[\phi]$, $\mathbb{E}[\phi]$ and $\phi(\partial\Omega)$.

$$+ \sup_{\substack{t_1, t_2 \in [0, T] \\ t_1 \neq t_2}} \sup_{x \in \bar{\Omega}} \frac{|u(t_1, x) - u(t_2, x)|}{|t_1 - t_2|^{\frac{1+\alpha}{2}}} < +\infty.$$

If Ω is of class $C^{1,\alpha}$, we can use the local parametrization of $\partial\Omega$ to define the space $C^{\frac{1+\alpha}{2}; 1+\alpha}([0, T] \times \partial\Omega)$ in the natural way. Similarly, we can define the spaces $C^{m,\alpha}(M)$ and $C^{\frac{m+\alpha}{2}; m+\alpha}([0, T] \times M)$, $m = 0, 1$ on a manifold M of class $C^{m,\alpha}$ imbedded in \mathbb{R}^n (for details, see Appendix A).

Finally, we use the subscript 0 to denote a subspace consisting of functions that are zero at $t = 0$. For example,

$$C_0^{\alpha;\beta}([0, T] \times \mathbb{D}) := \left\{ u \in C^{\alpha;\beta}([0, T] \times \mathbb{D}) : u(0, x) = 0 \quad \forall x \in \mathbb{D} \right\}.$$

Then the spaces $C_0^{\frac{1+\alpha}{2}; 1+\alpha}([0, T] \times \bar{\Omega})$, $C_0^{\frac{1+\alpha}{2}; 1+\alpha}([0, T] \times \partial\Omega)$, and $C_0^{\frac{m+\alpha}{2}; m+\alpha}([0, T] \times M)$ are similarly defined.

For a comprehensive introduction to parabolic Schauder spaces we refer the reader to classical monographs on the field, for example Ladyženskaja, Solonnikov and Ural'ceva [21, Ch. 1] (see also [25, 26]).

2.3 The class of diffeomorphisms $\mathcal{A}_{\partial\Omega}$

We now introduce the class of diffeomorphisms that we use to model the domains' shape.

Definition 2.1. Let Ω be a bounded open connected subset of \mathbb{R}^n of class C^1 . We denote by $\mathcal{A}_{\partial\Omega}$ the set consisting of the functions of class $C^1(\partial\Omega, \mathbb{R}^n)$ that are injective and whose differential is injective at all points of $\partial\Omega$ and, similarly, we denote by $\mathcal{A}_{\bar{\Omega}}$ the set of functions of $C^1(\bar{\Omega}, \mathbb{R}^n)$ that are injective and whose differential is injective at all points of $\bar{\Omega}$.

We can verify that $\mathcal{A}_{\partial\Omega}$ and $\mathcal{A}_{\bar{\Omega}}$ are open in $C^1(\partial\Omega, \mathbb{R}^n)$ and $C^1(\bar{\Omega}, \mathbb{R}^n)$, respectively (see, e.g., Lanza de Cristoforis and Rossi [30, Lem. 2.2, p. 197] and [29, Lem. 2.5, p. 143]). If Ω has connected exterior Ω^- , then $\mathbb{R}^n \setminus \partial\Omega$ has two open connected components and thus the Jordan-Leray separation theorem ensures that $\mathbb{R}^n \setminus \phi(\partial\Omega)$ has exactly two open connected components for all $\phi \in \mathcal{A}_{\partial\Omega}$ (see, e.g., Deimling [14, Thm 5.2, p.26]). One of these open connected components is bounded, and we denote it by $\mathbb{I}[\phi]$, the letter “I” standing for “interior.” The other one is unbounded, and we denote it by $\mathbb{E}[\phi]$, the letter “E” standing for “exterior.” (See Figure 1.)

We need to recall two technical lemmas of Lanza de Cristoforis and Rossi [30, §2], which show that a diffeomorphism on $\partial\Omega$ can be extended in a neighborhood of $\partial\Omega$ by means of a real analytic extension operator.

Lemma 2.2. Let $\alpha \in]0, 1[$. Let Ω be a bounded open connected subset of \mathbb{R}^n of class $C^{1,\alpha}$ such that Ω^- is connected. There exists $\omega \in C^{1,\alpha}(\partial\Omega, \mathbb{R}^n)$ such that $|\omega| = 1$ and $\omega \cdot \nu_\Omega > 1/2$ on $\partial\Omega$. Moreover, the following statements hold.

(i) There exists $\delta_\Omega \in]0, +\infty[$ such that the sets

$$\begin{aligned}\Omega_{\omega,\delta} &:= \{x + s\omega(x) : x \in \partial\Omega, s \in]-\delta, \delta[\}, \\ \Omega_{\omega,\delta}^+ &:= \{x + s\omega(x) : x \in \partial\Omega, s \in]-\delta, 0[\}, \\ \Omega_{\omega,\delta}^- &:= \{x + s\omega(x) : x \in \partial\Omega, s \in]0, \delta[\}\end{aligned}$$

are connected and of class $C^{1,\alpha}$. They have boundaries

$$\begin{aligned}\partial\Omega_{\omega,\delta} &= \{x + s\omega(x) : x \in \partial\Omega, s \in \{-\delta, \delta\}\}, \\ \partial\Omega_{\omega,\delta}^+ &= \{x + s\omega(x) : x \in \partial\Omega, s \in \{-\delta, 0\}\}, \\ \partial\Omega_{\omega,\delta}^- &= \{x + s\omega(x) : x \in \partial\Omega, s \in \{0, \delta\}\}.\end{aligned}$$

and we have $\Omega_{\omega,\delta}^+ \subseteq \Omega$ and $\Omega_{\omega,\delta}^- \subseteq \Omega^-$ for all $\delta \in]0, \delta_\Omega[$.

(ii) Let $\delta \in]0, \delta_\Omega[$. If $\Phi \in \mathcal{A}_{\overline{\Omega_{\omega,\delta}}}$, then $\phi := \Phi|_{\partial\Omega} \in \mathcal{A}_{\partial\Omega}$.

(iii) If $\delta \in]0, \delta_\Omega[$, then the set

$$\mathcal{A}'_{\overline{\Omega_{\omega,\delta}}} := \{\Phi \in \mathcal{A}_{\overline{\Omega_{\omega,\delta}}} : \Phi(\Omega_{\omega,\delta}^+) \subseteq \mathbb{I}[\Phi|_{\partial\Omega}]\}$$

is open in $\mathcal{A}_{\overline{\Omega_{\omega,\delta}}}$ and $\Phi(\Omega_{\omega,\delta}^-) \subseteq \mathbb{E}[\Phi|_{\partial\Omega}]$ for all $\Phi \in \mathcal{A}'_{\overline{\Omega_{\omega,\delta}}}$.

(iv) If $\delta \in]0, \delta_\Omega[$ and $\Phi \in C^{1,\alpha}(\overline{\Omega_{\omega,\delta}}, \mathbb{R}^n) \cap \mathcal{A}'_{\overline{\Omega_{\omega,\delta}}}$, then both $\Phi(\Omega_{\omega,\delta}^+)$ and $\Phi(\Omega_{\omega,\delta}^-)$ are open sets of class $C^{1,\alpha}$, and

$$\partial\Phi(\Omega_{\omega,\delta}^+) = \Phi(\partial\Omega_{\omega,\delta}^+), \quad \partial\Phi(\Omega_{\omega,\delta}^-) = \Phi(\partial\Omega_{\omega,\delta}^-).$$

Lemma 2.3. Let $\Omega, \omega, \delta_\Omega$ be as in Lemma 2.2. Let $\phi_0 \in C^{1,\alpha}(\partial\Omega, \mathbb{R}^n) \cap \mathcal{A}_{\partial\Omega}$. Then the following statements hold.

- (i) There exists $\delta_0 \in]0, \delta_\Omega[$ and $\Phi_0 \in C^{1,\alpha}(\overline{\Omega_{\omega,\delta_0}}, \mathbb{R}^n) \cap \mathcal{A}'_{\overline{\Omega_{\omega,\delta_0}}}$ such that $\phi_0 = \Phi_0|_{\partial\Omega}$.
- (ii) Let δ_0, Φ_0 be as in statement (i). Then there exist an open neighborhood \mathcal{W}_0 of ϕ_0 in $C^{1,\alpha}(\partial\Omega, \mathbb{R}^n) \cap \mathcal{A}_{\partial\Omega}$, and a real analytic extension operator \mathbf{E} from $C^{1,\alpha}(\partial\Omega, \mathbb{R}^n)$ to $C^{1,\alpha}(\overline{\Omega_{\omega,\delta_0}}, \mathbb{R}^n) \cap \mathcal{A}'_{\overline{\Omega_{\omega,\delta_0}}}$ which maps \mathcal{W}_0 to $C^{1,\alpha}(\overline{\Omega_{\omega,\delta_0}}, \mathbb{R}^n) \cap \mathcal{A}'_{\overline{\Omega_{\omega,\delta_0}}}$ and such that $\mathbf{E}[\phi_0] = \Phi_0$ and $\mathbf{E}[\phi]|_{\partial\Omega} = \phi$, for all $\phi \in \mathcal{W}_0$.

Finally, we present in the following technical lemma two real analyticity results, one for a map related to the change of variables in integrals, and one for the pullback of the outer normal field. For a proof we refer to Lanza de Cristoforis and Rossi [29, p. 166] and to Lanza de Cristoforis [24, Prop. 1]. Throughout the paper ν_ϕ denotes the exterior unit normal field to $\partial\mathbb{I}[\phi] = \phi(\partial\Omega)$.

Lemma 2.4. Let Ω be as in Lemma 2.2. Then the following statements hold.

(i) For each $\phi \in C^{1,\alpha}(\partial\Omega, \mathbb{R}^n) \cap \mathcal{A}_{\partial\Omega}$ there exists a unique $\tilde{\sigma}_n[\phi] \in C^{1,\alpha}(\partial\Omega)$ such that

$$\int_{\phi(\partial\Omega)} f(s) d\sigma_s = \int_{\partial\Omega} f \circ \phi(y) \tilde{\sigma}_n[\phi](y) d\sigma_y \quad \forall f \in L^1(\phi(\partial\Omega)).$$

Moreover, $\tilde{\sigma}_n[\phi] > 0$ and the map that takes ϕ to $\tilde{\sigma}_n[\phi]$ is real analytic from $C^{1,\alpha}(\partial\Omega, \mathbb{R}^n) \cap \mathcal{A}_{\partial\Omega}$ to $C^{0,\alpha}(\partial\Omega)$.

(ii) The map from $C^{1,\alpha}(\partial\Omega, \mathbb{R}^n) \cap \mathcal{A}_{\partial\Omega}$ to $C^{0,\alpha}(\partial\Omega, \mathbb{R}^n)$ that takes ϕ to $\nu_\phi \circ \phi$ is real analytic.

2.4 Layer heat potentials

In this section we collect some well-known facts on the layer heat potentials. For proofs and detailed references we refer to Ladyženskaja, Solonnikov and Ural'ceva [21] and [26, 34]. In Theorem 2.5 we introduce the double layer potential and describe some of its main properties.

Theorem 2.5. *Let $\alpha \in]0, 1[$. Let $T \in]0, +\infty[$. Let Ω be a bounded open subset of \mathbb{R}^n of class $C^{1,\alpha}$. The double layer potential with density $\mu \in C_0^{\frac{\alpha}{2};\alpha}([0, T] \times \partial\Omega)$ is the function $w[\mu]$ from $[0, T] \times \mathbb{R}^n$ to \mathbb{R} defined by*

$$w[\mu](t, x) := \int_0^t \int_{\partial\Omega} \frac{\partial}{\partial\nu_\Omega(y)} S_n(t - \tau, x - y) \mu(\tau, y) d\sigma_y d\tau \quad \forall (t, x) \in [0, T] \times \mathbb{R}^n.$$

For the double layer potential the following statements hold.

- (i) $w[\mu]$ solves the heat equation in $[0, T] \times (\mathbb{R}^n \setminus \partial\Omega)$.
- (ii) The restriction $w[\mu]|_{[0, T] \times \Omega}$ has a unique extension to a continuous function $w^+[\mu]$ from $[0, T] \times \overline{\Omega}$ to \mathbb{R} and the restriction $w[\mu]|_{[0, T] \times \Omega^-}$ has a unique extension to a continuous function $w^-[\mu]$ from $[0, T] \times \overline{\Omega^-}$ to \mathbb{R} .
- (iii) Unless $\mu = 0$, $w[\mu]$ is not continuous on the boundary $[0, T] \times \partial\Omega$ and we have the jump formula

$$w^\pm[\mu] = \mp \frac{1}{2} \mu + w[\mu] \quad \text{on } [0, T] \times \partial\Omega.$$

In addition, for the normal derivative of $w[\mu]$ we have

$$\frac{\partial}{\partial\nu_\Omega} w^+[\mu] = \frac{\partial}{\partial\nu_\Omega} w^-[\mu] \quad \text{on } [0, T] \times \partial\Omega.$$

- (iv) The map from $C_0^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times \partial\Omega)$ to $C_0^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times \overline{\Omega})$ that takes μ to $w^+[\mu]$ is linear and continuous. If $R > 0$ is such that $\overline{\Omega} \subseteq \mathbb{B}_n(0, R)$, then the map from $C_0^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times \partial\Omega)$ to $C_0^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times (\overline{\mathbb{B}_n(0, R)} \setminus \Omega))$ that takes μ to $w^-[\mu]|_{[0, T] \times (\overline{\mathbb{B}_n(0, R)} \setminus \Omega)}$ is linear and continuous.

In the following Theorem 2.6 we introduce the single layer potential.

Theorem 2.6. *Let $\alpha \in]0, 1[$. Let $T \in]0, +\infty[$. Let Ω be a bounded open subset of \mathbb{R}^n of class $C^{1,\alpha}$. The single layer potential with density $\mu \in C_0^{\frac{\alpha}{2};\alpha}([0, T] \times \partial\Omega)$ is the function $v[\mu]$ from $[0, T] \times \mathbb{R}^n$ to \mathbb{R} defined by*

$$v[\mu](t, x) := \int_0^t \int_{\partial\Omega} S_n(t - \tau, x - y) \mu(\tau, y) d\sigma_y d\tau \quad \forall (t, x) \in [0, T] \times \mathbb{R}^n.$$

For the single layer potential the following statements hold.

- (i) $v[\mu]$ solves the heat equation in $[0, T] \times (\mathbb{R}^n \setminus \partial\Omega)$.
- (ii) $v[\mu]$ is continuous in $[0, T] \times \mathbb{R}^n$.
- (iii) Let $v^+[\mu]$ and $v^-[\mu]$ denote the restriction of $v[\mu]$ to $[0, T] \times \overline{\Omega}$ and to $[0, T] \times \overline{\Omega^-}$. Then the following jump relations hold

$$\begin{aligned} \frac{\partial}{\partial\nu_\Omega(x)} v^\pm[\mu](t, x) &= \pm \frac{1}{2} \mu(t, x) + \int_0^t \int_{\partial\Omega} \frac{\partial}{\partial\nu_\Omega(x)} S_n(t - \tau, x - y) \mu(\tau, y) d\sigma_y d\tau, \\ \frac{\partial}{\partial x_r} v^\pm[\mu](t, x) &= \pm \frac{1}{2} \mu(t, x) (\nu_\Omega)_r(x) + \int_0^t \int_{\partial\Omega} \frac{\partial}{\partial x_r} S_n(t - \tau, x - y) \mu(\tau, y) d\sigma_y d\tau, \end{aligned}$$

for all $(t, x) \in [0, T] \times \partial\Omega$ and all $r \in \{1, \dots, n\}$.

(iv) The map from $C_0^{\frac{\alpha}{2};\alpha}([0, T] \times \partial\Omega)$ to $C_0^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times \overline{\Omega})$ that takes μ to $v^+[\mu]$ is linear and continuous. If $R > 0$ and $\overline{\Omega} \subseteq \mathbb{B}_n(0, R)$, then the map from $C_0^{\frac{\alpha}{2};\alpha}([0, T] \times \partial\Omega)$ to $C_0^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times (\overline{\mathbb{B}_n(0, R)} \setminus \Omega))$ that takes μ to $v^-[\mu]_{|[0, T] \times (\overline{\mathbb{B}_n(0, R)} \setminus \Omega)}$ is linear and continuous.

3 An auxiliary transmission problem

In the following Theorem 3.1 we introduce an auxiliary transmission problem and prove the existence and uniqueness of its solution. We will use such problem to characterize the layer heat potentials supported on $\Phi(\partial\Omega)$.

To shorten our statements we find convenient to introduce the following notation: If $\alpha \in]0, 1[$, $T \in]0, +\infty[$, Ω , ω and δ_Ω are as in Lemma 2.2, $\delta \in]0, \delta_\Omega[$, and $\Phi \in C^{1,\alpha}(\overline{\Omega_{\omega,\delta}}, \mathbb{R}^n) \cap \mathcal{A}'_{\Omega_{\omega,\delta}}$, then \mathcal{S}_Φ denotes the product space

$$\begin{aligned} \mathcal{S}_\Phi := & C_0^{\frac{1+\alpha}{2};\alpha} \left([0, T] \times \overline{\Phi(\Omega_{\omega,\delta}^+)} \right) \times C_0^{\frac{1+\alpha}{2};\alpha} \left([0, T] \times \overline{\Phi(\Omega_{\omega,\delta}^-)} \right) \\ & \times C^{\frac{\alpha}{2};\alpha} \left([0, T] \times \overline{\Phi(\Omega_{\omega,\delta}^+), \mathbb{R}^n} \right) \times C^{\frac{\alpha}{2};\alpha} \left([0, T] \times \overline{\Phi(\Omega_{\omega,\delta}^-, \mathbb{R}^n)} \right) \\ & \times C_0^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times \Phi(\partial\Omega)) \times C_0^{\frac{\alpha}{2};\alpha}([0, T] \times \Phi(\partial\Omega)) \\ & \times C_0^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times \Phi(\partial\Omega_{\omega,\delta}^+ \setminus \partial\Omega)) \times C_0^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times \Phi(\partial\Omega_{\omega,\delta}^- \setminus \partial\Omega)). \end{aligned} \quad (5)$$

Moreover, we set

$$B[\Phi](v^+, v^-) := (Dv^+)_{|[0, T] \times \Phi(\partial\Omega)} \nu_{\Phi|_{\partial\Omega}} - (Dv^-)_{|[0, T] \times \Phi(\partial\Omega)} \nu_{\Phi|_{\partial\Omega}},$$

for all $(v^+, v^-) \in C_0^{\frac{1+\alpha}{2};1+\alpha} \left([0, T] \times \overline{\Phi(\Omega_{\omega,\delta}^+)} \right) \times C_0^{\frac{1+\alpha}{2};1+\alpha} \left([0, T] \times \overline{\Phi(\Omega_{\omega,\delta}^-)} \right)$.

We are now ready to state Theorem 3.1.

Theorem 3.1. *Let $\alpha \in]0, 1[$, $T \in]0, +\infty[$. Let Ω , ω , δ_Ω be as in Lemma 2.2. Let $\delta \in]0, \delta_\Omega[$ and $\Phi \in C^{1,\alpha}(\Omega_{\omega,\delta}, \mathbb{R}^n) \cap \mathcal{A}'_{\Omega_{\omega,\delta}}$. Then the transmission problem*

$$\begin{cases} \partial_t v^+ - \Delta v^+ = \partial_t f_0^+ + \operatorname{div} f_1^+ & \text{in }]0, T[\times \Phi(\Omega_{\omega,\delta}^+), \\ \partial_t v^- - \Delta v^- = \partial_t f_0^- + \operatorname{div} f_1^- & \text{in }]0, T[\times \Phi(\Omega_{\omega,\delta}^-), \\ v^+ - v^- = g & \text{on } [0, T] \times \Phi(\partial\Omega), \\ B[\Phi](v^+, v^-) = g_1 & \text{on } [0, T] \times \Phi(\partial\Omega), \\ v^+ = h^+ & \text{on } [0, T] \times \Phi(\partial\Omega_{\omega,\delta}^+ \setminus \partial\Omega), \\ v^- = h^- & \text{on } [0, T] \times \Phi(\partial\Omega_{\omega,\delta}^- \setminus \partial\Omega), \\ v^+(0, \cdot) = 0 & \text{in } \overline{\Phi(\Omega_{\omega,\delta}^+)}, \\ v^-(0, \cdot) = 0 & \text{in } \overline{\Phi(\Omega_{\omega,\delta}^-)}, \end{cases} \quad (6)$$

has a unique solution (v^+, v^-) in $C_0^{\frac{1+\alpha}{2};1+\alpha} \left([0, T] \times \overline{\Phi(\Omega_{\omega,\delta}^+)} \right) \times C_0^{\frac{1+\alpha}{2};1+\alpha} \left([0, T] \times \overline{\Phi(\Omega_{\omega,\delta}^-)} \right)$ for each given $(f_0^+, f_0^-, f_1^+, f_1^-, g, g_1, h^+, h^-) \in \mathcal{S}_\Phi$.

We observe that the heat equations in (6) are to be understood in the weak sense of distributions in every instance.

Proof. First we consider the existence of a solution of the transmission problem (6). Let $(f_0^+, f_0^-, f_1^+, f_1^-, g, g_1, h^+, h^-) \in \mathcal{S}_\Phi$. In particular $f_0^\pm \in C_0^{\frac{1+\alpha}{2};\alpha} \left([0, T] \times \overline{\Phi(\Omega_{\omega,\delta}^\pm)} \right)$. By

[35, Thm 4.4] there exist functions $P[\partial_t f_0^\pm] \in C_0^{\frac{1+\alpha}{2}; 1+\alpha} \left([0, T] \times \overline{\Phi(\Omega_{\omega, \delta}^\pm)} \right)$ —which are the heat volume potentials with densities $\partial_t f_0^\pm$ —that solve the equations

$$\partial_t u - \Delta u = \partial_t f_0^\pm \quad \text{in }]0, T[\times \Phi(\Omega_{\omega, \delta}^\pm)$$

in the sense of distributions. Then, the proof of the existence of a solution of problem (6) can be reduced to that of the existence of a solution for

$$\begin{cases} \partial_t v^+ - \Delta v^+ = \operatorname{div} f_1^+ & \text{in }]0, T[\times \Phi(\Omega_{\omega, \delta}^+), \\ \partial_t v^- - \Delta v^- = \operatorname{div} f_1^- & \text{in }]0, T[\times \Phi(\Omega_{\omega, \delta}^-), \\ v^+ - v^- = g & \text{on } [0, T] \times \Phi(\partial\Omega), \\ B[\Phi](v^+, v^-) = g_1 & \text{on } [0, T] \times \Phi(\partial\Omega), \\ v^+ = h^+ & \text{on } [0, T] \times \Phi(\partial\Omega_{\omega, \delta}^+ \setminus \partial\Omega), \\ v^- = h^- & \text{on } [0, T] \times \Phi(\partial\Omega_{\omega, \delta}^- \setminus \partial\Omega), \\ v^+(0, \cdot) = 0 & \text{in } \overline{\Phi(\Omega_{\omega, \delta}^+)}, \\ v^-(0, \cdot) = 0 & \text{in } \overline{\Phi(\Omega_{\omega, \delta}^-)}, \end{cases} \quad (7)$$

with a possibly different array of data $(f_1^+, f_1^-, g, g_1, h^+, h^-)$. By known results in parabolic theory (see Lunardi and Vespi [32, Thm. 4.3] and Lieberman [31, Thm. 6.48]) there exists a pair $(\tilde{v}^+, \tilde{v}^-) \in C_0^{\frac{1+\alpha}{2}; 1+\alpha} \left([0, T] \times \overline{\Phi(\Omega_{\omega, \delta}^+)} \right) \times C_0^{\frac{1+\alpha}{2}; 1+\alpha} \left([0, T] \times \overline{\Phi(\Omega_{\omega, \delta}^-)} \right)$ such that

$$\begin{cases} \partial_t \tilde{v}^+ - \Delta \tilde{v}^+ = \operatorname{div} f_1^+ & \text{in }]0, T[\times \Phi(\Omega_{\omega, \delta}^+), \\ \tilde{v}^+ = g & \text{on } [0, T] \times \Phi(\partial\Omega), \\ \tilde{v}^+ = h^+ & \text{on } [0, T] \times \Phi(\partial\Omega_{\omega, \delta}^+ \setminus \partial\Omega), \\ \tilde{v}^+(0, \cdot) = 0 & \text{in } \overline{\Phi(\Omega_{\omega, \delta}^+)}, \end{cases}$$

and

$$\begin{cases} \partial_t \tilde{v}^- - \Delta \tilde{v}^- = \operatorname{div} f_1^- & \text{in }]0, T[\times \Phi(\Omega_{\omega, \delta}^-), \\ \tilde{v}^- = 0 & \text{on } [0, T] \times \Phi(\partial\Omega), \\ \tilde{v}^- = h^- & \text{on } [0, T] \times \Phi(\partial\Omega_{\omega, \delta}^- \setminus \partial\Omega), \\ \tilde{v}^-(0, \cdot) = 0 & \text{in } \overline{\Phi(\Omega_{\omega, \delta}^-)}. \end{cases}$$

Moreover, by Theorem 2.6, the pair of functions $(u^+, u^-) \in C_0^{\frac{1+\alpha}{2}; 1+\alpha} \left([0, T] \times \overline{\Phi(\Omega_{\omega, \delta}^+)} \right) \times C_0^{\frac{1+\alpha}{2}; 1+\alpha} \left([0, T] \times \overline{\Phi(\Omega_{\omega, \delta}^-)} \right)$ defined by

$$u^\pm := v^\pm[\mu] \quad \text{in } [0, T] \times \overline{\Phi(\Omega_{\omega, \delta}^\pm)},$$

with

$$\mu := -g_1 + B[\Phi](\tilde{v}^+, \tilde{v}^-) \quad \text{on } [0, T] \times \Phi(\partial\Omega),$$

is a solution of the transmission problem

$$\begin{cases} \partial_t u^+ - \Delta u^+ = 0 & \text{in }]0, T[\times \Phi(\Omega_{\omega, \delta}^+), \\ \partial_t u^- - \Delta u^- = 0 & \text{in }]0, T[\times \Phi(\Omega_{\omega, \delta}^-), \\ u^+ - u^- = 0 & \text{on } [0, T] \times \Phi(\partial\Omega), \\ B[\phi](u^+, u^-) = -g_1 + B[\Phi](\tilde{v}^+, \tilde{v}^-) & \text{on } [0, T] \times \Phi(\partial\Omega), \\ u^+(0, \cdot) = 0 & \text{in } \overline{\Phi(\Omega_{\omega, \delta}^+)}, \\ u^-(0, \cdot) = 0 & \text{in } \overline{\Phi(\Omega_{\omega, \delta}^-)}. \end{cases}$$

Then, if the boundary value problem

$$\begin{cases} \partial_t \hat{v}^+ - \Delta \hat{v}^+ = 0 & \text{in }]0, T[\times \Phi(\Omega_{\omega, \delta}^+), \\ \partial_t \hat{v}^- - \Delta \hat{v}^- = 0 & \text{in }]0, T[\times \Phi(\Omega_{\omega, \delta}^-), \\ \hat{v}^+ - \hat{v}^- = 0 & \text{on } [0, T] \times \Phi(\partial\Omega), \\ B[\phi](\hat{v}^+, \hat{v}^-) = 0 & \text{on } [0, T] \times \Phi(\partial\Omega), \\ \hat{v}^+ = u^+ & \text{on } [0, T] \times \Phi(\partial\Omega_{\omega, \delta}^+ \setminus \partial\Omega), \\ \hat{v}^- = u^- & \text{on } [0, T] \times \Phi(\partial\Omega_{\omega, \delta}^- \setminus \partial\Omega), \\ \hat{v}^+(0, \cdot) = 0 & \text{in } \overline{\Phi(\Omega_{\omega, \delta}^+)}, \\ \hat{v}^-(0, \cdot) = 0 & \text{in } \overline{\Phi(\Omega_{\omega, \delta}^-)}, \end{cases} \quad (8)$$

has a solution $(\hat{v}^+, \hat{v}^-) \in C_0^{\frac{1+\alpha}{2}; 1+\alpha} \left([0, T] \times \overline{\Phi(\Omega_{\omega, \delta}^+)} \right) \times C_0^{\frac{1+\alpha}{2}; 1+\alpha} \left([0, T] \times \overline{\Phi(\Omega_{\omega, \delta}^-)} \right)$, it follows that the pair (v^+, v^-) defined by

$$v^\pm := \hat{v}^\pm + \tilde{v}^\pm - u^\pm \quad \text{in } [0, T] \times \overline{\Phi(\Omega_{\omega, \delta}^\pm)},$$

belongs to $C_0^{\frac{1+\alpha}{2}; 1+\alpha} \left([0, T] \times \overline{\Phi(\Omega_{\omega, \delta}^+)} \right) \times C_0^{\frac{1+\alpha}{2}; 1+\alpha} \left([0, T] \times \overline{\Phi(\Omega_{\omega, \delta}^-)} \right)$ and solves problem (7). Thus, in order to show the existence of a solution of problem (6), it remains to show the existence of a solution (\hat{v}^+, \hat{v}^-) of problem (8). It is classical that there exists a solution $\hat{v} \in C_0^{\frac{1+\alpha}{2}; 1+\alpha} \left([0, T] \times \overline{\Phi(\Omega_{\omega, \delta})} \right)$ of the Dirichlet problem

$$\begin{cases} \partial_t \hat{v} - \Delta \hat{v} = 0 & \text{in }]0, T[\times \Phi(\Omega_{\omega, \delta}), \\ \hat{v} = u^+ & \text{on } [0, T] \times \Phi(\partial\Omega_{\omega, \delta}^+ \setminus \partial\Omega), \\ \hat{v} = u^- & \text{on } [0, T] \times \Phi(\partial\Omega_{\omega, \delta}^- \setminus \partial\Omega), \\ \hat{v}(0, \cdot) = 0 & \text{in } \overline{\Phi(\Omega_{\omega, \delta})}. \end{cases}$$

A proof of this fact can be found for example in Baderko [1, Thm 5.1], Lunardi and Vespri [32, Thm. 4.3], and Lieberman [31, Thm. 6.48]. Then, if we set

$$\hat{v}^\pm := \hat{v} \Big|_{[0, T] \times \overline{\Phi(\Omega_{\omega, \delta}^\pm)}},$$

we have that the pair (\hat{v}^+, \hat{v}^-) belongs to $C_0^{\frac{1+\alpha}{2}; 1+\alpha} \left([0, T] \times \overline{\Phi(\Omega_{\omega, \delta}^+)} \right) \times C_0^{\frac{1+\alpha}{2}; 1+\alpha} \left([0, T] \times \overline{\Phi(\Omega_{\omega, \delta}^-)} \right)$ and solves problem (8). Thus the existence of a solution of (6) is proved.

Now we turn to the proof of the uniqueness of the solution of (6). Since we are dealing with a linear problem, it suffices to show that, whenever we take homogeneous data in the right-hand sides, the only solution is $(v^+, v^-) = (0, 0)$. So let $(f_0^+, f_0^-, f_1^+, f_1^-, g, g_1, h^+, h^-) = (0, 0, 0, 0, 0, 0, 0, 0)$ and let

$$(v^+, v^-) \in C_0^{\frac{1+\alpha}{2}; 1+\alpha} \left([0, T] \times \overline{\Phi(\Omega_{\omega, \delta}^+)} \right) \times C_0^{\frac{1+\alpha}{2}; 1+\alpha} \left([0, T] \times \overline{\Phi(\Omega_{\omega, \delta}^-)} \right)$$

be a solution of problem (6). Let

$$e^\pm(t) := \int_{\Phi(\Omega_{\omega, \delta}^\pm)} (v^\pm(t, y))^2 dy \quad \forall t \in [0, T].$$

Since v^\pm are uniformly continuous in $[0, T] \times \overline{\Phi(\Omega_{\omega, \delta}^\pm)}$, we can see that $e^\pm \in C^0([0, T])$. But the functions e^\pm are actually more regular than that. Indeed, an argument based on classical differentiation theorems for integrals depending on a parameter and on a specific

approximation of the support of integration (see Verchota [47, Thm. 1.12, p. 581]) shows that $e^\pm \in C^1(]0, T[)$. A detailed proof of this fact can be found in [33, Lemma 5 and Prop. 2]. Following the argument in the same reference we can prove that

$$\frac{d}{dt}e^+(t) = -2 \int_{\Phi(\Omega_{\omega,\delta}^+)} |Dv^+(t, y)|^2 dy + 2 \int_{\partial\Phi(\Omega_{\omega,\delta}^+)} v^+(t, y) \frac{\partial}{\partial\nu_{\Phi(\Omega_{\omega,\delta}^+)}(y)} v^+(t, y) d\sigma_y.$$

Then we have

$$\begin{aligned} \frac{d}{dt}e^+(t) &= -2 \int_{\Phi(\Omega_{\omega,\delta}^+)} |Dv^+(t, y)|^2 dy + 2 \int_{\Phi(\partial\Omega_{\omega,\delta}^+ \setminus \partial\Omega)} v^+(t, y) \frac{\partial}{\partial\nu_{\Phi(\Omega_{\omega,\delta}^+)}(y)} v^+(t, y) d\sigma_y \\ &\quad + 2 \int_{\Phi(\partial\Omega)} v^+(t, y) \frac{\partial}{\partial\nu_{\Phi(\Omega_{\omega,\delta}^+)}(y)} v^+(t, y) d\sigma_y \\ &= -2 \int_{\Phi(\Omega_{\omega,\delta}^+)} |Dv^+(t, y)|^2 dy + 2 \int_{\Phi(\partial\Omega)} v^+(t, y) \frac{\partial}{\partial\nu_{\Phi(\partial\Omega)}(y)} v^+(t, y) d\sigma_y, \end{aligned}$$

for all $t \in]0, T[$. Indeed the boundary condition on $\Phi(\partial\Omega_{\omega,\delta}^+ \setminus \partial\Omega)$ implies that

$$\int_{\Phi(\partial\Omega_{\omega,\delta}^+ \setminus \partial\Omega)} v^+(t, y) \frac{\partial}{\partial\nu_{\Phi(\Omega_{\omega,\delta}^+)}(y)} v^+(t, y) d\sigma_y = 0 \quad \forall t \in]0, T[.$$

Similarly

$$\begin{aligned} \frac{d}{dt}e^-(t) &= -2 \int_{\Phi(\Omega_{\omega,\delta}^-)} |Dv^-(t, y)|^2 dy + 2 \int_{\Phi(\partial\Omega_{\omega,\delta}^-)} v^-(t, y) \frac{\partial}{\partial\nu_{\Phi(\Omega_{\omega,\delta}^-)}(y)} v^-(t, y) d\sigma_y \\ &= -2 \int_{\Phi(\Omega_{\omega,\delta}^-)} |Dv^-(t, y)|^2 dy + 2 \int_{\Phi(\partial\Omega_{\omega,\delta}^- \setminus \partial\Omega)} v^-(t, y) \frac{\partial}{\partial\nu_{\Phi(\Omega_{\omega,\delta}^-)}(y)} v^-(t, y) d\sigma_y \\ &\quad + 2 \int_{\Phi(\partial\Omega)} v^-(t, y) \frac{\partial}{\partial\nu_{\Phi(\Omega_{\omega,\delta}^-)}(y)} v^-(t, y) d\sigma_y \\ &= -2 \int_{\Phi(\Omega_{\omega,\delta}^-)} |Dv^-(t, y)|^2 dy - 2 \int_{\Phi(\partial\Omega)} v^-(t, y) \frac{\partial}{\partial\nu_{\Phi(\partial\Omega)}(y)} v^-(t, y) d\sigma_y, \end{aligned}$$

for all $t \in]0, T[$. Then, if we set $e := e^+ + e^-$ in $[0, T]$, by the transmission conditions we get

$$\begin{aligned} \frac{d}{dt}e(t) &= -2 \int_{\Phi(\Omega_{\omega,\delta}^+)} |Dv^+(t, y)|^2 dy - 2 \int_{\Phi(\Omega_{\omega,\delta}^-)} |Dv^-(t, y)|^2 dy \\ &\quad + 2 \int_{\Phi(\partial\Omega)} v^+(t, y) \frac{\partial}{\partial\nu_{\Phi(\partial\Omega)}(y)} v^+(t, y) d\sigma_y - 2 \int_{\Phi(\partial\Omega)} v^-(t, y) \frac{\partial}{\partial\nu_{\Phi(\partial\Omega)}(y)} v^-(t, y) d\sigma_y \\ &= -2 \left(\int_{\Phi(\Omega_{\omega,\delta}^+)} |Dv^+(t, y)|^2 dy + \int_{\Phi(\Omega_{\omega,\delta}^-)} |Dv^-(t, y)|^2 dy \right), \end{aligned}$$

for all $t \in]0, T[$. Hence $\frac{d}{dt}e \leq 0$ in $]0, T[$. Since $e \geq 0$ and $e(0) = 0$, it follows that $e(t) = 0$ for all $t \in [0, T]$. Then $v^+ = 0$ and $v^- = 0$. \square

We can now characterize the layer potentials as specific solutions of the transmission problem (6). Indeed, by the previous theorem and by Theorem 2.6, we can see that the pair

$$(v^+[\mu]_{|[0, T] \times \overline{\Phi(\Omega_{\omega,\delta}^+)}, v^-[\mu]_{|[0, T] \times \overline{\Phi(\Omega_{\omega,\delta}^-)}}) \quad (9)$$

is the unique solution of problem (6) with data

$$\begin{aligned} & (f_0^+, f_0^-, f_1^+, f_1^-, g, g_1, h^+, h^-) \\ & = (0, 0, 0, 0, 0, \mu, v^+[\mu]_{|[0,T] \times \Phi(\partial\Omega_{\omega,\delta}^+ \setminus \partial\Omega)}, v^-[\mu]_{|[0,T] \times \Phi(\partial\Omega_{\omega,\delta}^- \setminus \partial\Omega)}). \end{aligned}$$

Hence, problem (6) with such data uniquely identifies the pair (9). Similarly, the pair

$$(w^+[\mu]_{|[0,T] \times \overline{\Phi(\Omega_{\omega,\delta}^+)}}), w^-[\mu]_{|[0,T] \times \overline{\Phi(\Omega_{\omega,\delta}^-)}})$$

is the unique solution of problem (6) with data

$$\begin{aligned} & (f_0^+, f_0^-, f_1^+, f_1^-, g, g_1, h^+, h^-) \\ & = (0, 0, 0, 0, -\mu, 0, w^+[\mu]_{|[0,T] \times \Phi(\partial\Omega_{\omega,\delta}^+ \setminus \partial\Omega)}, w^-[\mu]_{|[0,T] \times \Phi(\partial\Omega_{\omega,\delta}^+ \setminus \partial\Omega)}). \end{aligned}$$

However, to proceed with our analysis we need to characterize the the pulled-back pairs

$$(v^+[\mu] \circ \Phi^T, v^-[\mu] \circ \Phi^T)$$

and

$$(w^+[\mu] \circ \Phi^T, w^-[\mu] \circ \Phi^T)$$

rather than (v^+, v^-) and (w^+, w^-) . To do so, we will Φ -pullback problem (6) and obtain a new problem defined on the fixed domain $\Omega_{\omega,\delta}$. With the right set of data the new problem will have the pulled-back pairs as unique solutions.

4 Pulling-back the problem

Now our aim is to pull-back problem (6) to have a problem on the fixed domain $\Omega_{\omega,\delta}$. It appears, however, that in this process we cannot keep the strong formulation of the heat operator $\partial_t - \Delta$ that we have in the first and second equations of (6). This is because the Laplace operator Δ is a second order operator and to pull-back it we should take two derivatives of Φ . We should then assume Φ at least of class C^2 , but in our paper Φ is in $C^{1,\alpha}(\overline{\Omega_{\omega,\delta}}, \mathbb{R}^n)$. Not even the weak formulation of the heat operator would suffice. Indeed, that would yeald to pulled-back operator

$$\partial_t u - \operatorname{div} \left((D\Phi)^{-1} (D\Phi)^{-\top} (Du)^\top \right) - \frac{1}{|\det D\Phi|} D(|\det D\Phi|) (D\Phi)^{-1} (D\Phi)^{-\top} (Du)^\top$$

(see Chapko, Kress and Yoon [5, Eq. (2.10), p. 858]) and for the term $D(|\det D\Phi|)$ to be a function we still need Φ of class C^2 .

Then we need to adopt a different approach. In particular, we will place the problem in a specific quotient space. This space is introduced in the following lemma, which can be verified immediately.

Lemma 4.1. *Let $\alpha \in]0, 1[$ and $T \in]0, +\infty[$. Let Ω be a bounded open connected subset of \mathbb{R}^n of class $C^{1,\alpha}$. Then the set*

$$\begin{aligned} \mathcal{Y}_\Omega := & \left\{ w = (w_0, w_1) \in C_0^{\frac{1+\alpha}{2};\alpha}([0, T] \times \overline{\Omega}) \times C^{\frac{\alpha}{2};\alpha}([0, T] \times \overline{\Omega}, \mathbb{R}^n) \right. \\ & \left. : \int_0^T \int_\Omega w_0 \partial_t \varphi + (D\varphi) w_1 \, dx dt = 0 \quad \forall \varphi \in \mathcal{D}([0, T] \times \Omega) \right\} \end{aligned}$$

is a closed linear subspace of $C_0^{\frac{1+\alpha}{2};\alpha}([0, T] \times \overline{\Omega}) \times C^{\frac{\alpha}{2};\alpha}([0, T] \times \overline{\Omega}, \mathbb{R}^n)$ and the quotient space

$$\mathcal{X}_\Omega := C_0^{\frac{1+\alpha}{2};\alpha}([0, T] \times \overline{\Omega}) \times C^{\frac{\alpha}{2};\alpha}([0, T] \times \overline{\Omega}, \mathbb{R}^n) / \mathcal{Y}_\Omega,$$

equipped with the quotient norm, is a Banach space. We denote by Π the canonical projection from $C_0^{\frac{1+\alpha}{2};\alpha}([0, T] \times \bar{\Omega}) \times C^{\frac{\alpha}{2};\alpha}([0, T] \times \bar{\Omega}, \mathbb{R}^n)$ to \mathcal{X}_Ω .

In the following Lemma 4.2 we see how we can exploit the projection Π to transform a heat equation defined on a Φ -dependent set $[0, T] \times \Phi(\Omega)$ into an equation on the fixed set $[0, T] \times \Omega$. To do so, we need some more notation. If Ω is a bounded open connected subset of \mathbb{R}^n of class $C^{1,\alpha}$, $T \in]0, +\infty[$, and $\alpha \in]0, 1[$, then

$$B_\Omega : (C^{1,\alpha}(\bar{\Omega}, \mathbb{R}^n) \cap \mathcal{A}_{\bar{\Omega}}) \times C_0^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times \bar{\Omega}) \rightarrow C_0^{\frac{1+\alpha}{2};\alpha}([0, T] \times \bar{\Omega}) \times C^{\frac{\alpha}{2};\alpha}([0, T] \times \bar{\Omega}, \mathbb{R}^n)$$

is the map that takes a pair (Φ, u) to

$$B_\Omega[\Phi, u] := \left(-|\det D\Phi| u, (D\Phi)^{-1}(D\Phi)^{-\top}(Du)^\top |\det D\Phi| \right)$$

and

$$A_\Omega : (C^{1,\alpha}(\bar{\Omega}, \mathbb{R}^n) \cap \mathcal{A}_{\bar{\Omega}}) \times C_0^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times \bar{\Omega}) \rightarrow \mathcal{X}_\Omega$$

is the map that takes a pair (Φ, u) to

$$A_\Omega[\Phi, u] := \Pi B_\Omega[\Phi, u].$$

Lemma 4.2. *Let $\alpha \in]0, 1[$ and $T \in]0, +\infty[$. Let Ω be a bounded open connected subset of \mathbb{R}^n of class $C^{1,\alpha}$. Let $(\tilde{f}_0, \tilde{f}_1) \in C_0^{\frac{1+\alpha}{2};\alpha}([0, T] \times \bar{\Omega}) \times C^{\frac{\alpha}{2};\alpha}([0, T] \times \bar{\Omega}, \mathbb{R}^n)$ and $(\Phi, u) \in (C^{1,\alpha}(\bar{\Omega}, \mathbb{R}^n) \cap \mathcal{A}_{\bar{\Omega}}) \times C_0^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times \bar{\Omega})$. Then equality*

$$A_\Omega[\Phi, u] = \Pi(\tilde{f}_0, \tilde{f}_1) \tag{10}$$

holds true if and only if

$$\begin{aligned} & \partial_t \left(u \circ (\Phi^T)^{(-1)} \right) - \Delta \left(u \circ (\Phi^T)^{(-1)} \right) \\ &= \partial_t \left\{ \tilde{f}_0 \circ (\Phi^T)^{(-1)} \left| \det D(\Phi^{(-1)}) \right| \right\} + \operatorname{div} \left\{ ((D\Phi)\tilde{f}_1) \circ (\Phi^T)^{(-1)} \left| \det D(\Phi^{(-1)}) \right| \right\} \end{aligned} \tag{11}$$

in the sense of distributions in $]0, T[\times \Phi(\Omega)$.

Proof. By the definition of A_Ω and B_Ω , equation (10) is equivalent to

$$\Pi \left(-|\det D\Phi| u, (D\Phi)^{-1}(D\Phi)^{-\top}(Du)^\top |\det D\Phi| \right) = \Pi(\tilde{f}_0, \tilde{f}_1),$$

which in turn we can rewrite as

$$\begin{aligned} & \int_0^T \int_\Omega -|\det D\Phi| u \partial_t \varphi + (D\varphi)(D\Phi)^{-1}(D\Phi)^{-\top}(Du)^\top |\det D\Phi| \, dxdt \\ &= \int_0^T \int_\Omega \tilde{f}_0 \partial_t \varphi + (D\varphi)\tilde{f}_1 \, dxdt \quad \forall \varphi \in \mathcal{D}([0, T[\times \Omega). \end{aligned}$$

Then, by a change of variables in the integrals, we obtain

$$\begin{aligned} & \int_0^T \int_{\Phi(\Omega)} -(u \circ (\Phi^T)^{(-1)}) \partial_t (\varphi \circ (\Phi^T)^{(-1)}) + (D(\varphi \circ (\Phi^T)^{(-1)}))(D(u \circ (\Phi^T)^{(-1)}))^\top \, dxdt \\ &= \int_0^T \int_{\Phi(\Omega)} (\tilde{f}_0 \circ (\Phi^T)^{(-1)}) \left| \det D(\Phi^{(-1)}) \right| \partial_t (\varphi \circ (\Phi^T)^{(-1)}) \\ & \quad + (D(\varphi \circ (\Phi^T)^{(-1)}))((D\Phi)\tilde{f}_1) \circ (\Phi^T)^{(-1)} \left| \det D(\Phi^{(-1)}) \right| \, dxdt \end{aligned} \tag{12}$$

for all $\varphi \in \mathcal{D}([0, T[\times \Omega)$. Now, if $\psi \in \mathcal{D}([0, T[\times \Phi(\Omega))$ we take a sequence $\{\varphi_j\}_j$ in $\mathcal{D}([0, T[\times \Omega)$ that converges to $\psi \circ \Phi^T$ in $C^1([0, T[\times \overline{\Omega})$ and we see that (12) implies that

$$\begin{aligned} & \int_0^T \int_{\Phi(\Omega)} -(u \circ (\Phi^T)^{(-1)}) \partial_t \psi + (D\psi)(D(u \circ (\Phi^T)^{(-1)}))^\top dx dt \\ &= \int_0^T \int_{\Phi(\Omega)} (\tilde{f}_0 \circ (\Phi^T)^{(-1)}) |\det D(\Phi^{(-1)})| \partial_t \psi \\ & \quad + (D\psi)((D\Phi)\tilde{f}_1) \circ (\Phi^T)^{(-1)} |\det D(\Phi^{(-1)})| dx dt. \end{aligned}$$

Thus, equation (11) holds in the sense of distributions in $]0, T[\times \Phi(\Omega)$.

We have shown that (10) implies (11). The proof that (11) implies (10) is similar: we have to follow the above argument backward. \square

So much for the pull-back of the heat equation on a set $\Phi(\Omega)$, we can now go back to the transmission problem (6), which is defined on the Φ -dependent pair of sets $(\Phi(\Omega_{\omega, \delta}^+), \Phi(\Omega_{\omega, \delta}^-))$, and transform it into a problem on $(\Omega_{\omega, \delta}^+, \Omega_{\omega, \delta}^-)$. For the first two equations we can use Lemma 4.2, so it remains to see what to do with the other four equations. To shorten our notation, we set

$$\begin{aligned} \mathcal{Z} &:= \mathcal{X}_{\Omega_{\omega, \delta}^+} \times \mathcal{X}_{\Omega_{\omega, \delta}^-} \times C_0^{\frac{1+\alpha}{2}; 1+\alpha}([0, T] \times \partial\Omega) \times C_0^{\frac{\alpha}{2}; \alpha}([0, T] \times \partial\Omega) \\ & \quad \times C_0^{\frac{1+\alpha}{2}; 1+\alpha}([0, T] \times (\partial\Omega_{\omega, \delta}^+ \setminus \partial\Omega)) \times C_0^{\frac{1+\alpha}{2}; 1+\alpha}([0, T] \times (\partial\Omega_{\omega, \delta}^- \setminus \partial\Omega)), \end{aligned}$$

(cf. (4.1) for the definition of the quotient space \mathcal{X}_Ω). Then we have the following.

Theorem 4.3. *Let $T \in]0, +\infty[$, $\alpha \in]0, 1[$. Let Ω , ω and δ_Ω be as in Lemma 2.2. Let $\delta \in]0, \delta_\Omega[$. For $\Phi \in C^{1, \alpha}(\overline{\Omega_{\omega, \delta}}, \mathbb{R}^n) \cap \mathcal{A}'_{\Omega_{\omega, \delta}}$, let R_Φ denote the map from*

$$C_0^{\frac{1+\alpha}{2}; 1+\alpha}([0, T] \times \overline{\Omega_{\omega, \delta}^+}) \times C_0^{\frac{1+\alpha}{2}; 1+\alpha}([0, T] \times \overline{\Omega_{\omega, \delta}^-})$$

to \mathcal{Z} defined by

$$\begin{aligned} R_\Phi[U^+, U^-] &:= \left(A_{\Omega_{\omega, \delta}^+}[\Phi, U^+], A_{\Omega_{\omega, \delta}^-}[\Phi, U^-], U_{|[0, T] \times \partial\Omega}^+ - U_{|[0, T] \times \partial\Omega}^-, \right. \\ & \quad \left. J_\Phi[U^+, U^-], U_{|[0, T] \times (\partial\Omega_{\omega, \delta}^+ \setminus \partial\Omega)}^+, U_{|[0, T] \times (\partial\Omega_{\omega, \delta}^- \setminus \partial\Omega)}^- \right), \end{aligned} \quad (13)$$

where

$$J_\Phi[U^+, U^-] := DU^+(D\Phi)^{-1} \mathbf{n}[\Phi] - DU^-(D\Phi)^{-1} \mathbf{n}[\Phi] \quad \text{on } [0, T] \times \partial\Omega,$$

and

$$\mathbf{n}[\Phi] := \frac{(D\Phi(x))^{-\top} \nu_\Omega(x)}{|(D\Phi(x))^{-\top} \nu_\Omega(x)|} \quad \forall x \in \partial\Omega.$$

Then the following statements hold.

(i) *Let $\Phi \in C^{1, \alpha}(\overline{\Omega_{\omega, \delta}}, \mathbb{R}^n) \cap \mathcal{A}'_{\Omega_{\omega, \delta}}$. Let $(F^+, F^-, G, G_1, H^+, H^-) \in \mathcal{Z}$. Then $(U^+, U^-) \in C_0^{\frac{1+\alpha}{2}; 1+\alpha}([0, T] \times \overline{\Omega_{\omega, \delta}^+}) \times C_0^{\frac{1+\alpha}{2}; 1+\alpha}([0, T] \times \overline{\Omega_{\omega, \delta}^-})$ is a solution of the equation*

$$R_\Phi[U^+, U^-] = (F^+, F^-, G, G_1, H^+, H^-) \quad (14)$$

if and only if the pair

$$(U^+ \circ (\Phi^T)^{(-1)}, U^- \circ (\Phi^T)^{(-1)}) \in C_0^{\frac{1+\alpha}{2}; 1+\alpha}([0, T] \times \overline{\Phi(\Omega_{\omega, \delta}^+)}) \times C_0^{\frac{1+\alpha}{2}; 1+\alpha}([0, T] \times \overline{\Phi(\Omega_{\omega, \delta}^-)})$$

is a solution of problem (6) with data

$$\begin{aligned} f_0^\pm &:= \tilde{f}_0^\pm \circ (\Phi^T)^{(-1)} |\det D(\Phi^{(-1)})|, \\ f_1^\pm &:= ((D\Phi)\tilde{f}_1^\pm) \circ (\Phi^T)^{(-1)} |\det D(\Phi^{(-1)})|, \\ g &:= G \circ (\Phi^T)^{(-1)}, \quad g_1 := G_1 \circ (\Phi^T)^{(-1)}, \quad h^\pm := H^\pm \circ (\Phi^T)^{(-1)}, \end{aligned} \quad (15)$$

and with $(\tilde{f}_0^\pm, \tilde{f}_1^\pm) \in C_0^{\frac{1+\alpha}{2};\alpha}([0, T] \times \overline{\Omega_{\omega,\delta}^\pm}) \times C^{\frac{\alpha}{2};\alpha}([0, T] \times \overline{\Omega_{\omega,\delta}^\pm}, \mathbb{R}^n)$ such that $\Pi(\tilde{f}_0^\pm, \tilde{f}_1^\pm) = F^\pm$.

(ii) Let $\Phi \in C^{1,\alpha}(\overline{\Omega_{\omega,\delta}}, \mathbb{R}^n) \cap \mathcal{A}'_{\Omega_{\omega,\delta}}$. Then R_Φ is a linear homeomorphism from

$$C_0^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times \overline{\Omega_{\omega,\delta}^+}) \times C_0^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times \overline{\Omega_{\omega,\delta}^-})$$

onto \mathcal{Z} .

Proof. We first consider statement (i). Let $(U^+, U^-) \in C_0^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times \overline{\Omega_{\omega,\delta}^+}) \times C_0^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times \overline{\Omega_{\omega,\delta}^-})$ satisfy equation (14). Lemmas 4.1 and 4.2 imply that there exist $(\tilde{f}_0^\pm, \tilde{f}_1^\pm) \in C_0^{\frac{1+\alpha}{2};\alpha}([0, T] \times \overline{\Omega_{\omega,\delta}^\pm}) \times C^{\frac{\alpha}{2};\alpha}([0, T] \times \overline{\Omega_{\omega,\delta}^\pm}, \mathbb{R}^n)$ such that $\Pi(\tilde{f}_0^\pm, \tilde{f}_1^\pm) = F^\pm$ and such that

$$\begin{aligned} (\partial_t - \Delta)(U^\pm \circ (\Phi^T)^{(-1)}) &= \operatorname{div} \left\{ ((D\Phi)\tilde{f}_1^\pm) \circ (\Phi^T)^{(-1)} |\det D(\Phi^{(-1)})| \right\} \\ &\quad + \partial_t \left\{ \tilde{f}_0^\pm \circ (\Phi^T)^{(-1)} |\det D(\Phi^{(-1)})| \right\} \quad \text{in }]0, T[\times \Phi(\Omega_{\omega,\delta}^\pm). \end{aligned}$$

Clearly we have that

$$U^+ \circ (\Phi^T)^{(-1)} - U^- \circ (\Phi^T)^{(-1)} = G \circ (\Phi^T)^{(-1)} = g \quad \text{on } [0, T] \times \Phi(\partial\Omega),$$

and

$$U^\pm \circ (\Phi^T)^{(-1)} = H^\pm \circ (\Phi^T)^{(-1)} = h^\pm \quad \text{on } [0, T] \times \Phi(\partial\Omega_{\omega,\delta}^\pm \setminus \partial\Omega).$$

Since

$$\nu_{\Phi|\partial\Omega} \circ \Phi|_{\partial\Omega} = \frac{(D\Phi)^{-\top} \nu_\Omega}{|(D\Phi)^{-\top} \nu_\Omega|} = \mathbf{n}[\Phi] \quad \text{on } \partial\Omega$$

(see, e.g., Lanza de Cristoforis and Rossi [30, Lem. 4.2, p. 207]), we have that

$$\begin{aligned} (DU^\pm) \circ (\Phi^T)^{(-1)} (D\Phi)^{-1} \circ \Phi^{(-1)} \mathbf{n}[\Phi] \circ \Phi^{(-1)} \\ = (DU^\pm) \circ (\Phi^T)^{(-1)} (D\Phi)^{-1} \nu_{\Phi|\partial\Omega} \\ = \frac{\partial}{\partial \nu_{\Phi|\partial\Omega}} (U^\pm \circ (\Phi^T)^{(-1)}) \quad \text{on } [0, T] \times \Phi(\partial\Omega). \end{aligned}$$

Accordingly, equality

$$DU^+(D\Phi)^{-1} \mathbf{n}[\Phi] - DU^-(D\Phi)^{-1} \mathbf{n}[\Phi] = G_1 \quad \text{on } [0, T] \times \partial\Omega,$$

implies that

$$\frac{\partial}{\partial \nu_{\Phi|\partial\Omega}} (U^+ \circ (\Phi^T)^{(-1)}) - \frac{\partial}{\partial \nu_{\Phi|\partial\Omega}} (U^- \circ (\Phi^T)^{(-1)}) = G_1 \circ (\Phi^T)^{(-1)} = g_1 \quad \text{on } [0, T] \times \Phi(\partial\Omega).$$

Thus, the pair $(U^+ \circ (\Phi^T)^{(-1)}, U^- \circ (\Phi^T)^{(-1)})$ solves the transmission problem (6) with data as in (15). The converse can be proved by reading backward the above argument.

We now prove statement (ii). We can readily see that R_Φ is linear and continuous. So, it suffices to show that R_Φ is bijective to deduce from the open mapping theorem that it is a homeomorphism. We start by proving that it is injective. Let $(U^+, U^-) \in C_0^{\frac{1+\alpha}{2}; 1+\alpha} \left([0, T] \times \overline{\Omega_{\omega, \delta}^+} \right) \times C_0^{\frac{1+\alpha}{2}; 1+\alpha} \left([0, T] \times \overline{\Omega_{\omega, \delta}^-} \right)$ and suppose that

$$R_\Phi[U^+, U^-] = (0, 0, 0, 0, 0, 0).$$

By statement (i), the pair

$$\left(U^+ \circ (\Phi^T)^{(-1)}, U^- \circ (\Phi^T)^{(-1)} \right)$$

solves problem (6) with data $(f_0^+, f_0^-, f_1^+, f_1^-, g, g_1, h^+, h^-) = (0, 0, 0, 0, 0, 0, 0, 0)$. Then, Theorem 3.1 implies that $(U^+ \circ (\Phi^T)^{(-1)}, U^- \circ (\Phi^T)^{(-1)}) = (0, 0)$ and accordingly we have $(U^+, U^-) = (0, 0)$. Now it remains to show that R_Φ is surjective. Let

$$(F^+, F^-, G, G_1, H^+, H^-) \in \mathcal{Z}.$$

It is easy to check that $(f_0^+, f_0^-, f_1^+, f_1^-, g, g_1, h^+, h^-)$ defined as in (15) belongs to \mathcal{S}_Φ (cf. (5)). Accordingly, Theorem 3.1 implies that there exists a pair

$$(v^+, v^-) \in C_0^{\frac{1+\alpha}{2}; 1+\alpha} \left([0, T] \times \overline{\Phi(\Omega_{\omega, \delta}^+)} \right) \times C_0^{\frac{1+\alpha}{2}; 1+\alpha} \left([0, T] \times \overline{\Phi(\Omega_{\omega, \delta}^-)} \right)$$

that solves the corresponding problem (6). Then statement (i) implies that

$$(U^+, U^-) := (v^+ \circ \Phi^T, v^- \circ \Phi^T) \in C_0^{\frac{1+\alpha}{2}; 1+\alpha} \left([0, T] \times \overline{\Omega_{\omega, \delta}^+} \right) \times C_0^{\frac{1+\alpha}{2}; 1+\alpha} \left([0, T] \times \overline{\Omega_{\omega, \delta}^-} \right)$$

is a solution of

$$R_\Phi[U^+, U^-] = (F^+, F^-, G, G_1, H^+, H^-).$$

□

We are now ready to characterize the pairs

$$(V_\Phi^+[\mu], V_\Phi^-[\mu]) := \left(v^+[\mu \circ (\Phi^T)^{(-1)}] \circ \Phi^T, v^-[\mu \circ (\Phi^T)^{(-1)}] \circ \Phi^T \right)$$

and

$$(W_\Phi^+[\mu], W_\Phi^-[\mu]) := \left(w^+[\mu \circ (\Phi^T)^{(-1)}] \circ \Phi^T, w^-[\mu \circ (\Phi^T)^{(-1)}] \circ \Phi^T \right)$$

as solutions of Φ -dependent equations defined on a fixed domain. This will be done in the following Theorem 4.4. The proof is a straightforward consequence of Theorem 4.3, of Theorems 2.5 and 2.6 on the properties of the layer heat potentials, and of Lemma 2.4 (i) on the change of variable in integrals over $\Phi(\partial\Omega)$.

Theorem 4.4. *Let $T \in]0, +\infty[$, $\alpha \in]0, 1[$. Let Ω , ω and δ_Ω be as in Lemma 2.2. Let $\delta \in]0, \delta_\Omega[$. Let $\Phi \in C^{1, \alpha}(\overline{\Omega_{\omega, \delta}}, \mathbb{R}^n) \cap \mathcal{A}'_{\Omega_{\omega, \delta}}$. Then the following statements hold.*

(i) *Let $\mu \in C_0^{\frac{\alpha}{2}; \alpha}([0, T] \times \partial\Omega)$. Then*

$$(V_\Phi^+[\mu], V_\Phi^-[\mu]) = R_\Phi^{(-1)}(0, 0, 0, \mu, \mathcal{V}_{\delta, \Phi}^+[\mu], \mathcal{V}_{\delta, \Phi}^-[\mu]) \quad (16)$$

with

$$\begin{aligned} \mathcal{V}_{\delta, \Phi}^\pm[\mu](t, x) &:= \int_0^t \int_{\partial\Omega} S_n(t - \tau, \Phi(x) - \Phi(y)) \mu(\tau, y) \tilde{\sigma}_n[\Phi|_{\partial\Omega}](y) d\sigma_y d\tau \\ &\quad \forall (t, x) \in [0, T] \times (\partial\Omega_{\omega, \delta}^\pm \setminus \partial\Omega). \end{aligned}$$

(ii) Let $\mu \in C_0^{\frac{1+\alpha}{2}; 1+\alpha}([0, T] \times \partial\Omega)$. Then

$$(W_{\Phi}^+[\mu], W_{\Phi}^-[\mu]) = R_{\Phi}^{(-1)}(0, 0, -\mu, 0, \mathcal{W}_{\delta, \Phi}^+[\mu], \mathcal{W}_{\delta, \Phi}^-[\mu]) \quad (17)$$

with

$$\begin{aligned} \mathcal{W}_{\delta, \Phi}^{\pm}[\mu] := & - \int_0^t \int_{\partial\Omega} DS_n(t - \tau, \Phi(x) - \Phi(y)) \nu_{\Phi|_{\partial\Omega}}(\Phi(y)) \mu(\tau, y) \tilde{\sigma}_n[\Phi|_{\partial\Omega}](y) d\sigma_y d\tau \\ & \forall (t, x) \in [0, T] \times (\partial\Omega_{\omega, \delta}^{\pm} \setminus \partial\Omega). \end{aligned}$$

5 Dependence of the heat layer potentials upon shape perturbations

In this section we prove our main result. That is, we prove that the maps V_{ϕ} , $V_{l, \phi}$, $W_{*, \phi}$, and W_{ϕ} in (1)–(4) depend smoothly on the shape parameter ϕ . To do so, we first use equalities (16) and (17) to show that the maps $\Phi \mapsto V_{\Phi}^{\pm}$ and $\Phi \mapsto W_{\Phi}^{\pm}$ are of class C^{∞} . So we have to understand the regularity of the terms appearing in these equations, and we begin with the map that takes Φ to $R_{\Phi}^{(-1)}$.

Proposition 5.1. *Let $T \in]0, +\infty[$, $\alpha \in]0, 1[$. Let Ω , ω and δ_{Ω} be as in Lemma 2.2. Let $\delta \in]0, \delta_{\Omega}[$. The map that takes $\Phi \in C^{1, \alpha}(\overline{\Omega_{\omega, \delta}}, \mathbb{R}^n) \cap \mathcal{A}'_{\overline{\Omega_{\omega, \delta}}}$ to*

$$R_{\Phi}^{(-1)} \in \mathcal{L} \left(\mathcal{Z}, C_0^{\frac{1+\alpha}{2}; 1+\alpha} \left([0, T] \times \overline{\Omega_{\omega, \delta}^+} \right) \times C_0^{\frac{1+\alpha}{2}; 1+\alpha} \left([0, T] \times \overline{\Omega_{\omega, \delta}^-} \right) \right)$$

is real analytic.

Proof. By the definition of R_{Φ} in (13), to prove that the map that takes $\Phi \in C^{1, \alpha}(\overline{\Omega_{\omega, \delta}}, \mathbb{R}^n) \cap \mathcal{A}'_{\overline{\Omega_{\omega, \delta}}}$ to

$$R_{\Phi} \in \mathcal{L} \left(C_0^{\frac{1+\alpha}{2}; 1+\alpha} \left([0, T] \times \overline{\Omega_{\omega, \delta}^+} \right) \times C_0^{\frac{1+\alpha}{2}; 1+\alpha} \left([0, T] \times \overline{\Omega_{\omega, \delta}^-} \right), \mathcal{Z} \right)$$

is real analytic it suffices to check that the maps that take Φ to

$$A_{\Omega_{\omega, \delta}^{\pm}}[\Phi, \cdot] \in \mathcal{L} \left(C_0^{\frac{1+\alpha}{2}; 1+\alpha} \left([0, T] \times \overline{\Omega_{\omega, \delta}^{\pm}} \right), \mathcal{X}_{\Omega_{\omega, \delta}^{\pm}} \right)$$

and to

$$J_{\Phi} \in \mathcal{L} \left(C_0^{\frac{1+\alpha}{2}; 1+\alpha} \left([0, T] \times \overline{\Omega_{\omega, \delta}^+} \right) \times C_0^{\frac{1+\alpha}{2}; 1+\alpha} \left([0, T] \times \overline{\Omega_{\omega, \delta}^-} \right), C_0^{\frac{\alpha}{2}; \alpha}([0, T] \times \partial\Omega) \right)$$

are real analytic. This follows from the real analyticity of the map that takes an invertible matrix with Schauder entries to its inverse (cf., e.g., [29, Lemma 2.1], see also [22]) and from the real analyticity of the map that takes Φ to $\mathbf{n}[\Phi] = \nu_{\Phi|_{\partial\Omega}} \circ \Phi|_{\partial\Omega}$ (cf. Lemma 2.4). Then, to complete the proof of the proposition it suffices to remember that the set of invertible operators is open, the map that takes an invertible operator to its inverse is real analytic, and the composition of real analytic maps is real analytic (see, e.g., Hille and Phillips [20, Theorems 4.3.2 and 4.3.4] and Prodi and Ambrosetti [44, Theorem 11.1]). \square

We now turn to the maps $\Phi \mapsto \mathcal{V}_{\delta, \Phi}^{\pm}$ and $\Phi \mapsto \mathcal{W}_{\delta, \Phi}^{\pm}$ of Theorem 4.4.

Proposition 5.2. *Let $T \in]0, +\infty[$, $\alpha \in]0, 1[$. Let Ω , ω and δ_Ω be as in Lemma 2.2. Let $\delta \in]0, \delta_\Omega[$. Then the maps that take $\Phi \in C^{1,\alpha}(\overline{\Omega_{\omega,\delta}}, \mathbb{R}^n) \cap \mathcal{A}'_{\Omega_{\omega,\delta}}$ to*

$$\mathcal{V}_{\delta,\Phi}^\pm \in \mathcal{L} \left(C_0^{\frac{\alpha}{2};\alpha}([0, T] \times \partial\Omega), C_0^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times (\partial\Omega_{\omega,\delta}^\pm \setminus \partial\Omega)) \right)$$

and to

$$\mathcal{W}_{\delta,\Phi}^\pm \in \mathcal{L} \left(C_0^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times \partial\Omega), C_0^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times (\partial\Omega_{\omega,\delta}^\pm \setminus \partial\Omega)) \right)$$

are of class C^∞ .

Proof. We verify the statement for $\mathcal{V}_{\delta,\Phi}^+$. The map from $C^{1,\alpha}(\overline{\Omega_{\omega,\delta}}, \mathbb{R}^n) \cap \mathcal{A}'_{\Omega_{\omega,\delta}}$ to $C^{1,\alpha}((\partial\Omega_{\omega,\delta}^+ \setminus \partial\Omega) \times \partial\Omega, \mathbb{R}^n \setminus \{0\})$ that takes Φ to the map Ψ defined by

$$\Psi(x, y) := \Phi(x) - \Phi(y) \quad \forall (x, y) \in (\partial\Omega_{\omega,\delta}^+ \setminus \partial\Omega) \times \partial\Omega,$$

is linear and continuous and therefore of class C^∞ . Since the composition of two C^∞ maps is of class C^∞ , Lemma A.3 of the Appendix on the regularity of a superposition operator implies that the map from $C^{1,\alpha}(\overline{\Omega_{\omega,\delta}}, \mathbb{R}^n) \cap \mathcal{A}'_{\Omega_{\omega,\delta}}$ to $C_0^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times ((\partial\Omega_{\omega,\delta}^+ \setminus \partial\Omega) \times \partial\Omega))$ that takes Φ to the function defined by

$$S_n(t, \Phi(x) - \Phi(y)) \quad \forall (t, x, y) \in [0, T] \times ((\partial\Omega_{\omega,\delta}^+ \setminus \partial\Omega) \times \partial\Omega)$$

is of class C^∞ . Finally, Lemma 2.4 on the real analyticity of $\tilde{\sigma}_n[\cdot]$ and Lemma A.2 of the Appendix on the linearity and continuity of a time dependent integral operator, imply the validity of the statement for $\mathcal{V}_{\delta,\Phi}^+$. The proof of the statement for $\mathcal{V}_{\delta,\Phi}^-$ and for $\mathcal{W}_{\delta,\Phi}^\pm$ are very similar and therefore omitted. \square

We observe that, in the proof of Proposition 5.2, it is the regularity of the fundamental solution S_n to prevent $\Phi \mapsto \mathcal{V}_{\delta,\Phi}^\pm$ and $\Phi \mapsto \mathcal{W}_{\delta,\Phi}^\pm$ from being real analytic. Indeed, for $\xi \neq 0$ the function $t \mapsto S_n(t, \xi)$ belongs to $C^\infty(\mathbb{R})$ but is not real analytic.

We can now go back to the maps $\Phi \mapsto \mathcal{V}_{\delta,\Phi}^\pm$ and $\Phi \mapsto \mathcal{W}_{\delta,\Phi}^\pm$ and prove that they are smooth.

Theorem 5.3. *Let $T \in]0, +\infty[$, $\alpha \in]0, 1[$. Let Ω , ω and δ_Ω be as in Lemma 2.2. Let $\delta \in]0, \delta_\Omega[$. Then the maps that take $\Phi \in C^{1,\alpha}(\overline{\Omega_{\omega,\delta}}, \mathbb{R}^n) \cap \mathcal{A}'_{\Omega_{\omega,\delta}}$ to*

$$\mathcal{V}_{\delta,\Phi}^\pm \in \mathcal{L} \left(C_0^{\frac{\alpha}{2};\alpha}([0, T] \times \partial\Omega), C_0^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times \overline{\Omega_{\omega,\delta}^\pm}) \right)$$

and to

$$\mathcal{W}_{\delta,\Phi}^\pm \in \mathcal{L} \left(C_0^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times \partial\Omega), C_0^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times \overline{\Omega_{\omega,\delta}^\pm}) \right)$$

are of class C^∞ .

Proof. It follows from equalities (16) and (17), from Propositions 5.1 and 5.2, and because the composition of a real analytic map with a C^∞ map is of class C^∞ . \square

As a corollary of Theorem 5.3, we are now ready to prove our main result.

Theorem 5.4. *Let $\alpha \in]0, 1[$, $T \in]0, +\infty[$. Let Ω be a bounded open subset of \mathbb{R}^n of class $C^{1,\alpha}$ such that both Ω and Ω^- are connected. Then the maps that take $\phi \in C^{1,\alpha}(\partial\Omega, \mathbb{R}^n) \cap \mathcal{A}_{\partial\Omega}$ to*

$$\begin{aligned} V_\phi &\in \mathcal{L} \left(C_0^{\frac{\alpha}{2};\alpha}([0, T] \times \partial\Omega), C_0^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times \partial\Omega) \right), \\ V_{l,\phi} &\in \mathcal{L} \left(C_0^{\frac{\alpha}{2};\alpha}([0, T] \times \partial\Omega), C_0^{\frac{\alpha}{2};\alpha}([0, T] \times \partial\Omega) \right) \quad \text{with } l \in \{1, \dots, n\}, \\ W_{*,\phi} &\in \mathcal{L} \left(C_0^{\frac{\alpha}{2};\alpha}([0, T] \times \partial\Omega), C_0^{\frac{\alpha}{2};\alpha}([0, T] \times \partial\Omega) \right), \\ W_\phi &\in \mathcal{L} \left(C_0^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times \partial\Omega), C_0^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times \partial\Omega) \right) \end{aligned}$$

are of class C^∞ .

Proof. It clearly suffices to show that the maps in the statement are of class C^∞ in a neighborhood of a function $\phi_0 \in C^{1,\alpha}(\partial\Omega, \mathbb{R}^n) \cap \mathcal{A}_{\partial\Omega}$. By the definitions (1)–(4), by the jump relations of the layer potentials of Theorems 2.5 and 2.6, and by the extension result of Lemma 2.3, there exists an open neighborhood \mathcal{W}_0 of ϕ_0 in $C^{1,\alpha}(\partial\Omega, \mathbb{R}^n) \cap \mathcal{A}_{\partial\Omega}$ such that

$$\begin{aligned} V_\phi[\mu] &= v^+[\mu \circ (\phi^T)^{(-1)}] \circ \phi^T = V_{\mathbf{E}[\phi]}^+[\mu], \\ V_{l,\phi}[\mu] &= -\frac{\mathbf{n}_l[\mathbf{E}[\phi]]}{2}\mu + \frac{\partial}{\partial x_l}(v^+[\mu \circ (\phi^T)^{(-1)}] \circ \phi^T) \\ &= -\frac{\mathbf{n}_l[\mathbf{E}[\phi]]}{2}\mu + ((DV_{\mathbf{E}[\phi]}^+[\mu]) \cdot (D\mathbf{E}[\phi])^{-1})_l, \\ W_{*,\phi}[\mu] &= -\frac{1}{2}\mu + ((Dv^+[\mu \circ (\phi^T)^{(-1)}] \circ \phi^T)\nu_\phi \circ \phi) \\ &= -\frac{1}{2}\mu + ((DV_{\mathbf{E}[\phi]}^+[\mu]) \cdot (D\mathbf{E}[\phi])^{-1}) \cdot \mathbf{n}[\mathbf{E}[\phi]] \end{aligned}$$

for all $\phi \in \mathcal{W}_0$ and all $\mu \in C_0^{\frac{\alpha}{2};\alpha}([0, T] \times \partial\Omega)$, and such that

$$W_\phi[\mu] = \frac{1}{2}\mu + W_{\mathbf{E}[\phi]}^+[\mu]$$

for all $\phi \in \mathcal{W}_0$ and all $\mu \in C_0^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times \partial\Omega)$. Thus, the statement follows by Theorem 5.3, by Lemma 2.3, and by standard calculus in Banach spaces. \square

Appendix A

In this appendix we collect a few auxiliary and technical results on the regularity of certain composition and nonlinear time-dependent integral operators. First of all, we introduce some notation and some definitions. Let $n, s \in \mathbb{N} \setminus \{0\}$, $\alpha \in]0, 1[$, $1 \leq s \leq n$. We set $\mathbb{B}_s := \{x \in \mathbb{R}^s : |x| < 1\}$. We say that a subset M of \mathbb{R}^n is a differential manifold (or simply a manifold) of dimension s and of class $C^{1,\alpha}$ imbedded in \mathbb{R}^n if, for every $P \in M$, there exist a neighborhood W of P in \mathbb{R}^n and a parametrization $\psi \in C^{1,\alpha}(\overline{\mathbb{B}_s}, \mathbb{R}^n)$ such that ψ is a homeomorphism of \mathbb{B}_s onto $W \cap M$, $\psi(0) = P$, and $D\psi$ has rank s at all points of $\overline{\mathbb{B}_s}$. If we further assume that M is compact, then there exist $P_1, \dots, P_r \in M$ and parametrizations $\{\psi_i\}_{i=1,\dots,r}$ with $\psi_i \in C^{1,\alpha}(\overline{\mathbb{B}_s}, \mathbb{R}^n)$ such that $\bigcup_{i=1}^r \psi_i(\overline{\mathbb{B}_s}) = M$.

Let M be a compact manifold of dimension s and of class $C^{1,\alpha}$ imbedded in \mathbb{R}^n . We can use the local parametrizations to define the Banach space $C^{1,\alpha}(M)$ (see, e.g., Lanza de Cristoforis and Rossi [29, p. 142]). Similarly, we can define the parabolic counterpart

of $C^{1,\alpha}(M)$. Let $T \in]0, +\infty[$. The space $C^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times M)$ is the space of functions f from $[0, T] \times M$ to \mathbb{R} such that

$$f \circ \psi_i^T \in C^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times \overline{\mathbb{B}_s}) \quad \forall i = 1, \dots, r,$$

where $\{\psi_i\}_{i=1,\dots,r}$ is a parametrization of M . On $C^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times M)$ we define a norm by setting

$$\|f\|_{C^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times M)} := \sup_{i=1,\dots,r} \|f \circ \psi_i^T\|_{C^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times \overline{\mathbb{B}_s})} \quad \forall f \in C^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times M).$$

We can verify that, with a different finite family of parametrizations of M , we obtain an equivalent norm. Moreover, $C^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times M)$ with the norm $\|\cdot\|_{C^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times M)}$ is a Banach space. Then

$$C_0^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times M) := \left\{ f \in C^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times M) : f(0, x) = 0 \quad \forall x \in M \right\},$$

is a Banach subspace of $C^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times M)$. The regularity of maps with values in $C^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times M)$ can be described using the local parametrization. More precisely, we have the following lemma, which can be proved by exploiting the definition of norm in the spaces $C^{\frac{1+\alpha}{2};1+\alpha}$.

Lemma A.1. *Let \mathcal{X} be a Banach space, and let \mathcal{O} be an open subset of \mathcal{X} . Let $\alpha \in]0, 1[$. Let $T \in]0, +\infty[$. Let M be a compact manifold of dimension $1 \leq s \leq n$ of class $C^{1,\alpha}$ imbedded in \mathbb{R}^n . Let N be a map from \mathcal{O} to $C^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times M)$. Let $\{\psi_i\}_{i=1,\dots,r}$ be a parametrization of M . For $i \in \{1, \dots, r\}$, let C_{ψ_i} be the composition operator from $C^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times M)$ to $C^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times \overline{\mathbb{B}_s})$ defined by*

$$C_{\psi_i}[f] := f \circ \psi_i^T \quad \forall f \in C^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times M).$$

Let $h \in \mathbb{N} \cup \{\infty\}$. Then N is of class C^h if and only if the operator $C_{\psi_i} \circ N$ is of class C^h for all $i = 1, \dots, r$.

Next, we turn to a time dependent integral operator with kernel in a parabolic Schauder space and summable density function.

Lemma A.2. *Let $n_1, n_2, s_1, s_2 \in \mathbb{N}$, $1 \leq s_1 \leq n_1$, $1 \leq s_2 \leq n_2$, $\alpha \in]0, 1[$. Let $T \in]0, +\infty[$. Let M_1, M_2 be two compact manifolds of dimension s_1, s_2 and of class $C^{1,\alpha}$ imbedded in $\mathbb{R}^{n_1}, \mathbb{R}^{n_2}$, respectively. Then the bilinear map K from $C_0^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times (M_1 \times M_2)) \times L^1([0, T] \times M_2)$ to $C_0^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times M_1)$ defined by*

$$K[G, f](t, x) := \int_0^t \int_{M_2} G(t - \tau, x, y) f(\tau, y) d\sigma_y d\tau \quad \forall (t, x) \in [0, T] \times M_1,$$

for all $(G, f) \in C_0^{\frac{1+\alpha}{2};1+\alpha}([0, T] \times (M_1 \times M_2)) \times L^1([0, T] \times M_2)$ is continuous.

Proof. Let $\{\phi_i\}_{i=1,\dots,r_1}$ and $\{\psi_j\}_{j=1,\dots,r_2}$ be local parametrizations of class $C^{1,\alpha}$ for M_1 and M_2 , respectively. We can suppose that $\bigcup_{i=1}^{r_1} \phi_i(\mathbb{B}_{s_1}/2) = M_1$ and $\bigcup_{j=1}^{r_2} \psi_j(\mathbb{B}_{s_2}/2) = M_2$. Let $\{\theta_j\}_{j=1,\dots,r_2}$ be a partition of unity subordinated to the parametrization $\{\psi_j\}_{j=1,\dots,r_2}$. Let π_1 and π_2 be the canonical projections of $\mathbb{R}^{n_1} \times \mathbb{R}^{n_2}$ onto \mathbb{R}^{n_1} and \mathbb{R}^{n_2} , respectively. Clearly $M_1 \times M_2$ is a compact manifold of dimension $s_1 + s_2$ and of class $C^{1,\alpha}$ imbedded in $\mathbb{R}^{n_1} \times \mathbb{R}^{n_2}$, and

$$\left\{ (\phi_i \circ \pi_1, \psi_j \circ \pi_2) \right\}_{\substack{i=1,\dots,r_1 \\ j=1,\dots,r_2}}$$

is a local parametrization of maps in $C^{1,\alpha}(\overline{\mathbb{B}_{s_1+s_2}}, \mathbb{R}^{n_1+n_2})$ for $M_1 \times M_2$. It suffices to show that there exists a constant $c > 0$ such that

$$\begin{aligned} & \sup_{i=1,\dots,r_1} \|K[G, f] \circ \phi_i^T\|_{C_0^{\frac{1+\alpha}{2}; 1+\alpha}([0, T] \times \overline{\mathbb{B}_{s_1}})} \\ & \leq c \sup_{\substack{i=1,\dots,r_1 \\ j=1,\dots,r_2}} \|G \circ (\phi_i \circ \pi_1, \psi_j \circ \pi_2)^T\|_{C_0^{\frac{1+\alpha}{2}; 1+\alpha}([0, T] \times \overline{\mathbb{B}_{s_1+s_2}})} \\ & \quad \times \sum_{j=1}^{r_2} \int_0^T \int_{\mathbb{B}_{s_2}} |f(\tau, \psi_j(\omega)) \theta_j(\psi_j(\omega))| |(D\psi_j^t \cdot D\psi_j)(\omega)|^{1/2} d\omega d\tau, \end{aligned}$$

for all $(G, f) \in C_0^{\frac{1+\alpha}{2}; 1+\alpha}([0, T] \times (M_1 \times M_2)) \times L^1([0, T] \times M_2)$. The above inequality follows by the equality

$$\begin{aligned} & K[G, f](t, \phi_i(\xi)) \\ & = \sum_{j=1}^{r_2} \int_0^t \int_{\mathbb{B}_{s_2}} G(t - \tau, \phi_i(\xi), \psi_j(\omega)) f(\tau, \psi_j(\omega)) \theta_j(\psi_j(\omega)) |(D\psi_j^t \cdot D\psi_j)(\omega)|^{1/2} d\omega d\tau \\ & \quad \forall (t, \xi) \in [0, T] \times \overline{\mathbb{B}_{s_1}}, \end{aligned}$$

that holds for all $(G, f) \in C_0^{\frac{1+\alpha}{2}; 1+\alpha}([0, T] \times (M_1 \times M_2)) \times L^1([0, T] \times M_2)$, by classical differentiation theorems for integrals depending on a parameter, and by the continuity of the linear map that takes a summable function to its integral. \square

The last result of this appendix shows a regularity result for a time dependent superposition operator.

Lemma A.3. *Let $n_1, n_2, s \in \mathbb{N}$, $1 \leq s \leq n_2$, $n_1 \geq 1$, $\alpha \in]0, 1[$. Let $T \in]0, +\infty[$. Let M be a compact manifold of dimension s and of class $C^{1,\alpha}$ imbedded in \mathbb{R}^{n_2} . Let Ω be an open subset of \mathbb{R}^{n_1} . Let F be a C^∞ function from $] -\infty, T] \times \Omega$ to \mathbb{R} such that $F(t, x) = 0$ for all $(t, x) \in] -\infty, 0] \times \Omega$. Then the set*

$$\mathcal{O} := \{\varphi \in C^{1,\alpha}(M, \mathbb{R}^{n_1}) : \varphi(M) \subseteq \Omega\}$$

is open in $C^{1,\alpha}(M, \mathbb{R}^{n_1})$ and the superposition operator T_F of \mathcal{O} to $C_0^{\frac{1+\alpha}{2}; 1+\alpha}([0, T] \times M)$ defined by

$$T_F[\varphi] := F \circ \varphi^T \quad \forall \varphi \in \mathcal{O}$$

is of class C^∞ .

Proof. The set \mathcal{O} is open in $C^{1,\alpha}(M, \mathbb{R}^{n_1})$ because the $C^{1,\alpha}$ -norm is stronger than the norm of the uniform convergence. Let $\{\psi_j\}_{j=1,\dots,r}$ be a local parametrization of class $C^{1,\alpha}$ for M . Lemma A.1 implies that, to prove the lemma, it suffices to show that the operator $C_{\psi_j} \circ T_F$ from \mathcal{O} to $C_0^{\frac{1+\alpha}{2}; 1+\alpha}([0, T] \times \overline{\mathbb{B}_s})$ given by

$$C_{\psi_j} \circ T_F[\varphi] = F \circ (\varphi \circ \psi_j)^T \quad \forall \varphi \in \mathcal{O}$$

is of class C^∞ for all $j = 1, \dots, r$. The map from $C^{1,\alpha}(M, \mathbb{R}^{n_1})$ to $C^{1,\alpha}(\overline{\mathbb{B}_s}, \mathbb{R}^{n_1})$ that takes ϕ to $\phi \circ \psi_j$ is linear and continuous and then of class C^∞ . Accordingly, it suffices to prove that the superposition operator from

$$\mathcal{O}' := \{\phi \in C^{1,\alpha}(\overline{\mathbb{B}_s}, \mathbb{R}^{n_1}) : \phi(\overline{\mathbb{B}_s}) \subseteq \Omega\}$$

to $C_0^{\frac{1+\alpha}{2}; 1+\alpha}([0, T] \times \overline{\mathbb{B}_s})$ that takes ϕ to $F \circ \phi^T$ is of class C^∞ . This fact is a consequence of known results on composition operators (for instance, see Böhme and Tomi [2, p. 10],

Henry [18, p. 29], and Valent [46, Thm 4.4, p. 35]). Indeed, exploiting [46, Thm. 4.4, p. 35] we obtain that the superposition operator that takes ϕ to $F \circ \phi^T$ is of class C^∞ from

$$\mathcal{O}' := \{\phi \in C^{1,\alpha}(\overline{\mathbb{B}_s}, \mathbb{R}^{n_1}) : \phi(\overline{\mathbb{B}_s}) \subseteq \Omega\}$$

to

$$C_0^{1,\alpha}([0, T] \times \overline{\mathbb{B}_s}) := \{f \in C^{1,\alpha}([0, T] \times \overline{\mathbb{B}_s}) : f(0, x) = 0 \ \forall x \in M\}.$$

Finally, to complete the proof we note that the embedding of $C_0^{1,\alpha}([0, T] \times \overline{\mathbb{B}_s})$ into $C_0^{\frac{1+\alpha}{2}; 1+\alpha}([0, T] \times \overline{\mathbb{B}_s})$ is linear and continuous. \square

In our paper we will apply Lemma A.3 to the fundamental solution $S_n(t, x)$ of the heat equation. We observe that $S_n(t, x)$ is real analytic in x for each t fixed, but it is only C^∞ with respect to the pair (t, x) (the problem being for $t = 0$). This fact prevents us from proving a real analytic result for the layer potentials (for example, we can not use Valent [46, Thm. 5.2]).

Acknowledgment

The authors are members of the ‘Gruppo Nazionale per l’Analisi Matematica, la Probabilità e le loro Applicazioni’ (GNAMPA) of the ‘Istituto Nazionale di Alta Matematica’ (INdAM). P.L. acknowledges the support of the Project BIRD191739/19 ‘Sensitivity analysis of partial differential equations in the mathematical theory of electromagnetism’ of the University of Padova.

References

- [1] E. A. Baderko, Parabolic problems and boundary integral equations. *Math. Methods Appl. Sci.* **20** (1997), no. 5, 449–459.
- [2] R. Böhme, F. Tomi, Zur Struktur der Lösungsmenge des Plateuproblems. *Math Z.* **133** (1973), 1–29.
- [3] R. Bramati, M. Dalla Riva, P. Luzzini, and P. Musolino, The Functional Analytic Approach for quasi-periodic boundary value problems for the Helmholtz equation, submitted (2022).
- [4] D. Bucur, G. Buttazzo, Variational methods in shape optimization problems. Progress in Nonlinear Differential Equations and their Applications, 65. Birkhäuser Boston, Inc., Boston, MA, 2005. viii+216 pp.
- [5] R. Chapko, R. Kress, and J. R. Yoon, On the numerical solution of an inverse boundary value problem for the heat equation. *Inverse Problems* **14** (1998), no. 4, 853–867.
- [6] R. Chapko, R. Kress, and J. R. Yoon, An inverse boundary value problem for the heat equation: the Neumann condition. *Inverse Problems* **15** (1999), no. 4, 1033–1046.
- [7] R. R. Coifman and Y. Meyer, Lavrentiev’s curves and conformal mappings, Report No. 5, Institut Mittag-Leffler, 1983.
- [8] M. Dalla Riva and M. Lanza de Cristoforis, A perturbation result for the layer potentials of general second order differential operators with constant coefficients. *J. Appl. Funct. Anal.* **5** (2010), no. 1, 10–30.
- [9] M. Dalla Riva, P. Luzzini, and P. Musolino, Multi-parameter analysis of the obstacle scattering problem, *Inverse Problems* **38** (2022), no. 5, Paper No. 055004, 17 pp.

- [10] M. Dalla Riva, P. Luzzini, and P. Musolino, Shape analyticity and singular perturbations for layer potential operators. *ESAIM Math. Model. Numer. Anal.* 56 (2022), no. 6, 1889–1910.
- [11] M. Dalla Riva, P. Luzzini, P. Musolino, and R. Pukhtaievych, Dependence of effective properties upon regular perturbations. In I. Andrianov, S. Gluzman, V. Mityushev, Editors, *Mechanics and Physics of Structured Media*. pages 271–301, Elsevier, 2022.
- [12] D. Daners, Domain perturbation for linear and nonlinear parabolic equations. *J. Differential Equations* **129** (1996), no. 2, 358–402.
- [13] D. Daners, Domain perturbation for linear and semi-linear boundary value problems, *Handbook of differential equations: stationary partial differential equations*, Vol. VI, 1-81, *Handb. Differ. Equ.*, Elsevier/North-Holland, Amsterdam, 2008.
- [14] K. Deimling, *Nonlinear Functional Analysis*, Springer-Verlag, Berlin, 1985. xiv+450 pp.
- [15] M. C. Delfour and J. P. Zolésio, *Shapes and geometries. Metrics, analysis, differential calculus, and optimization*. Second edition. *Advances in Design and Control*, 22. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2011.
- [16] D. Gilbarg and N. S. Trudinger, *Elliptic partial differential equations of second order*. Elliptic partial differential equations of second order. Reprint of the 1998 edition. *Classics in Mathematics*. Springer-Verlag, Berlin, 2001. xiv+517 pp.
- [17] A. Henrot and M. Pierre, *Variation et optimisation de formes*, Vol. 48 of *Mathématiques & Applications (Berlin) [Mathematics & Applications]*, Springer, Berlin, 2005. xii+334 pp.
- [18] D. Henry, *Topics in nonlinear analysis*, Universidade de Brasilia, *Trabalho de Matemática* **192** (1982).
- [19] F. Henríquez and C. Schwab, Shape holomorphy of the Calderón projector for the Laplacian in \mathbb{R}^2 . *Integral Equations Operator Theory* 93 (2021), no. 4, Paper No. 43, 40 pp.
- [20] E. Hille and R.S. Phillips. *Functional analysis and semi-groups*. American Mathematical Society Colloquium Publications, vol. 31. American Mathematical Society, Providence, R.I., 1957. rev. ed.
- [21] O. A. Ladyženskaja, V. A. Solonnikov, and N. N. Ural’ceva, *Linear and quasi-linear equations of parabolic type*. (Russian) Translated from the Russian by S. Smith. *Translations of Mathematical Monographs*, 23 American Mathematical Society, Providence, R.I. 1968.
- [22] M. Lanza de Cristoforis, Properties and pathologies of the composition and inversion operators in Schauder spaces, *Accad. Naz. delle Sci. detta dei XL* **15** (1991), 93–109.
- [23] M. Lanza de Cristoforis, A domain perturbation problem for the Poisson equation. *Complex Var. Theory Appl.* **50** (2005), no. 7-11, 851–867.
- [24] M. Lanza de Cristoforis, Perturbation problems in potential theory, a functional analytic approach, *J. Appl. Funct. Anal.* **2** (2007), no. 3, 197–222.
- [25] M. Lanza de Cristoforis and P. Luzzini, Time dependent boundary norms for kernels and regularizing properties of the double layer heat potential. *Eurasian Math. J.* **8** (2017), no. 1, 76–118.
- [26] M. Lanza de Cristoforis and P. Luzzini, Tangential derivatives and higher order regularizing properties of the double layer heat potential, *Analysis (Berlin)* **8** (2019), no. 4, 167–193.

- [27] M. Lanza de Cristoforis and P. Musolino, A perturbation result for periodic layer potentials of general second order differential operators with constant coefficients. *Far East J. Math. Sci.* **52** (2011), no. 1, 75–120.
- [28] M. Lanza de Cristoforis and L. Preciso, On the analyticity of the Cauchy integral in Schauder spaces, *J. Integral Equations Appl.* **11** (1999), no. 3, 363–391.
- [29] M. Lanza de Cristoforis and L. Rossi, Real analytic dependence of simple and double layer potentials upon perturbation of the support and of the density, *J. Integral Equations Appl.* **16** (2004), no. 2, 137–174.
- [30] M. Lanza de Cristoforis and L. Rossi, Real analytic dependence of simple and double layer potentials for the Helmholtz equation upon perturbation of the support and of the density, in: *Analytic methods of analysis and differential equations: AMADE 2006*, Camb. Sci. Publ., Cambridge, 2008, pp. 193–220.
- [31] G. M. Lieberman, *Second order parabolic differential equations*. World Scientific Publishing Co., Inc., River Edge, NJ, 1996. xii+439 pp.
- [32] A. Lunardi and V. Vespri, Hölder regularity in variational parabolic non-homogeneous equations, *J. Differential Equations* **94** (1991) no. 1, 1–40.
- [33] P. Luzzini, Regularizing properties of space-periodic layer heat potentials and applications to boundary value problems in periodic domains. *Math. Methods Appl. Sci.* **43** (2020), no. 8, 5273–5294.
- [34] P. Luzzini, Regularizing properties of the double layer heat potential and shape analysis of a periodic problem. *Ph.D. Dissertation*, Università degli Studi di Padova (2019).
- [35] P. Luzzini, A mapping property of the heat volume potential, submitted (2022).
- [36] P. Luzzini and P. Musolino, Perturbation analysis of the effective conductivity of a periodic composite. *Netw. Heterog. Media* **15** (2020), no. 4, 581–603.
- [37] P. Luzzini, P. Musolino, and R. Pukhtaievych, Shape analysis of the longitudinal flow along a periodic array of cylinders. *J. Math. Anal. Appl.* **477** (2019), no. 2, 1369–1395.
- [38] A.A. Novotny and J. Sokolowski, *Topological derivatives in shape optimization, Interaction of Mechanics and Mathematics*, Springer, Heidelberg, 2013. xxii+412 pp.
- [39] O. Pironneau, *Optimal shape design for elliptic systems*. Springer Series in Computational Physics. Springer-Verlag, New York, 1984. xii+168 pp.
- [40] R. Potthast, Domain derivatives in electromagnetic scattering, *Math. Methods Appl. Sci.* **19** (1996), no. 15, 1157–1175.
- [41] R. Potthast, Fréchet differentiability of boundary integral operators in inverse acoustic scattering, *Inverse Problems* **10** (1994), no. 2, 431–447.
- [42] R. Potthast, Fréchet differentiability of the solution to the acoustic Neumann scattering problem with respect to the domain, *J. Inverse Ill-Posed Probl.* **4** (1996), no. 1, 67–84.
- [43] R. Potthast and I. G. Stratis, On the domain derivative for scattering by impenetrable obstacles in chiral media. *IMA J. Appl. Math.* **68** (2003), no. 6, 621–635.
- [44] G. Prodi and A. Ambrosetti. *Analisi non lineare. I quaderno*. Pisa: Scuola Normale Superiore Pisa, Classe di Scienze, 1973.
- [45] J. Sokolowski and J.P. Zolésio, *Shape sensitivity analysis*. Springer Series in Computational Mathematics, 16. Springer-Verlag, Berlin, 1992. ii+250 pp.

- [46] T. Valent, Boundary value problems of finite elasticity. Local theorems on existence, uniqueness, and analytic dependence on data. Springer Tracts in Natural Philosophy, 31. Springer-Verlag, New York, 1988. xii+191 pp.
- [47] G. Verchota, Layer potentials and regularity for the Dirichlet problem for Laplace's equation in Lipschitz domains. *J. Funct. Anal.* **59** (1984), no. 3, 572-611.
- [48] S. Wu, Analytic dependence of Riemann mappings for bounded domains and minimal surfaces, *Comm. Pure Appl. Math.* **46** (1993), no. 10, 1303–1326.